

Epidemiological-clinical, Therapeutic and Evolutionary Profile of Children Treated for Bacterial Meningitis at CHU-MEL (Benin)

Annatou Yakoubou ^{a*}, Linda Ayedadjou ^a, Nicole Enianloko Tchiakpe ^a,
Omoyélé Ladipo ^a, Eurydice Yerima ^a, Adolphe Adjanohoun ^b, Haziz Sina ^b
and Lamine Baba-Moussa ^b

^a *Pediatric Unit, Hospital and University Center for Mothers and Children Lagune (CHU-MEL) of Cotonou, Benin.*

^b *Department of Biochemistry and Cell Biology, Laboratory of Biology and Molecular Typing in Microbiology, University of Abomey-Calavi, Benin.*

Authors' contributions

This work was carried out in collaboration among all authors. Authors AY, LA, NET and OL designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors AY, LA, OL, AA, HS and LBM managed the analyses of the study. Authors AY, NET, EY and HS managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJPR/2022/v9i430272

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/91075>

Original Research Article

Received 09 July 2022
Accepted 03 September 2022
Published 08 September 2022

ABSTRACT

Aims: Bacterial meningitis remains a public health problem due to an unfavorable geographical environment and socioeconomic context. Our study aimed to evaluate the epidemiological-clinical, therapeutic and evolutionary profile of children treated for bacterial meningitis in the pediatric department of CHU-MEL in Benin.

Methodology: The methodology consisted in carrying out a retrospective and descriptive study of children admitted to the pediatric department of the CHU-MEL in Cotonou from January 2019 to December 2020. All children admitted in the study period were taken in consideration. Data collection (epidemiological, clinical, paraclinical, therapeutic and evolutionary) was made from registers and files and by telephone call.

Results: Our data reveal that 68 children were treated for bacterial meningitis between January

*Corresponding author: Email: annatouy@hotmail.com;

2019 to December 2020. It was also noted that the most represented age group was between 3 months and 3 years with a rate of 52.9%. Most (95.6%) of the children had no particular history. The antibiotics used were ampicillin (17.4%), ceftriaxone (52.2%), cefotaxime (4.3%) and ciprofloxacin (8.7%). For the exploration of cases of meningitis, apart from the cytobacteriological examination of the cerebrospinal fluid (CSF), the blood culture, the complete blood count (NFS) and the C-reactive protein (CRP) were the examinations commonly requested.

Conclusion: It is observed that children treated as meningitis have varying clinical signs. Meningitis cases have different ends. In case of bacterial meningitis, it is important to determine the germs to ensure proper antibiotic therapy to be conducted.

Keywords: Bacterial meningitis; antibiotics; management; pediatric; Benin.

1. INTRODUCTION

Bacterial meningitis (BM) is a major pediatric pathology and constitutes a public health problem due to the high mortality associated with it [1]. It causes a diagnostic and therapeutic emergency if not properly diagnosed and managed [2]. Thus, the annually worldwide occurrence of BM estimated at over 1.2 million cases [3] though the incidence and case-fatality rates vary by region, country, pathogen and age group [4]. The clinical signs of the BM are not specific but among untreated patients, more than 70% induce case-fatality and 20% of survivors have permanent sequelae [4]. Therefore, this infection is considered as one of the most feared childhood diseases with special recommendations for its detection in endemic countries in Africa [5-6].

The detection of bacterial meningitis is often difficult [7]. This is why several tools have been developed and validated to guide decision-making. These tools will help to limit the prolonged and hazardous use of antibiotics [7-9]. The commonest bacterial meningitis pathogens are mainly *Neisseria meningitidis*, *Streptococcus pneumoniae* and *Haemophilus influenzae* [3]. But the incidence of *N. meningitidis* is highest in sub-Saharan Africa region during the dry season [5,10].

In Benin, as in many other sub-Saharan countries, meningitis suspected children are routinely admitted to hospitals and treated with broad-spectrum antibiotics without any laboratory confirmation. But very few studies have looked into this disease in Benin. To fill this gap, the present study is undertaken with the aim of appreciating not only the epidemiological-clinical profile but also those therapeutic and evolutionary of the children taken care of as bacterial meningitis in the pediatric department of the CHU-MEL in Cotonou (Benin).

2. METHODOLOGY

2.1 Type and Period of Study

This was a retrospective and descriptive study of children admitted to the pediatric department of the CHU-MEL of Cotonou from January 2019 to December 2020 and treated for bacterial meningitis. Included in the study were all patients aged 1 month to 17 years in whom the diagnosis of bacterial meningitis was retained and who underwent a lumbar puncture. The non-inclusion criterion was the non-availability of a cytobacteriological and chemical examination of the CSF. Recruitment was exhaustive and all patients meeting the inclusion criteria were included.

2.2 Data Collection

Data were collected first from registers and files and then by telephone call. Consulting the registers, the collected were epidemiological, clinical, paraclinical, therapeutic and evolutionary. The epidemiological data related to age, sex, treatment before admission and pathological history. With regard to the clinical data, they focused on the presence of fever, neurological signs and signs of meningeal irritation. The assessments made with their results were the substance of the paraclinical data. The therapeutic data took into account the antibiotics used, the duration of the treatments and the poorly followed treatment. We define poorly followed treatment as non-compliance with the dose and/or frequency and/or duration of the prescribed antibiotic and/or the existence of a therapeutic window. Finally, the evolutionary data took into account the short-term evolution, the mode of discharge from the hospital and the medium-term evolution (1 year after hospitalization).

2.3 Data analysis and Processing

Data collected was process using Microsoft Excel 2013 software and the IBM SPSS statistics 26 statistical analysis tool. The qualitative variables were expressed as a frequency. Correlations between variables were tested by the chi-square test. We considered a p value < 0.05 to be statistically significant.

3. RESULTS AND DISCUSSION

3.1 Results

3.1.1 Population size

During the study period, 68 children were treated for bacterial meningitis with a cyto-bacteriological and biochemical examination of the CSF out of a total of 10,236 hospitalized children, i.e., a hospital frequency of 0.66%.

3.1.2 Epidemiological profile

The average age was 31.6 months with extremes of 1 month and 13 years. The most represented age group was between 3 months and 3 years with a rate of 52.9% (Table 1). The predominance is male with a sex ratio of 1.1.

3.1.3 Clinical profile

Regarding the history, the majority (95.6%) of the children had no particular history. Nevertheless, antibiotic therapy was revealed by 33.8% (23/68) of the children. The antibiotics used were ampicillin (17.4%), ceftriaxone (52.2%), cefotaxime (4.3%) and ciprofloxacin (8.7%).

On admission, the diagnosis of bacterial meningitis was suggested by the deterioration of the general condition, the infectious syndrome, the signs of meningeal irritation and/or the neurological signs. Neurological signs were convulsions or coma rarely incessant crying or bulging anterior fontanel in infants. Table 2 shows the distribution of children treated for bacterial meningitis according to the clinical signs presented.

3.1.4 Reports requested and executed

Among the 68 cases reported during our investigation the rate of children's that performed the specific examinations varies (Table 3). Thus, for the confirmation of meningitis cases, apart from the classic cyto-bacteriological cerebrospinal fluid (CSF) examination, the blood culture, complete blood count (NFS) and C-reactive protein (CRP) were the examinations commonly requested. In Table 3, we can observe that the complete blood count was the most performed requested examination (95.6%) followed by the blood culture (22.1%) and then C-reactive protein (7.4%). It should be noted that in the case of blood culture, about 20% (3/15) of the analyzes samples were positive. The germs found were *Haemophilus influenzae*, *Escherichia coli* and a Gram-negative bacillus.

Most of the children had cellulorachy less than 100 cells/mm³ (55.9%, n=38) high protein ≥ 0.40 g/l (91.2%; n=62) and high glycorrachia greater than 0.27g/l (82.3%; n=56). Table 4 presents the distribution of children treated as bacterial meningitis according to the number of leukocytes, proteinorachia and glycorrachia.

Table 1. Distribution of the population studied according to age

Age	Number	Percentage (100%)
≤ 3-months	12	17,7
]3-month, 3 years]	36	52.9
]3-years, 5 years]	10	14.7
>5-years	10	14.7
Total	68	100.0

Table 2. Distribution of children treated as bacterial meningitis according to the clinical signs presented

Signs	Number	Percentage (%)
Fever	68	100
Signs of meningeal irritation	6	9.7
Neurological signs	61	89.7

Table 3. Distribution of children treated for bacterial meningitis according to the requested examinations

Reports carried out	Number	Percentage (%)
Blood culture	15	22.06
Complete blood count (NFS)	65	95.59
C-reactive protein (CRP)	5	7.35

Table 4. Distribution of children treated as bacterial meningitis according to the number of leukocytes, proteinorachia and glycorrachia

		Number	Percentage (%)
Cellulorachy	[10 cells / mm ³ – 50 cells / mm ³]	31	45.59%
	[50 cells/mm ³ – 100 cells / mm ³]	7	10.30%
	[100 cells/mm ³ – 500 cells/mm ³]	22	32.35%
	[500 cells/mm ³ – 1000 cells /mm ³]	2	2.94%
	≥ 1000 cells / mm ³	6	8.82%
Total		68	100.00%
Proteinorachia	<0.40 g/l	6	8.82%
	≥ 0.40 g/l	62	91.18%
	Total	68	100.00%
Glycorrachia	< 0.27 g/l	12	17.65%
	≥ 0.27 g/l	56	82.35%
	Total	68	100.00%

3.1.5 Therapeutic profile

The preferred treatment in the department was cefotaxime at 200 mg/kg/24 h three times plus gentamicin at 5 mg/kg/24 h once. In the absence of financial means from the parents, ceftriaxone 100 mg/kg/24 h twice is proposed then the treatment is adapted according to the germ found. In the absence of germs, ceftriaxone or cefotaxime is continued for ten days and gentamicin for 5 days. Finally, in the absence of

improvement after ten days of treatment, other antibiotics are proposed. Table 5 shows the combinations of antibiotics made in hospitalization.

The combinations of antibiotics used were diverse and varied. Overall, treatment was poorly followed by 60.3% (41/68) of children treated as bacterial meningitis. Most of the children (69.1%; n=47) completed 10 days of treatment (minimum treatment time for meningitis in the service).

Table 5. Distribution of children treated for meningitis according to the antibiotics used

Antibiotics	Number	Percentage (%)
Ceftriaxone + gentamicin	17	25.00%
Cefotaxime + gentamicin	33	48.50%
Ceftriaxone sulbactam + gentamicin	2	2.90%
Ceftriaxone + gentamicin then vancomycin	1	1.50%
Cefotaxime + gentamicin + ampicillin	2	2.90%
Cefotaxime then ceftriaxone + gentamicin then ciprofloxacin	2	2.90%
Cefotaxime + gentamicin then ciprofloxacin	4	5.90%
Cefotaxime + gentamicin then ciprofloxacin + CTM + phenicols	4	5.90%
Cefotaxime + gentamicin then ceftriaxone	1	1.50%
Cefotaxime + gentamicin then Ceftriaxone sulbactam + vancomycin	1	1.50%
Cefotaxime + gentamicin then ciprofloxacin + ampicillin	1	1.50%
Total	68	100.00%

3.1.6 Scalable profile

3.1.6.1 Short-term development

Table 6 shows the distribution of children treated for meningitis according to the evolution of clinical signs. It is observed that children treated as meningitis have varying clinical signs. We can note a predominance of children who have a persistence of fever beyond 5 days (33.3%).

Fig. 1 shows the mode of discharge of children treated as bacterial meningitis at CHU-MEL. An intra-hospital lethality of 7.3% (5/68) is noted. Uncured children are children who returned against medical advice or who did not benefit from a lumbar puncture control before the execution.

Fig. 2 shows the duration of hospitalization of children treated for bacterial meningitis. The majority of children (83.8%) spent more than 10 days in hospital with an average hospital stay of 18 days.

3.1.6.2 Medium-term development

Fig. 3 presents the medium-term evolution (1 year after hospitalization) of children treated

for bacterial meningitis. In this case, we see that 51 of the 68 parents were reachable for the assessment of the medium-term development of the children. Of these parents contacted, the majority expressed a favorable evolution of their child.

Table 7 below presents the evolutionary modalities of the children who had an unfavorable evolution. The majority (37.5%) had presented a regression of psychomotor acquisitions.

The overall lethality was 12.5% (two children died after exeat: the first during an intervention for hydrocephalus and the second in a context of altered consciousness after exeat). The sequelae are in the order of 8.9% (5/56). In total, 23.2% (13/56) of children treated for presumed bacterial meningitis experienced an unfavorable medium-term outcome.

The factors associated with an unfavorable evolution of children treated as bacterial meningitis are presented in Table 8.

Table 6. Distribution of children treated for meningitis according to the evolution of clinical signs

	Number	Percentage (%)	Total
Number of children with persistent fever for more than 5 days	22	33.3	66
Number of children who had persistent signs of meningeal irritation beyond 5 days	1	16.7	6
Number of children with persistence of neurological signs beyond 5 days	9	15.2	59

NB: Two children died before the fifth day




Fig. 1. Mode of discharge of children treated as bacterial meningitis at CHU-MEL




Fig. 2. Duration of hospitalization of children treated as bacterial meningitis




Fig. 3. Medium-term evolution of children treated for bacterial meningitis

Table 7. Evolutionary modalities of children who had an unfavorable evolution

Unfavorable development	Number	Percentage (%)
Death after enforcement	2	25.0
Hydrocephalus	1	12.5
Recurrent bacterial meningitis	1	12.5
Regression of psychomotor acquisitions	3	37.5
Deafness + blindness	1	12.5
Total	8	100

Table 8. Factors associated with the unfavorable evolution of children treated for bacterial meningitis

Risk factors		Favorable evolution	Unfavorable evolution	p
Age (n=56)	0- 3 months	7	1	0.197
	[3 months – 3 years]	21	10	
	[3 years -5 years]	6	2	
	> 5 years	9	0	
Cellulorachia (n=56)	0-50	25	3	0.083
	50-100	3	2	
	≥ 100	15	8	
Proteinorachia (n=56)	< 0.40	4	1	0.858
	≥ 0.40	39	12	

Risk factors		Favorable evolution	Unfavorable evolution	p
Glycorachia (n=56)	≤ 0.27	4	5	0.012
	> 0.27	39	8	
Poorly followed treatment (n=56)	Yes	21	13	0.001
	No	22	0	
Duration of treatment (n=56)	10 days	33	6	0.094
	21 days	6	5	
	> 21 days	4	2	
Disappearance of fever (n=51)	< 5 days	33	2	0.004
	≥ 5 days	10	6	
Disappearance of neurological signs (n=46)	< 5 days	35	4	0.027
	≥ 5 days	4	3	
Mode of exit (n=51)	Healed	37	2	0.000
	Not Healed	6	6	

3.2 Discussion

3.2.1 Epidemiological characteristics

In our study, the hospital frequency of meningitis treated as bacterial was 0.66%. This prevalence is lower than the 1.07% found at the CHU Gabriel Touré in Mali by Maïga et al. [11] and the 1.4% reported by Kané et al. [12] in their study conducted in the pediatric department of the Mali hospital from 2012 to 2018. However, this value is higher than the 0.3% found by Tiftha et al. [13]. In this last study, only 61 cases of bacterial meningitis had been recruited at Sahloul University Hospital (Sousse-Tunisia) over a period of 11 years (2006 to 2016). Thus, the most represented age group is between 3 months and 3 years (52.9%) and the majority (85.3%) were under 5 years old. This observation corroborates that found by Maïga et al. [11] when they found a similar prevalence to ours; which shows the immune fragility of this age group.

3.2.2 Biological characteristics

Regarding the biological characteristics, in our series, a germ was isolated in the CSF in 4.4% of cases. Previous studies have reported much higher values with positivity thresholds of 39% in Mali [11], 66% in Morocco [14] and 67.2 in Tunisia [13]. The absence of germs in the Cerebrospinal Fluid (CSF) despite bacterial meningitis may be related to taking antibiotics before performing the lumbar puncture. But only 33.8% of the children in our study had received an antibiotic compared to the series by Maïga et al. [11] of which 66% of children had received antibiotic therapy before admission. CSF and PCR can be very helpful in the diagnosis of bacterial meningitis. Along the same lines, Kané et al. [12], in their study on the diagnosis of

meningitis in children used PCR (59.1%), Gram stain (39.8%), agglutination test (29.5%) and cultivation (5.7%). The absence of germs in the CSF in the event of bacterial meningitis, perhaps also linked to the conditions of transport and storage. Indeed, given the fragility of certain germs such as pneumococcus, CSF must reach the laboratory without delay at room temperature for immediate processing. The study of transport and storage conditions and the performance of PCR on CSF samples would help us at the CHU-MEL to better detect the germs involved in presumed bacterial meningitis. Additionally, in low-resource settings, many children may present with bacterial meningitis, but laboratory capacity may not be sufficient to isolate the causative agent and confirm the diagnosis [15]. Unless this is an overestimate of cases of bacterial meningitis by clinicians, because most of the children in our series had a cellularity less than 100 cells/mm³ (55.9%, n=38) and a higher high glycorrhachia to 0.27g/l (82.3%; n=56) or bacterial meningitis is often considered when the leukocytes are greater than 100 cells/mm³ and the glycorrhachia is less than 0.27g/l [16].

In our study, 7.4% of children performed CRP; 22.1% blood culture and 95.6% NFS. These assessments are not systematically carried out because they are the responsibility of the parents and their realization depends on their financial means. However, these assessments and especially the blood procalcitonin and the lactate level in the CSF have an orientation interest in the diagnosis of bacterial meningitis. In the case of a CSF lactate assay, a value below 3.2 mmol/l makes the diagnosis of bacterial meningitis very unlikely and a value below 0.5 ng/ml for the procalcitonin assay serum level makes the diagnosis of bacterial meningitis highly unlikely

[17]. Blood procalcitonin is considered to have higher diagnostic accuracy for detecting bacterial meningitis than serum C-reactive protein, leukocyte count, neutrophil count, CSF protein and glucose levels [18-19]. Often the combination of the results of these tests allows an accurate prediction of the likelihood of bacterial meningitis versus viral meningitis.

3.2.3 Therapeutic characteristics

In the case of bacterial meningitis, antibiotic therapy should ideally be initiated within one hour of arrival at the hospital, regardless of the time that has already elapsed since the presumed start of the meningitis [20]. In our investigation, all our patients received ceftriaxone or cefotaxime either in combination with gentamicin or other antibiotics. This corresponds to the recommendations for the management of bacterial meningitis, which recommend the use of a third-generation cephalosporin in the event of bacterial meningitis [21]. Unfortunately, 60.3% of the children treated for bacterial meningitis in our study did not follow the treatment well. This poor compliance is probably linked to the cost of antibiotics, the duration of treatment and the low purchasing power of populations. Poorly followed treatment is statistically associated with the development of children treated as bacterial meningitis ($p = 0.001$). Health insurance for all would be a great contribution to overcoming the therapeutic non-compliance of our patients. In addition, the combinations of antibiotics used are diverse and varied. It would be more appropriate to establish a protocol of antibiotics to be used in the service in the event of bacterial meningitis.

3.2.4 Scalable features

In the event of meningitis, surveillance focuses on fever and neurological signs. The fever often disappears after 48 hours and the neurological signs within 2 to 5 days [20]. About 1/3 of the children (33.3%) had a persistence of fever beyond 5 days and this is associated with an unfavorable outcome ($p=0.004$). Similarly, 15.2% of children had persistence of neurological signs beyond 5 days and this was also associated with an unfavorable outcome ($p=0.027$). The medium-term evolution was assessed after phone calls to parents, this underlines a lack of follow-up of children hospitalized for bacterial meningitis. In addition, sequelae were observed in 8.9% of children (5/56). These sequelae were of the hydrocephalus type (1.8%), regression of psychomotor acquisitions (5.3%) and deafness +

blindness (1.8%). In addition, one child had a recurrence of bacterial meningitis (1.8%). The sequelae rate is lower than the 13% reported by El Fakiri et al. [14] and 12.5% by Kané et al. [12]. The overall lethality was 12.5% (7/56). This rate is similar to that of Merabet et al. [22] who recorded a case fatality rate of 11.7% in the Tangier region of Morocco. Nevertheless, this rate is lower than that found in the pediatric department of the Mali Hospital where it was 18.2% [12] and that found by El Amrani et al. [23] who found a rate of 19.2% but it is on the other hand much higher than that found by El Fakiri et al. [14] which was 1%.

Despite the limitations of our study with regard to the epidemiological data obtained from medical records, which did not allow us to study the socio-economic level of the parents and the vaccination status of the children and the evolution in the medium term which could not be appreciated in all the children (12 parents of children were unreachable by telephone), our study tried to objectively assess the state of play on the management of supposedly bacterial meningitis in our center while indicating some areas for improvement.

4. CONCLUSION

Bacterial meningitis is a serious condition characterized by high mortality and significant sequelae. In our study, 23.2% of children treated for presumed bacterial meningitis experienced an unfavorable evolution with an overall lethality of 12.5% and sequelae in the order of 8.9%. To improve care, it is important to better determine the germs involved, to establish a care protocol and to ensure proper conduct of antibiotic therapy.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Saez-Llorens X, McCracken GH. Bacterial meningitis in children. *Lancet*. 2003;361:2139–2148.
2. Nudelman Y, Tunkel A. Bacterial meningitis: Epidemiology, pathogenesis and management update. *Drugs*. 2009;69(18):2577–2596.
3. World Health Organization and Office of Information. Epidemic meningococcal

- disease. WHO fact sheet. Geneva: World Health Organization; 1998
4. Rosenstein N. Meningococcal disease. *N Engl J Med.* 2001;344(18):1378–1388.
 5. World Health Organization and Office of Information. Recommendation of a consensus meeting on detection of meningitis epidemics in Africa. Paris: World Health Organization; 2000.
 6. World Health Organization. Detecting meningococcal meningitis epidemics in highly-endemic African countries: WHO recommendation. *Weekly Epidemiological Record= Relevé épidémiologique hebdomadaire.* 2000;75(38):306-309.
 7. Nigrovic L, Kuppermann N, Malley R. Development and validation of a multivariable predictive model to distinguish bacterial from aseptic meningitis in children in the post-Haemophilus influenzae era. *Pediatrics.* 2002;110(4):712–719.
 8. Hart A, Hopkins CA. 2001 ICD-9-CM Code Book. Reston: St. Anthony Publishing and West Valley City, UT: Ingenix; 2000.
 9. Nigrovic L. Clinical prediction rule for identifying children with cerebrospinal fluid pleocytosis at very low risk of bacterial meningitis. *JAMA.* 2007;297(1):52–60.
 10. World Health Organization. Managing meningitis epidemics in Africa a quick reference guide for health authorities and health-care workers. Geneva: World Health Organization; 2010.
 11. Maïga B, Sacko K, Diakité F, Dembélé A, Dicko Traoré F, Diakité A, Traoré F, Diall H, Touré A, Cissé M, Togo P, Doumbia A, Sanogo T, Coulibaly O, Traoré I, Coulibaly A, Konaté D, Diakité F, Doumbia A, Maïga L, Konaré H, Sylla M, Togo B. Méningites Bactériennes chez l'Enfant au Service de Pédiatrie du CHU Gabriel Toure. *Health Sci Dis.* 2019;20(4). Available: <http://hsd-fmsb.org/index.php/hsd/article/view/1515>
 12. Kané B, Abdou M, Koné O, Dembélé G, Wélé Diallo K, Fané B, Sangaré A, Coulibaly M, Togo B. Causes of bacterial meningitis in children from 1 month to 15 years in the pediatric department of the hospital in Mali from 2012 to 2018. *Rev Mali Infect Microbiol.* 2020;15:72-76
 13. Tifha M, Mallouli M, Sahli J, Ben Abed H, Chemli J, Zouari N, Mabrouk S, Ajmi H, Hassayoun S, Abroug S. Bacterial Meningitis *In* Infants And Children: 11-Year Report In A Tunisian Pediatric Tertiary Unit. *J I M Sfax.* 2018;29:54–63.
 14. El Fakiri K, Bourrous M, Diffo C, Rada N, Draïss G, Bouskraoui M. Les méningites du nourrisson et de l'enfant au centre hospitalier universitaire de Marrakech : Expérience d'une unité pédiatrique marocaine. *J Pédiatrie Puériculture.* 2016; 29(5):237-243 Available: <https://doi.org/10.1016/j.jpp.2016.08.002>
 15. Luksic I, Mulic R, Falconer R, Orban M, Sidhu S, Rudan I. Estimating global and regional morbidity from acute bacterial meningitis in children: assessment of the evidence. *Croatian Med J.* 2013;54(6): 510-518. Available: [10.3325/cmj.2013.54.510](https://doi.org/10.3325/cmj.2013.54.510)
 16. Organisation Mondiale de la Santé. Guide to pediatric hospital care: Management of common childhood ailments. WHO. 2015;540. Available: https://apps.who.int/iris/bitstream/handle/10665/187940/9789242548372_fre.pdf
 17. Société de pathologie infectieuse de langue française. 17th consensus conference. Consensus conference on bacterial meningitis. (Short text). *Med Mal Inf.* 2009 ;39(3):175-186. French. DOI: [10.1016/j.medmal.2008.12.001](https://doi.org/10.1016/j.medmal.2008.12.001)
 18. Kim H, Roh YH, Yoon SH. Blood procalcitonin level as a Diagnostic marker of Pediatric Bacterial Meningitis: A Systematic Review and Meta-Analysis. *Diagnostics (Basel).* 2021;11(5):846. DOI: [10.3390/diagnostics11050846](https://doi.org/10.3390/diagnostics11050846)
 19. Babenko D, Seidullayeva A, Bayesheva D, Turdalina B, Omarkulov B, Almabayeva A, Zhanaliyeva M, Kushugulova A, Kozhakhmetov S. Ability of procalcitonin and c-reactive protein for discriminating between bacterial and enteroviral meningitis in children using decision tree, *BioMed Res Int.* 2021;Article ID 5519436:7. Available: <https://doi.org/10.1155/2021/5519436>
 20. Bourrillon A, Aujard Y, Bingen E. Méningites purulentes du nouveau-né, du nourrisson, et de l'enfant. *EMC Pédiatrie et Maladies Infect.* 2006;4-210-B-10
 21. Tunkel AR, Hartman BJ, Kaplan SL, Kaufman BA, Roos KL, Scheld WM, Whitley RJ. Practice Guidelines for the Management of Bacterial Meningitis, *Clin Infect Dis.* 2004;39(9):1267–1284

- Available: <https://doi.org/10.1086/425368>
22. Merabet M, Aouragh R, Idrissi A. Les méningites bactériennes aiguës communautaires chez les enfants de moins de 5 ans à la région Tanger-Tétouan-Al Hoceima (Maroc) 2006-2015 : Profil épidémiologique, clinique et biologique. *Antropo*. 2018;40:1-11.
- Available: www.didac.ehu.es/antropo
23. El Amrani K, El Hafidi N, Barkia A, Jroundi I. Epidemiological, clinical characteristics and prognostic factors of bacterial meningitis in children admitted at the pediatric hospital of Rabat, Morocco. *Rev Marocaine Santé Publique*. 2016;3(5): 11-18.

© 2022 Yakoubou et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/91075>