



Effect of Micronutrients on Physico-Chemical Parameters of Fruit and Yield of Guava (*Psidium guajava* L.) cv. L-49

Monika Kumari Meena^{a*} and R. S. Verma^{b#}

^a Department of Horticulture, School for Agricultural Sciences & Technology, Babasaheb Bhimrao Ambedkar University, Lucknow-226025, India.

^b Department of Fruit Science, School for Agricultural Sciences & Technology, Babasaheb Bhimrao Ambedkar University, Lucknow-226025, India.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2022/v34i1731031

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/86097>

Original Research Article

Received 14 February 2022

Accepted 23 April 2022

Published 28 April 2022

ABSTRACT

This research was done to find out the effect of micronutrients on physico-chemical parameters of fruit and yield of guava (*Psidium guajava* L.). For this purpose, guava var. L-49 was used in this research. The experiment consisted of eight treatment combinations having Zn, Mn and Fe concentrations of 0.5, 0.5 and 0.4 % respectively, in Randomized Block Design with three replications. There was a significant effect of micronutrients (Zn, Mn and Fe). According to the results, significant increases of fruit size (cm), fruit weight (gm), volume of fruit (cm³), specific gravity and chemical parameters are Total soluble solids (TSS °Brix), acidity (%), ascorbic acid (Vitamin 'C') (mg/100gm pulp), of fruit were determined with increasing micronutrients application. This experimental research carried out at Horticulture Research Farm, Department of Horticulture, Babasaheb Bhimrao Ambedkar University, Vidya-Vihar, Rae Bareilly Road, Lucknow 226 025 (U.P.) during the summer season of 2019-20.

Keywords: Guava; micronutrients; physico-chemical parameters; yield; Zn; Mn and Fe.

[≡]M.Sc. (Fruit Science);

[#]Assistant Professor;

^{*}Corresponding author: E-mail: monikahorticos@gmail.com;

1. INTRODUCTION

Guava (*Psidium guajava* L.) "Apple of the tropics" is an important fruit crop of the India, not because of its large area and production but due to its wider edapho-climatic adaptability, hardly to various biotic and abiotic stresses, precocious and prolific bearing habit, quality fruit with high nutritive value, medicinal attribute, use both as fresh fruit and after processing in different value-added products and considered as a multipurpose tree due to its utility as fruit, fuel, fodder, timber and it is a highly remunerative crop. It is a very rich source of vitamin C and A along with minerals like iron, calcium, and phosphorus. Guava is fourth the most important fruit crop in India after Mango, Banana and Citrus (Ray 2002). It also contains substantial quantities of carbohydrates, sugars and pectin. Owing to excellent taste and flavour, high nutritional value and wide availability at a moderate price the fruit is often called "Poor man's apple". The conventional products of guava are jelly, jam, pulp, concentrate, juice, cheese, toffee, dehydrated guava and canned guava. Subhash Chandra [1].

Guava is commercially grown throughout the India, particularly in Maharashtra, Uttar Pradesh, Bihar, Orissa, Punjab, Uttarakhand, Gujarat, Madhya Pradesh, and West Bengal. Uttar Pradesh is considered the most important guava-producing state of India, in which Allahabad (Prayagraj) region has the reputation of growing the best quality guava in the country as well as all over the world. It is cover around 3.7% (2.7 lakh ha) of the total area under fruit crops and contributes 3.3% (41.07 MT) of total fruit production (NHB 2017-18).

The spring flowering is called "Ambe Bahar", June or monsoon flowering is called "Mrig Bahar" and the third flowering which comes in October is called "Hast Bahar". Ambe Bahar fruits ripen from July to September and Mrig Bahar fruits ripen from November to February however, Hast Bahar fruits ripen in the spring season which is also known as the summer season [2-4].

The fruit is an excellent source of vitamin C (210 mg/100g) and pectin (0.60 %) but has low energy (66 cal /100g), protein content (1%) and dry matter (17%) and moisture (83%). The fruit is also rich in minerals like phosphorus (24-37mg/100g), calcium (14-30 mg/100g) and iron (0.6-1.4mg/100g) as well as vitamins like niacin, pantothenic acid, thiamine, riboflavin and vitamin A [5].

Foliar application is based on the principle that the nutrients are quickly absorbed by leaves and transported to different parts of the plant to the functional requirement of nutrition. This method is highly helpful for the correction of element deficiencies to restore disrupted nutrient supply, and overcome stress factors limiting their availability and it plays important role in improving fruit set, productivity and quality of fruits and recovery of nutritional and physiological disorders in fruit trees.

Zinc is an important constituent of several enzyme systems which regulate various metabolic reactions associated with water relations in the plant. Zinc is essential for auxin and protein synthesis, seed production and proper maturity. It also increases fruit size as well as yield. Zinc is essential for improving the vegetative growth of guava trees obtained in terms of terminal shoots, shoot diameter and number of leaves per shoot [6]. Boron is a constituent of cell membrane and essential for cell division. It acts as a regulator of potassium/calcium ratio in the plant and helps in nitrogen absorption and translocation of sugar in plant. Boron increases nitrogen availability to plants. It is involved in the synthesis of cell wall components. It has a central role in pollen viability and good fruit set. It increases the elongation growth of primary and lateral roots [7].

Iron play's a critical role in metabolic processes such as DNA synthesis, respiration, and photosynthesis. It serves as a component of many vital enzymes such as cytochromes of the electron transport chain, and it is thus required for a wide range of biological functions. In plants, iron is involved in the synthesis of chlorophyll, and it is essential for the maintenance of chloroplast structure and function [8].

Manganese plays an essential role in respiration and N metabolism in plants. Also act as an activator for the enzymes, nitrite reductase and hydroxylamine reductase. Manganese is directly or indirectly involved in chloroplast formation and probably in their multiplication and also involved in various oxidation-reduction reactions in photosynthesis [9].

2. MATERIAL AND METHODS

The present investigation was carried out at Horticulture Research Farm, Department of Horticulture, Babasaheb Bhimrao Ambedkar University (A Central University), Vidya-Vihar,

Rae Bareli Road, Lucknow 226 025 (U.P.), India during 2019-20. The Horticulture Research Farm, Department of Applied Plant Science Horticulture, Babasaheb Bhimrao Ambedkar University, Vidya- Vihar, Rae Bareli Road, Lucknow is situated at an elevation of 111 meters above mean sea level in the subtropical tracts of Central U.P. at 260-560 North latitude. The experiment is laid out in randomized block design with a factorial concept containing 8 treatment combinations (T₀, T₁, T₂, T₃, T₄, T₅, T₆, T₇) in which T₀: Control (Water Spray). T₁: Zinc Sulphate 0.5%, T₂: Manganese Sulphate 0.5%, T₃: Iron Sulphate 0.4% T₄: Zinc Sulphate + Manganese Sulphate 0.5% + 0.5%, T₅: Zinc Sulphate + Iron Sulphate 0.5% + 0.4%, T₆: Manganese Sulphate + Iron Sulphate 0.5% + 0.4%, T₇: Zinc Sulphate + Manganese Sulphate + Iron Sulphate 0.5% + 0.5% + 0.4%. Application of treatment was given in two doses 1st spray at Fruit Set and 2nd Spray 15 days after Fruit Set.

Plant parameters recorded during experiment are Physico-chemical characteristics., fruit size (cm), fruit weight (gm), volume of fruit (cm³), specific gravity and chemical parameters are Total soluble solids (TSS °Brix), acidity (%), ascorbic acid (Vitamin 'C') (mg/100gm pulp), reducing sugars (%), non-reducing sugar (%) and total sugar (%) are recorded. The data mended during the experimentation were statistically by randomized block design as suggested by Panse et al. [10].

3. RESULTS AND DISCUSSION

3.1 Fruit Characteristics (Physical) of Guava

The result revealed the application of micronutrients on Physico-chemical parameters and yield of guava. The parameters viz. Fruit length, Fruit breadth, Fruit weight, Fruit volume, Fruit set, Fruit drop, Total soluble solids (TSS),

total sugar, ascorbic acid content, number of fruits per plant, yield and yield efficiency.

Fruit length and breadth the data reveal that different micronutrients application exhibited a significant influence on fruit length and breadth during the study. Maximum fruit length (64.33 mm) was recorded in T₇. Whereas, the minimum fruit length (54.73 mm) was observed under treatment T₂. Maximum fruit breadth (68.03 mm) was recorded in T₇. Minimum fruit breadth (61.12 mm) was, however, observed in T₀ (control).

The data on fruit weight as influenced by different micronutrients application reveals that different micronutrients application had a significant effect on fruit weight which ranged from 116.93 g – 165.37 g. Maximum fruit weight (165.37 g) was recorded in T₇. The minimum fruit weight (116.93 g) was, however, observed in T₀ (control).

Different micronutrients affected fruit volume considerably whose value varied from 121.67 cc to 182.67 cc. The highest fruit volume (168.33 cc) was in T₇ and the minimum (121.67 cc) in T₀ (control). As indicated above, the application of different micronutrients alone or in combination has a significant effect on fruit size (length and breadth), weight and volume. The possible reason for this effect might be because mineral nutrients appear to have an indirect role in hastening the process of cell division and cell elongation due to which the size of the fruit might have improved. The increase in fruit weight may also be due to the rapid increase in the size of cells or it is also because foliar application of boron increased the fruit weight eventually by maintaining a lighter level of auxins in various parts of the fruits which helped in increasing the fruit growth [11]. The findings are similar to those reported by Bagali et al., [12], Pal et al., [13], Kumawat et al., [14] in guava and Meena et al., [15] in aonla who recorded maximum fruit size and weight with the foliar application of micronutrients.

Table 1. Treatment details

S. No.	Treatment notation	Treatment content
1.	T ₀	Control (Water Spray)
2.	T ₁	Zinc Sulphate (0.5 %)
3.	T ₂	Manganese Sulphate (0.5 %)
4.	T ₃	Iron Sulphate (0.4%)
5.	T ₄	Zinc Sulphate + Manganese Sulphate (0.5 % +0.5 %)
6.	T ₅	Zinc Sulphate + Iron Sulphate (0.5 % +0.4%)
7.	T ₆	Manganese Sulphate + Iron Sulphate (0.5 % +0.4%)
8.	T ₇	Zinc Sulphate + Manganese Sulphate + Iron Sulphate (0.5 % + 0.5 % +0.4%)

The fruit set was not much influenced by micronutrients application during the study (Table 2). However, the maximum fruit set (61.32 %) was recorded in T₇ and the minimum (60.03%) in T₁. As is evidenced from the data in Table 3, the micronutrients alone or in combinations had no significant effect on the fruit set. The possible reason for non-significant effect on the fruit set in the present study might be because the foliar application of micronutrients started after the fruit set.

It is evident from the data presented in Table 2 that those different treatments of micronutrient application exhibited a marked influence on the extent of fruit drop during the study. Minimum fruit drop (42.08 %) was recorded in T₇. Maximum fruit drop (58.60 %) was recorded in T₀ (control).

The data in Table 2 indicate that the foliar application of different micronutrients alone or in combination resulted in a significant effect on fruit drop. The minimum fruit drop was observed in trees sprayed with zinc sulphate @ 0.5% + manganese sulphate @ 0.5% + iron sulphate @ 0.4% (T₇). The results conform with those of Hada et al. (2014); Bagali et al. [12]; Balakrishnan (2000) who found a reduction in fruit drop in guava with foliar application of micronutrients. The reduction in fruit drop might be because the foliar application of micronutrients affected metabolic activities of the tree, improved the source-sink relationship and favourably influenced the metabolic status resulting in better control of drop and enhancing the retention of the fruits (Katiyar et al., 2008). Wright (1956) also suggested that the primitive effect of growth substances attributed to greater retention of fruit and reduction in fruit drop. There is a correlation between fruit drop and endogenous hormonal status, and the existence of a high level of internal auxin is useful for preventing fruit drop. Since a high level of

endogenous hormones might help in building up endogenous hormones at level that might be potent enough to reduce the fruit drop. The application of zinc could have promoted the auxin synthesis in the plant system which might have delayed the formation of the abscission layer during the early stages of fruit development (Nason and McElroy, 1963).

3.2 Chemical Parameters of Fruits

The result revealed the application of micronutrients to chemical parameters of guava. Data about the effect of micronutrients application on total soluble solids in fruits are given in Table 3. It is evident from the data that the application of micronutrients exerted a significant influence on the total soluble solid contents of fruits. Maximum total soluble solids (10.63° B) were recorded in the fruits of the plants treated. The minimum total soluble solids (8.45° B) were, however, found in T₀ (control). The increase in TSS has also been recorded by Yadav et al. [16], Awasthi and Lal [17], and Singh et al. [18] in guava. An increase in total soluble solids might be because boron helped in trans-membrane sugar transport. Micronutrients play important role in vital plant metabolic functions such as chlorophyll synthesis, various enzymatic reactions, respiration and photosynthesis [19-22]. Given that the main product of photosynthesis is sugar, so increasing the photosynthesis, leads to an increase in the sugar compounds and causes more soluble solids in fruit juice (Ram and Bose, 2000).

The scrutiny of data given in Table 3 indicates that the application of different micronutrients and their combinations exerted a significant effect on total sugars during present study. The highest total sugars (7.51%) were induced in fruits under T₇. While the lowest total sugars (5.74%) were recorded with T₀ (control).

Table 2. Effect of micronutrients on fruit characteristics (physical) of guava

Treatments	Fruit length (mm)	Fruit breadth (mm)	Fruit weight(g)	Fruit volume(cc)	Fruit set (%)	Fruit drop (%)
T ₁	59.46	64.86	136.43	140.00	60.03	45.66
T ₂	54.73	63.22	123.33	127.33	60.08	48.84
T ₃	55.98	63.29	120.70	126.33	60.89	49.16
T ₄	62.65	66.63	155.07	158.00	61.01	43.62
T ₅	59.14	65.08	140.10	145.67	60.66	44.29
T ₆	55.12	63.45	125.53	130.67	60.33	48.52
T ₇	64.33	68.03	165.37	168.33	61.32	42.08
T ₀	55.16	61.12	116.93	121.67	61.34	58.60
CD _{0.05}	4.87	2.49	2.41	4.88	NS	1.91

Statistical analysis of data about the ascorbic acid content of guava provides substantiation that different micronutrients application individually or in combinations pronouncedly increased the ascorbic acid content as per Table 3. Maximum ascorbic acid content (161.77 mg/100 g pulp) was recorded with T₇. Whereas, minimum ascorbic acid content (140.04 mg/100 g pulp) was found in T₀ (control).

The above observations of the present study indicate that the total sugars, reducing sugars, non-reducing sugars and ascorbic acid content were significantly affected by micronutrients applications alone or in combination (Table 3). These results conform to the results achieved by Kumar et al., (2015) and Jat and Kacha [23]. According to them, micronutrients application increased the quality parameters of guava fruit. The increase in the sugars content might be due to the more rapid translocation of sugars from leaves to developing fruits. Boron facilitated sugar transport within the plant. It was also reported that borate reacted with sugar to form a sugar-borate complex. Boron acted as a switcher in the degradation of glucose either by glycolysis or by pentose sugar pathway (Singh and Kaur 2016). The improvement in the quality of fruit might be due to the catalytic action of micronutrients particularly at higher concentrations. The foliar application of micronutrients quickly increased the uptake of macronutrients in the tissues and organs and improves fruit quality. The boron also plays an important role in activating the synthesis of ascorbic acid [24]. The present results on ascorbic acid conform to the results achieved by Jeyabaskaran and Pandey [25].

3.3 Yield Parameters

Data about the effect of different micronutrients and their combinations on several fruits per plant

in guava is summarized in Table 4. The data indicated that the number of fruits per plant in guava increased significantly with the foliar application of micronutrients. The maximum number of fruits per plant (45.33) was recorded under treatment T₇. The minimum number of fruits per plant (37.00) was recorded under treatment T₀ (control).

It is clear from the data given in Table 4 that the application of micronutrients exerted a significant effect on fruit yield per plant during study. Maximum fruit yield (7.50 kg/ tree) was observed under T₇. While the minimum fruit yield (4.33 kg/ tree) was recorded in treatment T₀ (control), which was significantly lower than all other treatments.

The application of micronutrients exerted a significant influence on yield efficiency (kg/cm² TCSA) during study. The data in the Table 3 showed that the highest yield efficiency of 0.35 kg/cm² TCSA was recorded in T₇ and the lowest yield efficiency of 0.16 kg/cm² TCSA was recorded in T₅. As far as the effect of micronutrients on guava yield and yield efficiency is concerned, the increase in fruit yield with the application of micronutrients may be attributed to increased fruit size, fruit weight, and minimum fruit drop resulting from the effects of micronutrients on cell division, cell elongation and translocation of photosynthetic and metabolites from leaves and others parts of plants to the developing fruits [26-28]. The highest fruit yield which was obtained by foliar spray of micronutrients may be attributed to better uptake and mobilization of nutrients to the sink which caused better fruit development. These findings are also supported by earlier reports by Bagali et al. [12], Rajkumar et al. (2014), Jat and Kacha [23], and Gaur et al. [29] who also found that foliar application of micronutrients increases the yield of guava.

Table 3. Effect of micronutrients on TSS and titratable acidity of guava

Treatments	TSS (° B)	Total sugar (%)	Ascorbic acid (mg/100 g pulp)
T1	10.63	7.12	164.24
T2	9.66	6.69	147.66
T3	9.50	6.76	146.79
T4	10.26	7.05	159.49
T5	9.90	6.66	156.40
T ₆	9.75	6.53	152.79
T ₇	10.56	7.51	161.77
T ₀	8.45	5.74	140.04
CD _{0.05}	0.40	0.35	6.46

Table 4. Effect of micronutrients on number of fruits per plant, yield, and yield efficiency of guava

Treatments	Number of fruits/plants	Yield (kg/tree)	Yield efficiency(kg/cm ² TCSA)
T ₁	42.67	5.82	0.23
T ₂	40.67	4.91	0.20
T ₃	44.33	6.87	0.27
T ₄	43.33	6.07	0.24
T ₅	41.33	5.19	0.16
T ₆	46.00	7.35	0.34
T ₇	45.33	7.50	0.35
T ₀ (control)	37.00	4.33	0.25
CD (5%)	1.91	0.31	0.02

1. CD has been calculated based on percentage value

4. CONCLUSION

On the basis of the results obtained in the present investigation, it is concluded the combined foliar application of zinc sulphate (ZnSO₄) @ 0.5% + manganese (MnSO₄) (0.5%) + iron sulphate (FeSO₄) (0.4%) (T₇). Therefore, combined spray of ZnSO₄ (0.5%) + MnSO₄ (0.5%) + FeSO₄ (0.4%) can be advocated to guava growers for maximum physical, chemical and yield parameters of guava.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Subhash Chander MC, Jain PK, Pareek PK, Bola RR, Meena YK, Sharma, Renuka. Effect of Foliar Feeding of Borax, Zinc sulphate and Urea on Fruiting and Yield of Guava (*Psidium guajava* L.) Cvs. Lalit and Shweta under high density planting system. Chem. Sci. Rev. 2017;6(22):874-883.
- Manoj Gaund, Ram D, Anand Singh Rawat, Ajendra Kumar. Response of foliar application of micronutrients and plant growth regulator on yield and economic feasibility of guava (*Psidium guajava* L.) CV. Shweta and Lalit. The Pharma Innovation Journal. 2022;11(3):1752-1756.
- NHB. State wise area, production and productivity of guava; 2017. Available: www.nhb.gov.in
- Ruby Rani, Brahmachari VS. Effect of growth substance and calcium compounds on fruit retention, growth and yield of Amrapali mango. Orissa J. Hort. 2004; 32(1):15-18.
- Bose TK, Mitra SK, Sanyal D. Fruits: tropical & sub-tropical. Naya Prakashan, Calcutta, West Bengal. 1990;838.
- Price CA, Clark HE, Funkhouser EA. Functions of micronutrients in plants. In: Micronutrients in agriculture. Soil Science Society America, Madison, Wisconsin. 1972;173-88.
- O' kelley JC. Boron effects on growth, oxygen uptake and sugar absorption by germinating pollen. American Journal of Botany. 1957;44:239-44.
- Rout GR and Sahoo S. Role of iron in plant growth and metabolism. Reviews in Agriculture Science. 2015;3:1-24.
- Nijjar GS. Nutrition of fruit trees. 2nd ed. Kalyani Publishers, Ludhiana. 1990;311.
- Panse VG, Sukhatme PV. Statistical methods for agricultural workers. Indian Council of Agricultural Research Publications, New Delhi. 1995;330.
- Kaur S. Effect of micronutrients and plant growth regulators on fruit set, fruit retention, yield and quality attributes in litchi cultivar Dehradun. Chemical Science Review and Letters. 2017;6:982-86.
- Bagali AN, Hulamani NC and Sulikeri GS. Effect of foliar application of zinc, magnesium and boron on growth and yield of guava (*Psidium guajava* L.) cv. Sardar. Karnataka Journal of Agricultural Sciences. 1993;6:137-41.
- Pal A, Pathak RK, Pal K and Singh T. Effect of foliar application of nutrients on yield and quality of guava (*Psidium guajava* L.) fruit cv. Sardar. Progressive Research. 2008;3:89- 90.
- Kumawat KL, Sarolia DK, Shukla AK. Growth, yield and quality of rejuvenated guava as influenced by thinning-bending and micronutrients. Indian Journal of Horticulture. 2012;69:478-83.

15. Meena D, Tiwari R, Singh OP. Effect of nutrient spray on growth, fruit yield and quality of aonla. *Annals of Plant and Soil Research*. 2014;16:242-45.
16. Yadav A, Ram RB, Verma RS, Kumar V, Yadav RK. Effect of foliar application of micronutrients on bio-chemical attributes of winter season guava (*Psidium guajava* L.) cv. Lalit. *Journal of Pharmacognosy and Phytochemistry*. 2018;7:3196-97.
17. Awasthi P, Lal S. Effect of calcium, boron and zinc foliar sprays on the yield and quality of guava (*Psidium guajava* L.). *Pantnagar Journal of Research*. 2009; 7:223-25.
18. Singh R, Chaturvedi OP, Singh R. Effect of pre-harvest spray of zinc, boron and calcium on the physico-chemical quality of guava fruit (*Psidium guajava* L.). *Internal Seminar on Recent Trend in Hi-Tech Horticulture and Post-Harvest Technology*. 2004;4-6.
19. Dhaliwal GS, Nanra NK, Rattanpal HS. Effect of chemicals on flower drop, fruit set and yield on rainy and winter season crops of guava. *Indian Journal of Horticulture*. 2002;59:31-33.
20. Gaur B, Hada TS, Beer K, Kanth N, Syamal MM. Studies on the effect of foliar application of nutrients and GA₃ on yield and reproductive parameters of winter season guava. *Trends in Biosciences*. 2014b ;7:3386-89.
21. Ghosh SN. Effect of magnesium, zinc and manganese on yield and fruit quality of guava. *South Indian Horticulture*. 1986;34:327-30.
22. Kaur G, Dhillon WS. Effect of foliar application of chemicals on physico and chemical characters of variety Allahabad Safeda of guava during winter. *Journal of Reseach*. 2006;43:114-16.
23. Jat G, Kacha HL. Response of guava to foliar application of urea and zinc on fruit set, yield and quality. *Journal of Agri Search*. 2014;1:86-91.
24. Baranwal D, Tomar S, Singh JP, Maurya JK. Effect of foliar application of zinc and boron on fruit growth, yield and quality of winter season guava (*Psidium guajava* L.). *International Journal of Current Microbiology and Applied Sciences*. 2017; 6:1525-29.
25. Jeyabaskaran KJ, Pandey SD. Effect of foliar spray of micronutrients in banana under high soil pH condition. *Indian Journal of Horticulture*. 2008;65:102-05.
26. Ali W, Pathak RA, Yadav AL. Effect of foliar application of nutrients on guava (*Psidium guajava* L.) cv. Allahabad Safeda. *Prog. Hort*. 1993;23(1-4):14-21.
27. Aulakh PS. Effect of seasonal variation on yield and fruit quality of some promising guava cultivars under arid irrigated region of Punjab. *Haryana J. Hort. Sci*. 2004;33:170-71.
28. Das BC, Chakraborty A, Chakraborty PK, Maiti A, Mandal S, Ghosh S. Comparative performance of guava cultivars under red and laterite soils of West Bengal. *Horticultural Journal*. 1995;8:141-46.
29. Gaur B, Beer K, Hada TS, Kanth N, Syamal MM. Studies on the effect of foliar application of nutrients and GA₃ on fruit yield and quality of winter season guava. *The Ecoscan*. 2014a;6:479-83.

© 2022 Meena and Verma; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/86097>