



A New Record to Rust Fungi of South Western Province Kandahar, Afghanistan

Hayatullah Ahmadi ^{a*}, L. I. Changtian ^a and Qudratullah Ehsan ^b

^a *Engineering Research Center of Chinese Ministry of Education for Edible and Medicinal Fungi, Jilin Agricultural University, Changchun, Jilin - 130118, China.*

^b *Department of Agronomy, Faculty of Plant Sciences, Afghanistan National Agricultural Science and Technology University (ANASTU), Kandahar, Afghanistan.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2022/v34i2331599

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/92165>

Original Research Article

Received 15 July 2022
Accepted 29 September 2022
Published 30 September 2022

ABSTRACT

A rust fungus occurring on *Centaurea iberica* leaves from Arghandab district, southwestern province Kandahar Afghanistan is described and illustrated. A critical morphological and microscopical examination revealed it to be *Puccinia calcitrapae*. After conducting a literature survey, it was found that this fungus is the first record to be reported from southwestern province Kandahar. A detailed taxonomic description of this fungus along with its distribution is provided. Weeds cause serious problems in agriculture in Afghanistan especially in southwestern province Kandahar. These plants reduce yield and the quality of crops by competing for water, nutrients and sunlight. The improper or excessive usage of herbicides has led to development of resistance in some weed species while contaminating the environment; therefore, biological control has an increasing role as an alternative method for controlling special weed. Previous studies in other countries revealed that this rust fungus are good biological control agents for these weeds.

Keywords: *Centaurea*; *Puccinales*; *systematics*; *taxonomy*.

1. INTRODUCTION

Rust fungi are highly obligate parasites of plants belongs to the order Puccinales

(previously Uredinales) in the sub phylum Pucciniomycotina, phylum Basidiomycotina and possess 7,800 species, 166 genera and 14 families worldwide [1]. Thirteen out of the 14

*Corresponding author: E-mail: Hayatullahahmadi114@gmail.com;

families are well described and accepted. [2]. The majority of rust species belong to two genera *Puccinia* and *Uromyces* [3].

Rust fungi are obligate parasites of vascular plants, i.e., ferns, gymnosperm and most families of angiosperm [3], while currently about 30 species are successfully cultured on artificial media [4-6]. Since some of the species caused economically important diseases of crops, vegetable, orchard and trees. In addition, these fungi exhibit a wider diversity and broader host range and their infection are not only limited to agricultural crops but also to non-agricultural plants including medicinal herbs, shrubs, and even weeds (dr. kiran r ranadive 2021). Biology and taxonomy of the rust fungi have been broadly studied and well cataloged from the beginning of last century. especially, the rust fungi of north america, europe, brazil, japan, australia, New Zealand, and argentina [2]. However, some important new species and perhaps genera are still expected, especially in tropical and subtropical regions of central and south america, africa, and south asia [2].

Rust fungi are the most devastating and economically important pathogen of both domesticated and wild plants worldwide [2]. Rust fungi are responsible for some of enormous economic losses of trees and crops, including pines, poplars, grape, wheat, corn, coffee, sugarcane, and soybean rusts which are greater than any other single biotic stress. The improper or excessive usage of herbicides leads to the development of resistance in the target weed species as well as contamination of the environment; therefore, biological control has become an alternative method in the case of controlling a single weed species [7]. Shortly biological control of weeds can offer alternatives on economically and environmentally safe approach to reach sustainable agricultural production.

Centaurea is a genus of herbaceous thistle like flowering plants comprise more than 300 species in the family of Asteraceae. Members of the genus are found in equator, mostly in the Eastern Hemisphere and the Middle East. There are 11 species of *Centaurea* including *Centaurea iberica* (Spreng., Syst Veg.3: 406.1826) identified in Afghanistan. Many species in particular those inhabiting more arid regions have a long top root system, mostly found in pastures or meadows, also found in mesotrophic grasslands. Due to their habit of

dominating ecosystem under good conditions, many *Centaurea* species can become invasive weeds in region where they are not native meanwhile, is inedible to most livestock due to its spines and apparently outright poisonous to horses and equines. *Centaurea iberica* as a weed is a serious problem in Kandahar province. Previous studies in other countries revealed that this rust fungus are good biological control agents for these weeds.

Rust survey were performed two seasons, spring and early fall (May 2019 & September 2020). We collected rust fungi on *Centaurea iberica* leaves from Arghandab, Dand and Daman districts of south western province Kandahar, Afghanistan. After critical morphological examination of the diseased leaf samples revealed it to be a species of *Puccinia*. A detailed and precise literature survey and comparative analyses [8-11] revealed that this fungus is the first record for southwestern province Kandahar. Therefore, a detailed taxonomic description of the species is provided.

2. MATERIALS AND METHODS

The survey of rust fungi was performed two times at two different seasons, spring (from May 05 to June 07, 2019) and early fall (from September 15 to October 05, 2020). Rust-infected leaf of *Centaurea iberica* were collected into plastic bags and were air-dried to make specimens of rust infected plants and taken to the laboratory for identification and taxonomical studies.

The rust sori were observed under stereo-microscope (meji techno, emz-5tr; japan). Free hand section of spores taken from the sori were mounted on a glass slide with lactic acid or lacto phenol for light microscopic observation. Thirty to 60 spores were randomly chosen from each specimen for observation under a light microscope (olympus bx50f4 or olympus bh2, japan). Morphological characteristics of spores such as length, width, wall thickness, shape, surface ornamentation, color, number and location of germ pores were recorded. In case of teliospores, number of cells and pedicel were also recorded. Paraphysis and other structures included in sori were also observed. Ten to 30 spores of each spore states were measured by using an image analyzer photoruler (onishi 2009.2010). All Identified specimen is kept in afghanistan agriculture and sciences and technology university plant sciences herbarium (anastu-plsh), kandahar, afghanistan.

3. RESULTS

Initial symptoms of rust fungi were observed during May to June 2019, as grayish brown spots in the upper surface (adaxial) of leaves which later coalesced to form spindle shaped yellow to orange- yellow powdery pustules were noticed on corresponding lower (adaxial) surface. The infection was limited to mature leaves and no rust symptoms were found on juvenile or young leaves. The development of fungal sori was observed as dark brown longitudinal streaks mainly on the adaxial surface of the leaves. The detailed description and illustrations (Fig. 1) of the fungi along with a discussion on its taxonomy and distribution are presented below.

3.1 Taxonomy

Puccinia calcitrapae DC.1805.

Spermogonia and acia are known in other countries but not known in Afghanistan. Uredinia are amphigenous brown to dark-brown without paraphysis; rediniospores are yellowish brown to cinnamon brown, mostly globose to sub globose and 20-28 × 18-24 μm thick. Walls are 1.5-2 μm thick pale yellow to pale brown, echinulate, germ pores 3 rarely 2 sometimes or 4 equatorials.

Telia are amphigenous, chocolate brown to blackish brown; teliospores are dark-brown, 28-37 × 19-24 μm in size and ellipsoid. Walls are smooth 1.5-2 μm thick at sides; pedicels are hyaline 10-15 μm long.

Known Distribution - worldwide in distribution (Asia, America, Europe, Africa, Australia). This rust has been reported in Afghanistan from Kabul, Maymana, Bamyan, Baghlan, Herat, Badakhshan, Kanduz and Balkh (Mazar-i-Sharif) (Fig. 2).

Material examined – Afghanistan, Kandahar (1010 m) on leaves of *Centaurea iberica*, June 15, 2019, Hayatullah Ahmadi, & Qudratullah Ehsan, (ANASTU-PLSH-0019).

4. DISCUSSION AND CONCLUSION

Rust fungus on *Centaurea* spp. collected in Afghanistan was identified *Puccinia calcitrapae* [8-12] or *P. centaureae* [13], Brandenburger and Steiner 1972, [11]. They are morphologically very similar and their taxonomic treatments were different among the uredinologists.

For example, Wilson and Henderson [9] adopted very wide species concept for the rust fungi with similar morphology on similar host plants (*Arctium*, *Carduus*, *Centaurea*, *Cirsium* and *Cardina* in the tribe Cynareae of the family Compositae) and treated them as *P. calcitrapae*, which contained numbers of races or special forms. *Puccinia centaureae* was considered as synonym of *P. calcitrapae*. Records of *P. calcitrapae* in Afghanistan by Henderson and Jørstad [8], Hederson [10] and Gjaerum [12] might be based on this wide species concept.




Fig. 1. *Puccinia calcitrapae*. (A) Urediniospores. (B-C) Teliospores. The scale bare for all pictures are 30 μm




Fig. 2. Distribution map of *Puccinia calcitrapae* DC. 1805 in Afghanistan (Current report)

Table 1. *Puccinia calcitrapae* recorded from Afghanistan, host, location and references

Species	Host	Location	References
<i>Puccinia calcitrapae</i> DC.1805	<i>Acroptilon austral</i>	Mymana	Henderson and Jørstad [8]
	<i>Acropilon repens</i>	Mazar-i-Sharif	Henderson, D. M. and Joerstad.[8]
	<i>Carduus pycnocephalus</i>	Fariab	Henderson,D.M. [10]
	<i>Centaurea iberica</i>	Balkh	Gjaerum, H. B.[12]
	<i>Centaurea iberica</i>	Mazar-i-Sharif	Henderson, D. M. and I. Joerstad.[8]
	<i>Centaurea iberica</i>	Unknown province	H. B. [8], Sydowia Durrieu, G.[11]
	<i>Centaurea pulchella</i>	Kabul	Henderson, D. M. and I. Joerstad.[8]
	<i>Centaurea sp.</i>	Kunduz	Gjaerum, H. B. [2]
	<i>Cousinia leptacatha</i>	Mazar-i-Sharif	Henderson D. M. and I. Joerstad. [8]
	<i>Centaurea mirocarpa</i>	Mymana	Henderson, D. M. and Joerstad. [8]
	<i>Centaurea mirocarpa</i>	Mazar-i-Sharif	Henderson, D. M. and I. Joerstad [8]
	<i>Cousinia sp.</i>	Badakhshan	Henderson D. M. [10]
	<i>Cousinia sp.</i>	Herat	Henderson, D. M. [10]

Species	Host	Location	References
	<i>Cousinia</i> sp.	Maymana	Henderson, D. M. and I. Joerstad. [8]
	<i>Cousinia</i> sp.		Henderson, D. M. and I. Joerstad. (1966)
	<i>Cousinia</i> sp.	Bamyan	Henderson, D. M. and I. Joerstad. [8]
	<i>Cousinia</i> sp.	Kabul	Henderson, D. M. and I. Joerstad. [8]
	<i>Cousinia</i> sp.	Qataghan	Henderson, D. M. and I. Joerstad. [8]
	<i>Centaurea codringtonii</i>	Unknown Province Baghlan	Durrieu, G. [11] Henderson, D. M. and I. Joerstad. [8]
	<i>Centaurea iberica</i> <i>Centaurea iberica</i>	Baghlan,	Brandenburger, W. and M. Steiner. (1972)
	<i>Cousinia</i> sp.	Unknown Province	Petrak, Franz. [13]
	<i>Cousinia</i> sp.	Bamyan	Petrak, F. [13]

Savile [14] considered these two species were distinguishable based on precise morphological composition and host plants. They differed in surface structures of urediniospores and teliospores, and germ pores position of urediniospores. *Puccinia calcitrapae* was found only on *Centaurea calcitrapeae* collected in Europe, while *P. centaureae* (as *P. centuraeae* var *centuraeae*) was found on many other *Centaureae* species in Europe and U.S.S.R. Durrieu [11] followed Savile's taxonomic treatment and identified the rust on *Centuraea iberica* as *P. calcitrapae* and the one on *C. condingtonii* as *P. centaureae*.

Cummins (1977) treated North American population on *Carthamus*, *Centaurea* and *Cirsium* as a variety, *P. calcitrapae* var. *centuraeae* (= *P. centaureae*). Hiratsuka et al. [15] followed Cummins and treated Japanese population on *Carduus*, *Carthamus* and *Cirsium* as *P. calcitrapae* var. *centuraeae*.

In the present study, two specimens were identified as *Puccinia calcitrapae sensu* Wilson and Henderson [9]. Sizes of urediniospores and teliospores on *Cardus crispus* were slightly larger than those on *Centaurea ibrrica*.

Centaurea ibrrica caused by rust fungi *puccinia calcitrapae* has a worldwide distribution including asia, america, europe, africa and australia. Eleven species of the genus *centaurea* has been reported from Afghanistan [16,17]. Many *centaurea* species can become invasive weeds

in region where they are not native meanwhile, is inedible to most livestock due to its spines and apparently outright poisonous to horses and equines. *Centaurea iberica* as weed is a serious problem in kandahar province. Previous studies in other countries revealed that this rust fungus are good biological control agents for these weeds [18].

ACKNOWLEDGEMENTS

The author gratefully thanks their respective organization for providing laboratory facilities to carry out the present work. We also express our thanks for encouragement and every possible support provided by Dr. LI, Changtian during this scientific study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Kirk PM, Cannon PF, Minter DW, Stalfer JA. (Eds). Dictionary of the Fungi, 10th edn. CABI Publishing, UK; 2008.
2. Cummins GB, Hiratsuka Y. Illustrated genera of rust fungi, 3rd Edn, APS press, St. Paul, MN; 2003.
3. Helfer S. Rust fungi and global change. New Phytologist. 2013;201:770-780.
4. Maclean DJ. Axenic culture and metabolism of rusr fungi. In Scott KJ, Chakravorty AK (Ed) The rust fungi Academic Press. 1982;37-120.

5. Williams PG. Obligate parasitism and axenic culture. 1984;399-430.
6. Yamaoka Y. Axenic culture of rust fungi (in Japanese). Riken Symposium, Yet-to-be cultured microorganism and cultural collection, Riken. 2002;14-17.
7. Muller-Scharer, Frantzen. 1966. Evaluation of rust fungi as biological control agents of weedy *Centaurea* in North America. *Weed Science*. 1986;34:7-10.
8. Henderson DM, Joerstad I. Studies in the Flora of Afghanistan. II Uredinales. Norwegian University press. 1966;4:2-18.
9. Wilson M, Henderson DM. British rust fungi; 1966.
10. Henderson DM. Notes on three british ascomycetes. Notes from the Royal Botanical Garden Edinburgh. 1970;30: 203-207.
11. Durrieu G. Micromycetes parasites d'Afghanistan. *Revue de Mycologie*. 1975;39:137-171.
12. Gjaerum HB. 1996. Rust fungi (Uredinales) collected on the finnish botanical expedition to West-Central Asia. *Lidia*. 1972;3:195-204.
13. Petrak F. Ein Beitrag zur pilzflora von Afghanistan. *Sydowia*. 1963;16:331-349.
14. Savile DBO. Some eurasian puccinia species attacking cardueae. *Canadian Journal of Botany*. 1970;48(9):1553-1566.
15. Hiratsuka N, Sato S, Katsuya K, Kakishima M, Hiratsuka Y, Kaneko S, Ono Y, Sato t, Harada Y, Hiratsuka T, Nakayama K. The rust flora of Japan. Tsukuba Shuppankai, Japan; 1992.
16. Gjaerum HB. Rust fungi (Uredinales) from Iran and Afghanistan. *Sydowia*. 1986;39: 68-100.
17. Gjaerum HB. Studies in rusts (Uredinales) on *Astragalus* (Fabaceae). *Edinburgh Journal of Botany*. 1991;48:393-401.
18. Gautam AK, Avasthi S, Verma RK, Devadatha B, Sushma, Ranadive KR, Bhadauria R2, Prasher IB and Kashyap PL. Current status of research on rust fungi (*Pucciniales*) in India. *Asian Journal of Mycology*. 2021;4(1):40-80.

© 2022 Ahmadi et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/92165>