



Ichthyofauna in the Lomami National Park and Its Hinterlands, Democratic Republic of the Congo

**Ekyamba Shabani ^{a*}, Ernest Lukusha Tambwe ^b,
Oscar Ndeo Wembo ^c, Alain Bolonga Bolonga ^d,
Roger Bolaya Lingofu ^d and Alidor Busanga Kankonda ^d**

^a *Department of Conservation and Management of Renewable Natural Resources, Faculty of Science and Technology, University of Goma, P.O. Box 204 Goma, DR Congo.*

^b *Département de Gestion des Ressources Naturelles Renouvelables, Institut Supérieur d'Etudes Agronomiques de Bengamisa, BP 202 Bengamisa, DR Congo.*

^c *Ruwenzori State University, P.O. Box 560 Butembo, DR Congo.*

^d *Laboratory of Hydrobiology and Aquaculture, Department of Hydrobiology, Faculty of Science, University of Kisangani, PO Box 2012 Kisangani, DR Congo.*

Authors' contributions

This work was carried out in collaboration among all authors. Authors' ES, ELT and ONW collected data. Authors' RBL and ABB identified species. Author ES performed the statistical analysis and wrote the first draft of the manuscript. Authors' ABK and ES managed the analyses of the study. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJFAR/2023/v25i3676

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/104110>

Original Research Article

Received: 05/06/2023

Accepted: 10/08/2023

Published: 30/09/2023

ABSTRACT

This investigation deals with freshwater fish populations in thirteen sampling rivers of Lomami National Park (LNP) and its hinterlands. The study aimed to inventory the ichthyofauna in freshwater bodies of LNP and its hinterlands; determine fish species with high economic value according to preference by riparian communities; and assess the conservation status of fish

*Corresponding author: Email: shabaniekyamba@unigom.ac.cd;

species. Intensive fish sampling was conducted thirteen rivers of LNP and its hinterlands using gill nets with stretched mesh sizes of 9.5, 10, 12, 15 and 20 mm in 2016. A survey questionnaire was administered to fishermen in order to determine fish species with high economic value. A total of 2985 fish were caught in thirteen rivers belonging to 129 species, 54 genera, 20 families and 8 orders. The family of Mormyridae dominated in the fish collection with 30 species followed by family of Alestidae with 23 species. *Clarias* spp., *Schilbe* spp., *Chrysichthys* spp., *Auchenoglanis occidentalis*, *Polypterus* spp., *Hydrocynus* spp., *Alestes* spp., *Parachana* spp. and species of Mormyridae were recorded as fish species with high economic value. The majority of species was least concerned, but one was reported as endangered species (*Nannothrissa sterwarti*) and another vulnerable (*Labeo rectipinnis*). The analysis displayed that all rivers were diverse with a minimum observed in the Onema River. The probability of catching two individuals of different species in the same river was high in all rivers, but it was higher in the Amboko, Lodja, Lomami and Yalo Rivers (96%); and it was slightly low in the Onema River (73%). The Kruskal–Wallis result showed statistical differences among sampling rivers ($P = 0.0000$). Our findings provided valuable information for the conservation and management of fish species in the freshwater of Lomami National Park and its hinterlands.

Keywords: Fishery resources; rivers; conservation status; Lomami National Park.

1. INTRODUCTION

Fish form the most important group of vertebrates of planet earth with about 30,000 species that 41% live in freshwater [1]. Fishing and fish related activities are important for most riparian communities. The resource is not only an important protein source but also an important revenue earner for commercial fishers [2,3].

Lomami National Park (LNP) is an area with high freshwater fish species. Unfortunately, there is very little information about the biodiversity and the distribution of freshwater fish of the LNP. However, there is considerable anthropogenic pressure due to deforestation, mining logging, hunting, fishing and polluted water untreated in the most part of Central Africa [3,4,5]. These effects are increased by climate change and population growth, which cause the increase of needs in natural resources, destruction of ecosystems and extinction of species. Furthermore, the conservation status of many freshwater fish from Lomami National Park cannot be assessed adequately due to lack of data.

Thus, the knowledge of ichthyofauna, as well as behavior, ecology and biology of reproduction of fish caught are vital prerequisites, and implementation of a good fish farming production, will widely contribute to the reduction of overexploitation of natural resources, profitability for hunters of commercial bushmeat and sustainable management of Lomami National Park and its hinterlands.

This investigation aimed to provide a list of fish species from freshwater bodies of LNP and its hinterlands; determine fish species with high economic value according to preference by riparian communities; and assess the conservation status of fish species.

2. MATERIALS AND METHODS

2.1 Area of Investigation

Lomami National Park is located in eastern part of Central basin of Democratic Republic of the Congo. It is a vast forest area that extends between 1° and 3° S latitude and between 24.5° and 25.5° E longitude. This massif also includes the basins of Tshuapa River in the west and Congo River (Lualaba) in the east [6]. Sampling was carried out in thirteen rivers in the southern part of Lomami National Park (Fig. 1), including Amboko, Baleke, Ename, Lodja, Loidjo, Lomami, Nvul'elongo, Nyamatende, Onema, Petchi, Shingi, Wele and Yalo.

2.2. Data Collection Procedure

Intensive fish sampling was conducted using gill nets with stretched mesh sizes of 9.5, 10, 12, 15 and 20 mm in 2016. Gill nets were set between 04:00 and 06:00 pm at strategic positions and retrieved the following morning between 06:30 and 08:30 am (Fig. 2). The collected fish specimens were counted, grouped according to the rivers and dates of sampling and placed in labelled plastic containers filled with 10% formalin to stop the digestion process postmortem. Samples were then transported to

laboratory of Hydrobiology and Aquaculture, University of Kisangani for further processing.

A survey questionnaire was administered to riparian fishermen in order to record information on fish exploitation in the Lomami National Park and its hinterlands.

2.3 Identification of Fish

In the laboratory, sampled fish were washed with running water and the taxonomic identification was initially determined based on individual morphology using the identification keys of Boulenger [7]; Poll [8,9]; Matthes [10]; Gosse

[11]; Roberts and Stewart [12]; Banister and Bailey [13]; Teugels [14]; Poll and Gosse [15]; Tshibwabwa [16]; Shumway *et al.* [17]; Mbenga and Teugels [18]; Skelton [19]; Stiassny *et al.* [20,21]. AMNH lower Congo keys, FishBase [22]. As well as Sullivan *et al.* [23]. and fish reference collection done by the same authors were also used.

An assessment according to Red List Categories of the International Union for Conservation of Nature (IUCN) was provided for each fish species where available referring to the Pan Africa Freshwater Fish Species List [<https://www.iucnredlist.org/>].




Fig. 1. Map showing fish collection sites in the Lomami National Park and its hinterlands



Fig. 2. Capture of fish in the rivers of Lomami National Park and its hinterlands

2.4 Data Analysis

We performed biological diversity indices, including Shannon-Wiener and Simpson in terms of fish species richness abundance using PAST software. We computed the non-parametric Kruskal-Wallis test using Rcmdr package in R software to test the significance of fish species richness abundance among thirteen sampling rivers.

3. RESULTS AND DISCUSSION

A total of 2985 fish individuals were caught in thirteen rivers belonging to 129 species, 54 genera, 20 families and 8 orders. It was observed that family of Mormyridae dominated in the collection with 13 genera and 30 species followed by family of Alestidae with 9 genera and 23 species (Table 1).

As shown in the Table 2 and Fig. 3a, many fish species were caught in the Lomami river with 66 species, followed by the Amboko River (62 species), Lodja river (60 species) and Yalo River (58 species). The lowest number of fish species (11 species) was recorded in the Wele River. We experienced identification problems for Cichlid specimens caught in the Nvul'elongo River, which were not identified to genus or species levels.

The highest relative abundance was reported in the Lomami and Loidjo Rivers with 16.60% and 16.19%, respectively (Fig. 3b), and the Kruskal-Wallis test showed statistical differences among sampling rivers ($P = 0.0000$).

Mulangu [24] identified 27 species in the Lubilanji River and its two tributaries. Among them, 5

species were reported in the Lomami National Park and its hinterlands, including *Petrocephalus christyi*, *Tylochromis lateralis*, *Auchenoglanis occidentalis*, *Clarias gariepinus*, and *Schilbe grenfelli*. He found also that Mormyridae was abundant in his collection. Tumitu and Mambo [25] identified 102 species in the Lomami National Park, however 47 species of them were not found in the checklist of fish collected in 2016, including *Bathyaethiops caudomaculatus*, *Bryconaethiops microstoma*, *Hydrocynus goliath*, *Micralestes humilis*, *Micralestes stormsi*, *Micralestes* sp., *Nanopetersius ansorgii*, *Citharinus congicus*, *Distichodus altus*, *Distichodus* sp.1, *Distichodus* sp.2, *Eugnathichthys eetveldii*, *Ichthyborus besse congolensis*, *Poecilothrissa centralis*, *Nannothrissa parva*, *Microthrissa congica*, *Barbus holotaenia*, *Barbus* sp., *Clypeobarbus congicus*, *Chelaethiops congicus*, *Campylomormyrus mirus*, *Campylomormyrus* sp., *Campylomormyrus curvirostris*, *Campylomormyrus alces*, *Campylomormyrus tamandua*, *Hippopotamyrus psittacus*, *Mormyrops attenuatus*, *Mormyrus caballus*, *Petrocephalus* sp., *Petrocephalus grandoculis*, *Petrocephalus microphthalmus*, *Pollimyrus isidori*, *Stomathorinus* sp., *Microctenopoma* sp., *Tylochromis robertsi*, *Tilapia* sp., *Polypterus* sp., *Clarias angolensis*, *Chrysichthys longibarbis*, *Chrysichthys ornatus*, *Chrysichtys* sp, *Parauchangolanis punctatus*, *Malapterurus electricus*, *Synodontis acanthomias*, *Synodontis nummifer*, *Schilbe debauwi*, and *Mastacembelus congicus*. The family of Mormyridae dominated in their checklist. Kisekelwa et al [26]. collected 4 Mormyrid species and 3 Alestid species in the Lowa River, Walikale territory. Among them, 2 Mormyrid species included *Pollimyrus* sp.,

Marcusenius sp. and 1 Alestidae species *Bryconaethiops boulengeri* were identified in our checklist. They reported that Alestidae species prefer torrential locations. Byanikiro et al [27]. collected 9 species of Mormyridae in their study on characterization and ecology of stands of Mormyrids in the Reserve of Yoko. Among them, one species *Cyphomyrus psittacus* was identified in our collection. Decru et al [28]. identified 320 species belonging to 28 families in the Itimbiri, Aruwimi, and Lindi/Tshopo Rivers. Among them, Mormyridae was the most dominant family in all three rivers. Of Mormyrid species, 17 were found in the Lomami National Park and its hinterlands, including

Campylomormyrus elephas, *Campylomormyrus numenius*, *Cyphomyrus psittacus*, *Cyphomyrus wilverthi*, *Genyomyrus donnyi*, *Gnathonemus petersii*, *Hippopotamyrus weeksii*, *Marcusenius greshoffi*, *Marcusenius kutuensis*, *Marcusenius monteiri*, *Marcusenius moorii*, *Mormyrops anguilloides*, *Mormyrops nigricans*, *Myomyrus macrops*, *Petrocephalus christyi*, *Petrocephalus sauvagii*, and *Pollimyrus osborni*. Kisekelwa et al [29]. collected 13 Mormyrid species in the river systems draining the Kahuzi-Biega National Park. Among them, 5 species were found in our checklist, including *Gnathonemus petersii*, *Marcusenius greshoffi*, *Marcusenius monteiri*, *Myomyrus macrops*, and *Pollimyrus osborni*.




Fig. 3. Species richness (a), Relative abundance (b), Shannon-Wiener's index (c), and Simpson's index (d) for thirteen sampling rivers

Table 1. Systematic broad outline of fish captured in thirteen rivers of LNP and its hinterlands

Order	Family	Genus	Species	
Characiformes	Alestidae	<i>Alestes</i>	<i>Alestes liebrechtsii</i> Boulenger, 1898 <i>Alestes macrophthalmus</i> Günther, 1867	
		<i>Alestopetersius</i>	<i>Alestopetersius brichardi</i> Poll, 1967 <i>Alestopetersius</i> cfr <i>brichardi</i> <i>Alestopetersius compressus</i> (Poll & Gosse, 1963)	
		<i>Bathyaethiops</i>	<i>Bathyaethiops greeni</i> Fowler, 1949	
		<i>Brachypetersius</i>	<i>Brachypetersius altus</i> (Boulenger, 1899) <i>Brachypetersius huloti</i> (Poll, 1954) <i>Brachypetersius pseudonummifer</i> Poll, 1967	
		<i>Brycinus</i>	<i>Brycinus bimaculatus</i> (Boulenger, 1899) <i>Brycinus</i> cfr <i>bimaculatus</i> <i>Brycinus</i> cfr <i>grandisquamis</i> <i>Brycinus</i> cfr <i>macrolepidotus</i> <i>Brycinus</i> cfr <i>macrolepidotus</i> 2 <i>Brycinus grandisquamis</i> (Boulenger, 1899) <i>Brycinus imberi</i> (Boulenger, 1899)	
		<i>Bryconaethiops</i>	<i>Bryconaethiops boulengeri</i> Pellegrin, 1900 <i>Bryconaethiops macrops</i> Boulenger, 1920	
		<i>Hydrocynus</i>	<i>Hydrocynus forskahlii</i> (Cuvier, 1819) <i>Hydrocynus vittatus</i> Castelnau, 1861	
		<i>Micralestes</i>	<i>Micralestes acutidens</i> (Peters, 1852) <i>Micralestes</i> cfr <i>sardina</i>	
		<i>Phenacogrammus</i>	<i>Phenacogrammus polli</i> Lambert, 1961	
		Citharinidae	<i>Citharinus</i>	<i>Citharinus gibbosus</i> Boulenger, 1899 <i>Citharinus macrolepis</i> Boulenger, 1899
			Distichodontidae	<i>Distichodus</i>

Order	Family	Genus	Species
			<i>Distichodus sexefasciatus</i> Boulenger, 1897
			<i>Distichodus</i> cfr <i>fasciolatus</i>
		<i>Eugnathichthys</i>	<i>Eugnathichthys macroterolepis</i> Boulenger, 1899
		<i>Mesoborus</i>	<i>Mesoborus crocodilus</i> Pellegrin, 1900
		<i>Phago</i>	<i>Phago boulengeri</i> Schilthuis, 1891
			<i>Phago intermedius</i> Boulenger, 1899
		<i>Xenocharax</i>	<i>Xenocharax spilurus</i> Günther, 1867
	Hepsetidae	<i>Hepsetus</i>	<i>Hepsetus</i> sp.
Clupeiformes	Clupeidae	<i>Nannothrissa</i>	<i>Nannothrissa stewarti</i> Poll & Roberts, 1976
Cypriniformes	Cyprinidae	<i>Enteromius</i>	<i>Enteromius miolepis</i> Boulenger, 1902
		<i>Labeo</i>	<i>Labeo cyclorhynchus</i> Boulenger, 1899
			<i>Labeo greenii</i> Boulenger, 1902
			<i>Labeo lineatus</i> Boulenger, 1898
			<i>Labeo parvus</i> Boulenger, 1902
			<i>Labeo rectipinnis</i> Tshibwabwa, 1997
			<i>Labeo</i> sp.
			<i>Labeo weeksii</i> Boulenger, 1909
		<i>Labeobarbus</i>	<i>Labeobarbus caudovittatus</i> (Boulenger, 1902
		<i>Raiamas</i>	<i>Raiamas buchholzi</i> (Peters, 1876)
		<i>Heterotis</i>	<i>Heterotis niloticus</i> (Cuvier, 1829)
Osteoglossiformes	Arapaimidae	<i>Campylomormyrus</i>	<i>Campylomormyrus elephas</i> (Boulenger, 1898
Mormyriiformes	Mormyridae		<i>Campylomormyrus numenius</i> (Boulenger, 1898)
		<i>Cyphomyrus</i>	<i>Cyphomyrus psittacus</i> (Boulenger, 1897)
			<i>Cyphomyrus</i> sp.
			<i>Cyphomyrus wilverthi</i> (Boulenger, 1898)
		<i>Genyomyrus</i>	<i>Genyomyrus donnyi</i> Boulenger, 1898
		<i>Gnathonemus</i>	<i>Gnathonemus petersii</i> (Günther, 1862)
		<i>Hippopotamyrus</i>	<i>Hippopotamyrus weeksii</i> (Boulenger, 1902)
		<i>Marcusenius</i>	<i>Marcusenius greshoffi</i> (Schilthuis, 1891)
			<i>Marcusenius kutuensis</i> (Boulenger, 1899)
			<i>Marcusenius monteiri</i> (Günther, 1873)
			<i>Marcusenius moorii</i> (Günther, 1867)
			<i>Marcusenius</i> sp.
			<i>Marcusenius</i> sp.2
		<i>Mormyrops</i>	<i>Mormyrops anguilloides</i> (Linnaeus, 1758)
			<i>Mormyrops nigricans</i> Boulenger, 1899

Order	Family	Genus	Species	
Perciformes		<i>Mormyrus</i>	<i>Mormyrus proboscirostris</i>	
		<i>Myomyrus</i>	<i>Myomyrus macrops</i> Boulenger, 1974	
		<i>Oxymormyrus</i>	<i>Oxymormyrus boulengeri</i>	
		<i>Petrocephalus</i>	<i>Petrocephalus</i> cfr <i>binotatus</i>	
			<i>Petrocephalus</i> cfr <i>boboto</i>	
			<i>Petrocephalus</i> cfr <i>sauvagii</i>	
			<i>Petrocephalus</i> cfr <i>sauvagii</i> 2	
			<i>Petrocephalus</i> cfr <i>valentini</i>	
			<i>Petrocephalus christyi</i> Boulenger, 1920	
			<i>Petrocephalus sauvagii</i> (Boulenger, 1887)	
			<i>Pollimyrus</i>	<i>Pollimyrus osborni</i> (Nichols & Grisom, 1917)
			<i>Pollimyrus</i> sp.	
			<i>Stomathorinus</i>	<i>Stomathorinus corneti</i> Boulenger, 1899
			<i>Stomathorinus kununguensis</i> Poll, 1945	
	<i>Xenomystus nigri</i> (Günther, 1868)			
	<i>Pantodon buchholzi</i> Peters, 1877			
	<i>Ctenopoma</i> cfr <i>acutirostre</i>			
	<i>Ctenopoma kingsleyae</i> Günther, 1896			
	<i>Ctenopoma ocellatum</i> Pelligrin, 1899			
	<i>Ctenopoma weeksii</i> Boulenger, 1896			
	<i>Parachanna obscura</i> (Günther, 1861)			
	<i>Hemichromis fasciatus</i>			
	<i>Lamprologus mocquardi</i> Pellegrin, 1903			
	<i>Tylochromis lateralis</i> (Boulenger, 1898)			
	Unknown			
Polypteriformes	Polypteridae	<i>Polypterus</i>	<i>Polypterus delhezi</i> Boulenger, 1899	
			<i>Polypterus polli</i> Gosse, 1988	
Siluriformes	Bagridae	<i>Bagrus</i>	<i>Bagrus ubangensis</i> Boulenger, 1902	
	Clariidae	<i>Clarias</i>	<i>Clarias buthupogon</i> Sauvage, 1879	
			<i>Clarias camerunensis</i> Lönnberg, 1895	
			<i>Clarias pachynema</i> Boulenger, 1903	
			<i>Clarias gariepinus</i> (Burchell, 1822)	
			<i>Clarias</i> sp.	
		<i>Heterobranchus</i>	<i>Heterobranchus longifilis</i> Valenciennes, 1840	
	Claroteidae	<i>Auchenoglanis</i>	<i>Auchenoglanis occidentalis</i> (Valenciennes, 1840)	
			<i>Auchenoglanis</i> cfr <i>occidentalis</i>	

Order	Family	Genus	Species
		<i>Chrysichthys</i>	<i>Chrysichthys brevibarbis</i> (Boulenger, 1899) <i>Chrysichthys habereri</i> Steindachner, 1912 <i>Chrysichthys longipinnis</i> (Boulenger, 1899)
	Malapteruridae	<i>Malapterurus</i>	<i>Malapterurus microstoma</i> Poll & Gosse, 1969
	Mochokidae	<i>Synodontis</i>	<i>Synodontis alberti</i> Schilthuis, 1891 <i>Synodontis angelicus</i> Schilthuis, 1891 <i>Synodontis batesii</i> Boulenger, 1907 <i>Synodontis</i> cfr <i>pleurops</i> <i>Synodontis congicus</i> Poll, 1971 <i>Synodontis contractus</i> Vinciguerra, 1928 <i>Synodontis decorus</i> Boulenger, 1899 <i>Synodontis flavitaenitus</i> Boulenger, 1919 <i>Synodontis greshoffi</i> Schilthuis, 1891 <i>Synodontis nigriventis</i> David, 1936 <i>Synodontis notatus</i> Vaillant, 1893 <i>Synodontis pleurops</i> Boulenger, 1899 <i>Synodontis schoutedeni</i> David, 1936 <i>Synodontis</i> sp.
	Schilbeidae	<i>Pareutropius</i> <i>Parailia</i> <i>Schilbe</i>	<i>Pareutropius debauwi</i> (Boulenger, 1900) <i>Parailia congica</i> Boulenger, 1899 <i>Schilbe congensis</i> (Leach, 1818) <i>Schilbe grenfelli</i> (Boulenger, 1900) <i>Schilbe intermedius</i> Rüppell, 1832 <i>Schilbe laticeps</i> (Boulenger, 1899) <i>Schilbe marmoratus</i> Boulenger, 1911 <i>Schilbe yangambianus</i> (Poll, 1954)

Table 2. List showing the presence (+) or absence (-) of fish species captured in thirteen sampling rivers

Fish species	Sampling rivers												
	Amboko	Baleke	Emame	Lodja	Loidjo	Lomami	Nvul'elongo	Nyamatende	Onema	Petchi	Shingi	Wele	Yalo
<i>Alestes liebrechtsii</i>	+	-	-	+	-	-	-	-	-	-	-	-	+
<i>Alestes macrophthalmus</i>	-	-	-	-	-	-	-	-	-	-	-	-	+
<i>Alestopetersius brichardi</i>	+	+	-	+	-	-	+	+	+	+	+	-	+
<i>Alestopetersius</i> cfr <i>brichardi</i>	-	+	-	-	-	-	-	-	-	-	+	-	-
<i>Alestopetersius compressus</i>	-	+	-	+	+	-	-	+	+	+	+	-	-
<i>Auchenoglanis</i> cfr <i>occidentalis</i>	-	-	-	-	-	+	-	-	-	-	-	-	-
<i>Auchenoglanis occidentalis</i>	+	-	-	-	-	+	+	-	-	-	-	-	+
<i>Bagrus ubangensis</i>	-	-	-	-	-	+	-	-	-	-	-	-	-
<i>Bathyaethiops greeni</i>	-	-	-	-	+	-	-	+	-	-	-	-	-
<i>Brachypetersius altus</i>	+	-	-	+	+	+	+	+	+	+	+	-	+
<i>Brachypetersius huloti</i>	+	-	-	+	-	-	-	-	-	-	-	-	+
<i>Brachypetersius pseudonummifer</i>	-	-	-	-	+	-	-	+	-	-	-	-	-
<i>Brycinus bimaculatus</i>	+	-	-	+	+	-	-	+	-	-	-	-	+
<i>Brycinus</i> cfr <i>bimaculatus</i>	-	-	-	-	+	-	-	-	-	-	-	-	-
<i>Brycinus</i> cfr <i>grandisquamis</i>	+	-	-	-	-	+	-	-	-	-	-	-	-
<i>Brycinus</i> cfr <i>macrolepidotus</i>	+	-	-	-	+	-	-	-	-	-	-	-	-
<i>Brycinus</i> cfr <i>macrolepidotus</i> 2	+	-	-	+	-	-	-	-	-	-	-	-	+
<i>Brycinus grandisquamis</i>	+	-	-	+	-	+	-	-	-	-	-	-	+
<i>Brycinus imberi</i>	+	-	-	+	-	+	-	-	-	+	+	+	+
<i>Bryconaethiops boulengeri</i>	+	-	-	+	+	+	+	-	-	-	-	-	-
<i>Bryconaethiops macrops</i>	+	-	-	-	-	+	-	-	-	-	-	-	-
<i>Campylomormyrus elephas</i>	-	-	-	-	-	+	-	-	-	-	-	-	-
<i>Campylomormyrus numenius</i>	-	-	-	+	-	-	-	-	-	-	-	-	-
<i>Chrysichthys brevibarbis</i>	-	-	-	-	-	+	-	-	-	-	-	-	-
<i>Chrysichthys habereri</i>	-	-	-	-	-	+	-	-	-	-	-	-	-
<i>Chrysichthys longipinnis</i>	+	-	-	-	-	+	-	-	-	-	-	-	+
<i>Citharinus gibbosus</i>	+	-	-	+	-	+	-	-	-	-	-	-	+
<i>Citharinus macrolepis</i>	-	-	-	+	-	+	-	-	-	-	-	-	+
<i>Clarias buthupogon</i>	-	-	-	-	+	-	-	-	-	-	-	-	-
<i>Clarias camerunensis</i>	-	-	-	-	+	-	+	-	+	-	+	-	+
<i>Clarias gariepinus</i>	-	-	-	-	-	-	+	-	+	-	-	-	-
<i>Clarias pachynema</i>	-	-	-	-	-	-	-	-	-	+	-	-	-

Fish species	Sampling rivers												
	Amboko	Baleke	Emame	Lodja	Loidjo	Lomami	Nvul'elongo	Nyamatende	Onema	Petchi	Shingi	Wele	Yalo
<i>Clarias</i> sp.	-	-	-	-	-	-	+	-	-	-	-	-	-
<i>Ctenopoma cfr acutirostre</i>	+	-	-	-	-	-	-	-	-	-	-	-	-
<i>Ctenopoma kingsleyae</i>	-	-	-	-	+	-	-	-	-	-	+	-	+
<i>Ctenopoma ocellatum</i>	+	-	-	+	-	-	-	-	-	-	-	-	+
<i>Ctenopoma weeksii</i>	-	-	-	+	-	-	-	-	-	-	-	-	+
<i>Cyphomyrus psittacus</i>	+	-	-	+	-	+	-	-	-	-	-	-	+
<i>Cyphomyrus</i> sp.	+	-	-	-	-	-	-	-	-	-	-	-	-
<i>Cyphomyrus wilverthi</i>	+	-	-	-	-	-	-	-	-	-	-	-	-
<i>Distichodus affinis</i>	+	-	-	+	-	+	-	-	-	-	-	-	+
<i>Distichodus antonii</i>	+	-	-	+	-	+	-	-	-	-	-	-	+
<i>Distichodus atroventralis</i>	-	-	-	-	-	+	-	-	-	-	-	-	-
<i>Distichodus fasciolatus</i>	+	-	-	+	-	+	-	-	-	-	-	-	+
<i>Distichodus lusosso</i>	-	-	-	+	-	+	-	-	-	-	-	-	+
<i>Distichodus noboli</i>	-	-	-	+	-	-	-	-	-	-	-	-	-
<i>Distichodus sexefasciatus</i>	-	-	-	+	-	+	-	-	-	-	-	-	-
<i>Distichodus cfr fasciolatus</i>	-	-	-	-	-	+	-	-	-	-	-	-	-
<i>Enteromius miolepis</i>	-	+	+	+	+	-	+	+	+	+	+	-	-
<i>Eugnathichthys macroterolepis</i>	+	-	-	+	-	+	-	-	-	-	-	-	+
<i>Genyomyrus donnyi</i>	+	-	-	+	-	-	+	-	-	-	-	-	+
<i>Gnathonemus petersii</i>	+	+	+	+	-	+	+	-	-	+	+	-	+
<i>Hemichromis fasciatus</i>	+	+	+	+	+	-	+	+	+	+	+	+	+
<i>Hepsetus</i> sp.	-	+	+	+	+	+	+	+	+	+	+	+	+
<i>Heterobranchus longifilis</i>	-	-	-	-	-	-	+	-	-	-	-	-	-
<i>Heterotis niloticus</i>	-	-	-	-	-	-	+	-	-	-	-	-	-
<i>Hippopotamyrus weeksii</i>	-	-	-	+	-	-	-	-	-	-	-	-	-
<i>Hydrocynus forskahlii</i>	-	-	-	+	-	+	-	-	-	-	-	-	-
<i>Hydrocynus vittatus</i>	+	-	-	+	-	+	-	-	-	-	-	-	-
Unknown	-	-	-	-	-	-	+	-	-	-	-	-	-
<i>Labeo cyclorhynchus</i>	-	-	-	-	-	+	-	-	-	-	-	-	-
<i>Labeo greenii</i>	-	-	-	-	-	+	-	-	-	-	-	-	-
<i>Labeo lineatus</i>	-	-	-	-	-	+	-	-	-	-	-	-	-
<i>Labeo parvus</i>	-	-	-	-	-	+	-	-	-	-	-	-	-
<i>Labeo rectipinnis</i>	-	-	-	-	-	+	+	-	-	-	-	-	-
<i>Labeo</i> sp.	-	-	-	-	-	-	+	-	-	-	-	-	-
<i>Labeo weeksii</i>	-	-	-	-	-	+	-	-	-	-	-	-	+

Fish species	Sampling rivers												
	Amboko	Baleke	Emame	Lodja	Loidjo	Lomami	Nvul'elongo	Nyamatende	Onema	Petchi	Shingi	Wele	Yalo
<i>Labeobarbus caudovittatus</i>	-	-	-	-	-	-	+	-	-	-	-	-	-
<i>Lamprologus mocquardi</i>	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Malapterurus microstoma</i>	+	-	-	+	-	-	-	-	-	-	-	-	-
<i>Marcusenius greshoffi</i>	+	-	-	+	-	+	-	-	-	-	-	+	+
<i>Marcusenius kutuensis</i>	+	-	+	+	+	+	+	-	-	-	+	+	+
<i>Marcusenius monteiri</i>	+	-	-	+	-	+	-	-	-	-	-	+	+
<i>Marcusenius moorii</i>	+	-	+	+	+	+	+	+	+	+	+	-	+
<i>Marcusenius sp.</i>	-	+	-	+	-	+	-	+	-	+	-	-	-
<i>Marcusenius sp.2</i>	+	+	+	+	+	+	+	+	+	+	+	-	+
<i>Mesoborus crocodilus</i>	+	-	-	+	-	-	-	-	-	-	-	-	-
<i>Micralestes acutidens</i>	+	-	-	+	-	+	-	-	-	-	-	+	+
<i>Micralestes cfr sardina</i>	+	-	-	-	-	-	-	-	-	-	-	-	-
<i>Mormyrops anguilloides</i>	+	-	-	+	-	-	-	-	-	-	-	-	+
<i>Mormyrops nigricans</i>	-	-	-	+	-	-	-	-	-	-	-	-	+
<i>Mormyrus probosciostris</i>	-	-	-	-	-	+	-	-	-	-	-	-	-
<i>Myomyrus macrops</i>	-	-	-	-	-	+	-	-	-	-	-	-	-
<i>Nannothrissa stewarti</i>	+	-	-	-	-	+	-	-	-	-	-	-	+
<i>Oxymormyrus boulengeri</i>	-	-	-	-	+	-	-	-	-	-	-	-	-
<i>Pantodon buchholzi</i>	-	+	+	+	+	-	-	-	-	+	-	-	-
<i>Parachanna obscura</i>	-	-	-	-	-	-	-	-	-	-	-	-	+
<i>Pareutropius debauwi</i>	-	+	-	+	+	-	-	+	-	+	+	-	-
<i>Parailia congica</i>	+	-	-	-	-	-	-	-	-	-	-	-	+
<i>Petrocephalus cfr binotatus</i>	+	-	-	+	-	-	-	-	-	-	-	-	+
<i>Petrocephalus cfr boboto</i>	+	-	-	-	+	+	-	+	-	-	-	-	+
<i>Petrocephalus cfr sauvagii</i>	-	-	-	-	+	+	-	+	-	-	-	-	-
<i>Petrocephalus cfr valentini</i>	-	-	+	+	-	-	-	+	-	-	+	-	-
<i>Petrocephalus christyi</i>	+	-	-	+	+	+	-	+	-	-	+	+	+
<i>Petrocephalus sauvagii</i>	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Petrocephalus cfr sauvagii 2</i>	-	-	-	-	-	+	-	-	-	-	-	-	-
<i>Phago boulengeri</i>	+	-	-	+	-	-	-	-	-	-	-	-	+
<i>Phago intermedius</i>	+	-	-	-	-	+	-	-	-	-	-	-	+
<i>Phenacogrammus polli</i>	+	+	-	-	-	+	+	-	+	+	+	-	-
<i>Pollimyrus osborni</i>	+	+	+	+	+	+	-	+	+	+	+	-	+
<i>Pollimyrus sp.</i>	-	-	-	-	-	+	-	-	-	-	-	-	-
<i>Polypterus delhezi</i>	-	-	-	-	+	-	-	-	-	-	+	-	-

Fish species	Sampling rivers												
	Amboko	Baleke	Emame	Lodja	Loidjo	Lomami	Nvul'elongo	Nyamatende	Onema	Petchi	Shingi	Wele	Yalo
<i>Polypterus polli</i>	+	-	+	-	-	-	-	-	-	-	-	-	+
<i>Raiamas buchholzi</i>	+	-	-	+	-	+	-	-	-	-	-	-	+
<i>Schilbe congensis</i>	-	+	-	-	-	-	-	-	+	-	-	-	-
<i>Schilbe grenfelli</i>	-	-	-	-	-	+	-	-	+	+	-	-	-
<i>Schilbe intermedius</i>	+	+	-	+	-	+	-	-	-	-	-	-	+
<i>Schilbe laticeps</i>	-	+	-	+	+	-	+	+	-	-	+	-	-
<i>Schilbe marmoratus</i>	+	+	-	+	+	+	+	-	-	+	-	+	+
<i>Schilbe yangambianus</i>	-	+	+	+	+	-	+	+	+	+	+	-	-
<i>Stomathorinus corneti</i>	-	-	-	-	-	-	-	-	-	-	+	-	-
<i>Stomathorinus kununguensis</i>	-	-	-	-	+	-	-	-	-	-	-	-	-
<i>Synodontis alberti</i>	+	-	-	-	-	-	-	-	-	-	-	-	+
<i>Synodontis angelicus</i>	+	-	-	-	-	-	-	-	-	-	-	-	+
<i>Synodontis batesii</i>	-	-	-	-	-	+	-	-	-	-	-	-	-
<i>Synodontis cfr pleurops</i>	-	-	-	-	-	+	-	-	-	-	-	-	-
<i>Synodontis congicus</i>	+	-	-	+	-	-	-	-	-	-	-	-	+
<i>Synodontis contractus</i>	-	-	+	-	+	-	-	+	-	-	+	-	-
<i>Synodontis decorus</i>	+	-	-	-	-	+	-	-	-	-	-	-	-
<i>Synodontis flavitaenitus</i>	+	-	-	-	-	-	-	-	-	-	-	-	-
<i>Synodontis greshoffi</i>	+	+	+	+	+	+	+	+	+	-	-	+	+
<i>Synodontis nigriventis</i>	-	-	-	-	+	-	-	-	-	-	-	-	-
<i>Synodontis notatus</i>	-	-	-	-	-	+	-	-	-	-	-	-	-
<i>Synodontis pleurops</i>	+	-	-	-	-	+	-	-	-	-	-	-	+
<i>Synodontis schoutedeni</i>	-	-	-	-	-	-	-	-	-	-	-	-	+
<i>Synodontis sp.</i>	-	-	+	-	-	-	-	-	-	-	-	-	-
<i>Tylochromis lateralis</i>	+	-	-	+	-	+	-	-	-	-	-	-	-
<i>Xenocharax spilurus</i>	+	-	-	+	+	+	-	-	-	-	-	-	+
<i>Xenomystus nigri</i>	+	-	+	+	+	+	-	+	-	-	+	-	+

The family Mormyridae, endemic to sub-Saharan Africa, dominates in African rivers in terms of specific diversity [30,31,32,33] and here more particularly in the upstream part of the small tributary rivers of the Congo River [27]. Some can measure up to 1.5 m in length although the majority are between 9 and 50 cm in size [27]. These fish are often found together, forming groups in the turbid waters of rivers. This behavior and adaptation are probably favored by the very particular physiology of their sensory organs [27,32].

Indeed, the dominance of the Mormyriiform fishes is not coincidental, as they are also the largely dominant order in Africa [34]. Our findings agreed with those found by Mbega [35] in the lower Ogooué basin. Note that the families of fishes of the Lomami National Park and its hinterlands are among the families with a wide distribution in the freshwaters of Africa [36], including the families of Clariidae, Schilbeidae, Bagridae, Claroteidae, Mormyridae, Cichlidae, Mastacembellidae, Cyprinidae, Alestidae and Distichodontidae [27].

The Fig. 3c & d displayed that in the Lomami National Park and its hinterlands, all rivers are diverse with a minimum observed in the Onema River. This could be due to a low catching effort in this river. The probability of catching two individuals of different species in the same river is high in all rivers, but it is higher in the Amboko,

Lodja, Lomami and Yalo Rivers (96%); and it is slightly low in the Onema River (73%).

The Table 3 shows that the majority of species was least concerned (74.42%), but reports that one was endangered species (*Nannothrissa stewarti*) and another vulnerable (*Labeo rectipinnis*). The results of Micha et al. [37]. showed the overfishing of *Nannothrissa stewarti* in the Lake Mai-Ndombe was due to the use of the purse beach seine with mosquito nets, a prohibited fishing gear, which leads to the scarcity and disappearance of *Nannothrissa stewarti* of that lake.

The Table 4 indicates that nine taxa were identified as fish with high economic due to their preference by local populations. A total of 23 riparian fishermen were interviewed; 78.26% of them said that the main destinations of fishery resources from LNP were Kindu and Lodja cities located in Democratic Republic of the Congo. While 21.74% reported that fish caught were locally sold and consumed. Local fishermen of the LNP and its hinterlands reported that the main fishing materials used were gill nets of 10 mm, 15 mm, 20 mm, 25 mm, 30 mm, 40 mm, 60 mm and 100 mm of mesh; cast nets of 10 mm, 20 mm and 30 mm of mesh; hooks (N° 1, 2, 4, 6, 8, 10, 14 and 16); hand-line; traps; and planked canoes. 95.65% of riparian fishermen were the owners of these fishing gears used in the LNP and its hinterlands.

Table 3. Conservation status of species red list category of IUCN (International Union for Conservation of Nature)

No	Species	IUCN Status
1	<i>Alestes liebrechtsii</i>	LC
2	<i>Alestes macrophthalmus</i>	LC
3	<i>Alestopetersius brichardi</i>	LC
4	<i>Alestopetersius cfr brichardi</i>	-
5	<i>Alestopetersius compressus</i>	LC
6	<i>Auchenoglanis cfr occidentalis</i>	-
7	<i>Auchenoglanis occidentalis</i>	LC
8	<i>Bagrus ubangensis</i>	LC
9	<i>Bathyaethiops greeni</i>	LC
10	<i>Brachypetersius altus</i>	LC
11	<i>Brachypetersius huloti</i>	LC
12	<i>Brachypetersius pseudonummifer</i>	LC
13	<i>Brycinus bimaculatus</i>	LC
14	<i>Brycinus cfr bimaculatus</i>	-
15	<i>Brycinus cfr grandisquamis</i>	-
16	<i>Brycinus cfr macrolepidotus</i>	-
17	<i>Brycinus cfr macrolepidotus 2</i>	-
18	<i>Brycinus grandisquamis</i>	LC
19	<i>Brycinus imberi</i>	LC

No	Species	IUCN Status
20	<i>Bryconaethiops boulengeri</i>	LC
21	<i>Bryconaethiops macrops</i>	LC
22	<i>Campylomormyrus elephas</i>	LC
23	<i>Campylomormyrus numenius</i>	LC
24	<i>Chrysichthys brevibarbis</i>	LC
25	<i>Chrysichthys habereri</i>	LC
26	<i>Chrysichthys longipinnis</i>	LC
27	<i>Citharinus gibbosus</i>	LC
28	<i>Citharinus macrolepis</i>	LC
29	<i>Clarias buthupogon</i>	LC
30	<i>Clarias camerunensis</i>	LC
31	<i>Clarias gariepinus</i>	LC
32	<i>Clarias pachynema</i>	LC
33	<i>Clarias</i> sp.	-
34	<i>Ctenopoma cfr acutirostre</i>	-
35	<i>Ctenopoma kingsleyae</i>	LC
36	<i>Ctenopoma ocellatum</i>	LC
37	<i>Ctenopoma weeksii</i>	LC
38	<i>Cyphomyrus psittacus</i>	LC
39	<i>Cyphomyrus</i> sp.	-
40	<i>Cyphomyrus wilverthi</i>	LC
41	<i>Distichodus affinis</i>	LC
42	<i>Distichodus antonii</i>	LC
43	<i>Distichodus atroventralis</i>	LC
44	<i>Distichodus fasciolatus</i>	LC
45	<i>Distichodus lusosso</i>	LC
46	<i>Distichodus noboli</i>	LC
47	<i>Distichodus sexefasciatus</i>	LC
48	<i>Distichodus cfr fasciolatus</i>	-
49	<i>Enteromius miolepis</i>	NE
50	<i>Eugnathichthys macroterolepis</i>	LC
51	<i>Genyomyrus donnyi</i>	LC
52	<i>Gnathonemus petersii</i>	LC
53	<i>Hemichromis fasciatus</i>	LC
54	<i>Hepsetus</i> sp.	-
55	<i>Heterobranchus longifilis</i>	LC
56	<i>Heterotis niloticus</i>	LC
57	<i>Hippopotamyrus weeksii</i>	NE
58	<i>Hydrocynus forskahlii</i>	LC
59	<i>Hydrocynus vittatus</i>	LC
60	Unknown	
61	<i>Labeo cyclorhynchus</i>	LC
62	<i>Labeo greenii</i>	LC
63	<i>Labeo lineatus</i>	LC
64	<i>Labeo parvus</i>	LC
65	<i>Labeo rectipinnis</i>	VU
66	<i>Labeo</i> sp.	-
67	<i>Labeo weeksii</i>	LC
68	<i>Labeobarbus caudovittatus</i>	LC
69	<i>Lamprologus mocquardi</i>	LC
70	<i>Malapterurus microstoma</i>	LC
71	<i>Marcusenius greshoffi</i>	LC
72	<i>Marcusenius kutuensis</i>	LC
73	<i>Marcusenius monteiri</i>	LC
74	<i>Marcusenius moorii</i>	LC
75	<i>Marcusenius</i> sp.	-

No	Species	IUCN Status
76	<i>Marcusenius</i> sp.2	-
77	<i>Mesoborus crocodilus</i>	LC
78	<i>Micralestes acutidens</i>	LC
79	<i>Micralestes</i> cfr <i>sardina</i>	-
80	<i>Mormyrops anguilloides</i>	LC
81	<i>Mormyrops nigricans</i>	LC
82	<i>Mormyrus proboscirostris</i>	NE
83	<i>Myomyrus macrops</i>	LC
84	<i>Nannothrissa stewarti</i>	EN
85	<i>Oxymormyrus bouleengeri</i>	LC
86	<i>Pantodon buchholzi</i>	LC
87	<i>Parachanna obscura</i>	LC
88	<i>Pareutropius debauwi</i>	LC
89	<i>Parailia congica</i>	LC
90	<i>Petrocephalus</i> cfr <i>binotatus</i>	-
91	<i>Petrocephalus</i> cfr <i>boboto</i>	-
92	<i>Petrocephalus</i> cfr <i>sauvagii</i>	-
93	<i>Petrocephalus</i> cfr <i>valentini</i>	-
94	<i>Petrocephalus christyi</i>	LC
95	<i>Petrocephalus sauvagii</i>	LC
96	<i>Petrocephalus</i> cfr <i>sauvagii</i> 2	-
97	<i>Phago bouleengeri</i>	LC
98	<i>Phago intermedius</i>	LC
99	<i>Phenacogrammus polli</i>	LC
100	<i>Pollimyrus osborni</i>	NE
101	<i>Pollimyrus</i> sp.	-
102	<i>Polypterus delhezi</i>	LC
103	<i>Polypterus polli</i>	LC
104	<i>Raiamas buchholzi</i>	LC
105	<i>Schilbe congensis</i>	LC
106	<i>Schilbe grenfelli</i>	LC
107	<i>Schilbe intermedius</i>	LC
108	<i>Schilbe laticeps</i>	LC
109	<i>Schilbe marmoratus</i>	LC
110	<i>Schilbe yangambianus</i>	LC
111	<i>Stomathorinus corneti</i>	NE
112	<i>Stomathorinus kununguensis</i>	NE
113	<i>Synodontis alberti</i>	LC
114	<i>Synodontis angelicus</i>	LC
115	<i>Synodontis batesii</i>	LC
116	<i>Synodontis</i> cfr <i>pleurops</i>	-
117	<i>Synodontis congicus</i>	LC
118	<i>Synodontis contractus</i>	LC
119	<i>Synodontis decorus</i>	LC
120	<i>Synodontis flavitaenitus</i>	LC
121	<i>Synodontis greshoffi</i>	LC
122	<i>Synodontis nigriventis</i>	LC
123	<i>Synodontis notatus</i>	LC
124	<i>Synodontis pleurops</i>	LC
125	<i>Synodontis schoutedeni</i>	LC
126	<i>Synodontis</i> sp.	-
127	<i>Tylochromis lateralis</i>	LC
128	<i>Xenocharax spilurus</i>	NE
129	<i>Xenomystus nigri</i>	LC

LC = Least Concern, EN = Endangered, NE = Not Evaluated, VU = Vulnerable [<https://www.iucnredlist.org/>]

Table 4. Scientific and local names of the most preferred fish species

No	Scientific name	Local name
1	<i>Clarias</i> spp.	Samba, Kambale, Ngolo
2	<i>Schilbe</i> spp.	Mpendakula
3	<i>Chrysichthys</i> spp.	Kalimba, Nyuvi
4	<i>Auchenoglanis occidentalis</i>	Feke
5	<i>Polypterus</i> spp.	Mokonga ou Mukunga
6	<i>Hydrocynus</i> spp.	Manda
7	<i>Parachanna</i> spp.	Mongusu, Singa
8	<i>Alestes</i> spp.	Pungululu, Mandekibere
9	Species of Mormyridae	Mipoto, Tale, Pono

4. CONCLUSION

This study was mainly focused on ichthyofauna of Lomami National Park and its hinterlands. The findings showed that 129 fish species were caught during this investigation in the southern part of Lomami National Park. Mormyridae dominated in the collection with 13 genera and 30 species followed by family of Alestidae with 9 genera and 23 species. There was a possibility to have new species in some genera, such as *Marcusenius*, *Brycinus* and *Petrocephalus*. It is reported that *Labeo rectipinnis* is vulnerable species and *Nannothrissa stewarti* is endangered species. Note that fish species with high economic value have been recorded, including *Clarias* spp., *Schilbe* spp., *Chrysichthys* spp., *Auchenoglanis occidentalis*, *Polypterus* spp., *Hydrocynus* spp., *Alestes* spp., *Parachanna* spp. and species of Mormyridae.

81% of fishermen recorded reported that the number of fishermen is currently increasing. This leads to the decline of fish caught. However fish has long been a major source of livelihood for most riparian communities. From the results stressed above of this study, we recommend to promote integrated fish farming in the villages surroundings the Lomami National Park in order to increase food security and reduce the pressure exerted by the fishermen on fishery resources of LNP and its hinterlands; study the biology and ecology of fish species with high economic value identified in this survey in view of their possible domestication in the fish farming. This will enable the fish farming to be sustainable; and conduct further monitoring on vulnerable and endangered species.

ACKNOWLEDGEMENT

This work was supported by a research fund from U.S. Fish and Wildlife Service, through

Lukuru Foundation. We would like to express our gratitude to the staff members of USFWS and Lukuru Foundation. Our thankful feelings are also addressed to the staff members of ICCN/Maniema who provided us with guards we needed during fieldwork. We sincerely thank the researchers of lab of Hydrobiology and Aquaculture, University of Kisangani for completing this study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Maître-Allain T. The new aquarium manual. Paris, France, Solar; 1992. English.
2. Edison VGD. The status of the conch fishery of the Bahamas. ACP-EU Fisheries Research Report; 1998.
3. Thieme ML, Abell R, Stiassny MLJ, Skelton P, Lehner B, Teugels GG, Dinnerstein E, Toham AK, Burgess N, Olson D. Freshwater ecoregions of Africa and Madagascar: A conservation assessment. Island Press, Washington; 2005.
4. Darwall WRT, Smith KG, Allen DJ, Holland RA, Harrison IJ, Brooks EGE. The diversity of life in African freshwaters: underwater, under threat: An analysis of the status and distribution of freshwater species throughout mainland Africa. Cambridge, United Kingdom and Gland, Switzerland: IUCN; 2011.
5. Anthony NM, Atteke C, Bruford MW, Dallmeier F, Freedman A, Hardy O, Ibrahim B, Jeffery KJ, Johnson M, Lahm SA, Lepengue N, Lowenstein JH, Maisels F, Mboumba J-F, Mickala P, Morgan K,

- Ntie S, Smith TB, Sullivan J-P, Verheyen E, Gonder MK. Evolution and conservation of Central African Biodiversity: Priorities for Future Research and Education in the Congo Basin and Gulf of Guinea. *Biotropica*. 2015;47(1):6–17.
6. Hart TB. Community drafting and acceptance of conservation zoning across the Tshuapa-Lomami-Lualaba (TL2) landscape. MID-Term Report, USFWS – Great Ape Conservation Fund, DRC; 2011.
 7. Boulenger GA. Fishes of the Congo Basin. Publication of the Congo Free State; 1901. English.
 8. Poll M. Non-cichlidae fish of Lake Tanganyika. Royal Institute of Natural Sciences of Belgium; 1953. English.
 9. Poll M. Classification of Lake Tanganyika Cichlidae: Tribes, Genera and Species. Torley-Rousseau Zoological Institute, Free University of Brussels; 1986. English.
 10. Matthes H. Fishes of Lake Tumba and the Ikela region: systematics and ecology. RMCA No. 126, Brussels; 1964. English.
 11. Kid JP. Fish from the Ubangui basin. *Annals of the Royal Museum for Central Africa. Zoological Science*. 1968;13:1-56. English.
 12. Roberts TR, Stewart DJ. An Ecological and Systematic survey of Fishes in rapids of the lower Zaire or Congo River. *Bulletin of the Museum of Comparative Zoology*. 1976;147(6):239-317.
 13. Banister KE, Bailey RG. Fishes collected by the Zaire River expedition 1974-1975. *Zoological Journal of the Linnean Society*. 1979;66(3):205-249.
 14. Teugels GG. A systematic outline of the African species of the genus *Clarias* (Pisces; Clariidae), with an annotated bibliography. *Annals of the Royal Museum for Central Africa*; 1982.
 15. Poll M, Gosse J-P. Genera of freshwater fish from Africa. Science class. Royal Academic of Belgium, in-8° collection, 3rd series. Volume IX;1995. English.
 16. Tshibwabwa S. Systematics of African species of the genus *Labeo* (Teleostei, Cyprinidae) in the ichthyological regions of Lower Guinea and Congo. Presses Universitaires de Namur Rampart of the Virgin, Belgium; 1997. English.
 17. Shumway C, Lévêque C, Paugy D, Teugels GG, Poll M, Gosse JP. Field guide to the fishes of the Democratic Republic of Congo, excluding Lake Tanganyika. Congo River Environment and Development Project (CREDP); 2002.
 18. Mbega JD, Teugels GG. Guide for determining the fish of the lower basin of the Ogooué. Ogooué, Gabon; 2003. English.
 19. Skelton PH. a complete guide to the freshwater fishes of Southern Africa. Grahamstown: Southern Book Publishers; 2004.
 20. Stiassny LJM, Teugels GG, Hopkins DC. Fresh and brackish water fish from Lower Guinea, West Central Africa. Research Institute for Development (IRD) (Paris, France), National Museum of Natural History (MNHN) (Paris, France) and Royal Museum for Central Africa (RMCA) (Tervuren, Belgium); 2007a. English.
 21. Stiassny LJM, Teugels GG, Hopkins DC. Fresh and brackish water fish from Lower Guinea, West Central Africa. Research Institute for Development (IRD) (Paris, France), National Museum of Natural History (MNHN) (Paris, France) and Royal Museum for Central Africa (RMCA) (Tervuren, Belgium); 2007b. English.
 22. AMNH lower Congo keys, FishBase. Worldwide web electronic publication; 2016. Available:www.fishbase.org.
 23. Sullivan JP, Friel JP, Kankonda BA, Tambwe EL, Tumitu JPU, Bulimwengu WA. Ichthyological study and collection-building at the University of Kisangani, Orientale, D.R. Congo. PAFFA Bujumbura; 2013.
 24. Mulangu K. Contribution to the ecological study and fish biodiversity of the Lubilanj River (Kasaï Oriental/ DR Congo). Doctoral thesis, Department of Zootechnics, Faculty of Agronomic Sciences, Official University of Mbuimayi; 2011. English.
 25. Tumitu J-PU, Mambo T. The fish fauna of the Lomami River and its tributaries, as well as a section of the Congo River from Kindu to Ubundu. Report, University of Kisangani; 2012. English.
 26. Kisekelwa T, Hyangya L, Masilya MP, Isumbisho M, Kaningini M. Contribution to the systematic inventory of fish in the Lowa River in Walikale territory. Cahiers du CERUKI, Special Issue Alphonse Byamungu; 2014. English.
 27. Byanikiro MR, Ndjaki NJ, Kankonda BA, Ulyel AJ, Micha JC. Characterization and ecology of mormyridae populations in the

- Yoko reserve (Kisangani, DR Congo). International Journal of Biological and Chemical Sciences. 2017;11(3):967-999. English.
28. Decru E, Vreven E, Danadu C, Walanga A, Mambo T, Snoeks J. Ichthyofauna of the Itimbiri, Aruwimi, and Lindi/Tshopo rivers (Congo basin): Diversity and distribution patterns. Acta Ichthyologica and Piscatoria. 2017;47(3):225–247.
29. Kisekelwa T, Snoeks J, Vreven E. An annotated checklist of the fish fauna of the river systems draining the Kahuzi-Biega National Park (Upper Congo: Eastern DR Congo). Journal of Fish Biology. 2020; 96:700–72.
30. Lalèyè P, Chikou A, Philippart J-C, Teugels G, Vandewalle P. Study of the ichthyological diversity of the Ouémé river basin in Benin (West Africa). Cybium. 2004;28(4):329-339. English.
31. Nelson JS. Fishes of the world. John Wiley and Sons, Inc. New York. 3rd edition; 2006.
32. Kramer B. Electric organ discharge. In: M.D. Binder N. Hirokawa U. Windhorst (Eds.). Encyclopedia of Neuroscience, Berlin Heidelberg: Springer Publishing House; 2009.
33. Aboua RDB, N'zikonan G, Kouamelan PE, Berte S, Bamba M. Spatial organization of fish population in Bandama. International Journal of Biological and Chemical Sciences. 2010;4(5): 1480–1493. English.
34. Lowe-McConnell RE. Ecological Studies in Tropical Communities. Cambridge University Press [PubMed]; 1987.
35. Mbega J. Fish biodiversity of the lower Ogooué basin. PhD thesis, FUNDP, Namur; 2003. English.
36. Greenwood PH. On *Macrolepodus*, *Chilotilapia* (Teleostei, Cichlidae), and the interrelationships of African cichlid species flocks. Bulletin of the British Museum (Natural History) Zoology [PubMed]. 1983;45(4):209-2 English.
37. Micha JC, Bangulu NB-L, Ibofa R, Mumba F, Mutambwe S, Zanga N, Willem E, Svennsson JE, Wilander A. An overexploited resource, *Nannothrissa stewarti*, the endemic sardine of Lake Maï-Ndombe (DR Congo), AN unexpected outcome of the National Malaria Control Programme. Report ; 2019. English.

© 2023 Shabani et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/104110>