



Performance of *Gladiolus* Genotypes for Growth and Flowering under Gangetic Region of Bihar, India

Kajal Kumari ^a, Paramveer Singh ^{a*}, Deepti Singh ^a,
Mainak Ghosh ^b, Subrat Keshori Behera ^c,
Ajay Bhardwaj ^a and R. B. Verma ^a

^a Department of Horticulture (Veg. and Flori.), Bihar Agricultural College, Bihar Agricultural University, Sabour-813210, Bhagalpur, Bihar, India.

^b Department of Agronomy, Bihar Agricultural College, Bihar Agricultural University, Sabour-813210, Bhagalpur, Bihar, India.

^c Department of S.M.C.A., BAC, Sabour, Bihar Agricultural College, Bihar Agricultural University, Sabour-813210, Bhagalpur, Bihar, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/v13i102938

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/105357>

Original Research Article

Received: 25/06/2023

Accepted: 31/08/2023

Published: 09/09/2023

ABSTRACT

An evaluation experiment on performance of gladiolus genotypes was conducted at Botanical Garden, Bihar Agricultural University, Sabour during 2021-2022 for the identification of suitable genotype for the cultivation in Bihar. Twenty genotypes of gladiolus were evaluated for vegetative, floral and vase life parameters under Gangetic region of Bihar conditions. The best genotypes identified in present investigation based on mean performance were White Prosperity (79.73 cm), Red Majesty (79.27 cm), Nova Lux (73.47 cm) and Rose Supreme (73.40 cm) for spike length.

*Corresponding author: E-mail: shekhawatdeep@rediffmail.com;

Earliness in flowering was observed in Rose Supreme (83.53 days) which was at par with Punjab Dawn (83.60 days). Genotype Kum Kum responded well for number of spikes per plant (2.73), whereas Pusa Shubham for corms production (3.33). Genotype White Prosperity (30.27 days) and Rose Supreme (28.47 days) were recorded long flowering duration.

Keywords: *Gladiolus*; genotypes; spike length; florets; corm; cormels.

1. INTRODUCTION

Gladiolus is one of the most cultivated, economically important and common flowering plants. It is commonly known as Sword lily or queen of the bulbous flowers for its excellent aesthetic value and display life [1]. It is native to South Africa and presently has been cultivated throughout the world due to its attractive characteristics, dazzling colors, varying sizes and long vase life [2]. Its demand is very high among consumers and florist because of its fascinating spikes with variety of colours that make it attractive for use in as cut flower, making bouquets, indoor decoration and in landscaping as herbaceous borders, bedding plants, in rockeries and pots. It has the 8th rank of cut flowers and first rank of bulbous flowers in the world trade [3]. The major gladiolus growing states are West Bengal, Maharashtra, Uttar Pradesh, Himachal Pradesh and Uttarakhand. It is one of the dominating cut flowers in the flower market of Bihar, but its cultivation is very negligible in Bihar and most of the cut flower procured in Bihar from West Bengal. However, many climatic conditions like abundant sunshine and favourable temperature for its growth are available in Bihar. Thus, there is much scope for increase in cultivation of gladiolus in Bihar.

The first important step in any crop improvement programme is evaluation. It is the basic tool for assessing the genetic variability present in any crop species, which could be exploited for its commercialization. The performance of any crop or genotype mainly depends on genotypic and environmental interaction. As a result, genotypes, which perform well in one region, may not perform same in other regions of changing climatic conditions. Hence, it becomes very much necessary to study the morphological variation to find out suitable genotype for a particular region. Since there is no information available on the performance of gladiolus genotypes in Gangetic region of Bihar. Thus, the present evaluation experiment was conducted to study the relative performance of twenty genotypes of gladiolus for growth, flowering, quality and yield traits.

2. MATERIALS AND METHODS

The present experiment was carried out at Botanical Garden, Department of Horticulture (vegetable & floriculture), BAU, Sabour during 2021-2022. The experiment was laid out in a randomized block design with twenty genotypes with three replications. The corms were treated with carbendazim (0.2%) before planting in field. The corms were planted in beds at a spacing of 30 cm x 30 cm. The recommended package and practices i.e., irrigation, hoeing, weeding and fertilizer application was given in time during the entire crop period for obtaining better quality spikes along with high yield. Observations were recorded in five randomly selected plants in each genotype for various vegetative and floral parameters.

3. RESULTS AND DISCUSSION

The findings of the research work conducted during 2021-2022, on the evaluation of twenty gladiolus genotypes revealed that the vegetative, floral and corms production parameters of the plants were greatly influenced due to genotypes. Mean data (Table 1) indicated that among the genotypes, White Prosperity had shown maximum plant height (82.47 cm) and minimum plant height was observed in genotypes Kum Kum (50.00 cm). The variation in plant height in different genotypes may be due to influence of genetic and environmental factors. Similar observations for plant height also reported by Singh et al. [4], Singh et al. [5] and Mushtaq et al. [6] in gladiolus.

Number of leaves per plant was observed maximum in genotypes Pusa Subham (12.73) followed by genotypes Punjab Dawn (10.80) and Traderhorn (10.80), while minimum number of leaves per plant was observed in genotypes Green Spire (6.87). The variation in number of leaves per plant might be due to genetic variation and difference in adaptation to the agro climatic conditions [7]. Similar variations in number of leaves had been reported by Swaroop et al. [8] and Susila [9] in gladiolus.

Table 1. Performance of gladiolus genotypes for growth and flowering under Gangetic region of Bihar

Genotypes	Plant height (cm)	Number of leaves per plant	Width of longest leaf (cm)	Days required for first spike emergence	Spike length (cm) at first floret open stage
Applouse	66.50	10.13	3.79	73.27	68.40
Pink Lady	62.40	7.13	2.51	72.73	68.07
Summer Sunshine	62.93	7.33	2.85	71.00	69.13
White Prosperity	82.47	7.80	2.81	77.93	79.73
Rose Supreme	68.90	7.53	2.85	63.40	73.40
Red Majesty	66.00	7.67	2.80	71.20	79.27
American Beauty	66.10	7.80	2.66	69.27	68.73
Nova Lux	62.00	8.20	2.67	70.87	73.47
Green Spire	65.17	6.87	2.71	68.00	68.13
Pusa Subham	58.00	12.73	2.87	73.40	67.93
Darshan	52.80	8.33	2.54	68.40	61.20
Gunjan	69.97	7.47	2.75	75.87	68.53
Algarve	58.32	7.53	2.87	70.47	67.73
Arka Poonam	65.83	8.00	2.60	71.20	68.37
Green Star	65.38	7.73	2.59	74.40	63.27
Punjab Dawn	64.77	10.80	2.55	64.20	66.53
Traderhorn	57.27	10.80	2.68	68.73	70.53
Arka Gold	59.73	7.20	2.55	74.47	67.29
Kum Kum	50.00	10.60	2.74	77.00	60.39
MPUAT Glad-1	62.77	8.93	2.85	78.53	68.07
S. Em. ±	1.929	0.293	0.116	2.420	2.513
CD. at 5%	5.524	0.839	0.334	6.928	7.197
CV. %	6.830	7.087	8.115	5.845	6.318

Width of longest leaf was recorded maximum in genotypes Applouse (3.79 cm) followed by genotypes Pusa Subham (2.87 cm) and Algarve (2.87 cm) while width of longest leaf recorded minimum in genotypes Pink Lady (2.51 cm). The similar variation reported by Azimi [10].

Earliest spike emergence was exhibited by genotypes Rose Supreme (63.40 days) followed by Punjab Dawn (64.20 days) and genotype MPUAT Glad-1 took maximum days for first spike emergence (78.53 days). Time required for spike emergence is an important varietal character in gladiolus that might be primarily governed by the genetic makeup of the genotypes. Spike emergence might have been primarily dependent on food reserves in plant that could be related to growth rate of plants regulating accumulation of the requisite level of carbohydrates for slipping. Similar results on differences for spike emergence among different genotypes have been reported by Singh et al. [11].

Spike length at first floret open stage was recorded maximum in genotype White Prosperity (79.73 cm) followed by Red Majesty (79.27 cm) and genotype Kum Kum (60.39 cm) was

observed to minimum spikes length at first floret open stage. This variation in spike length might be attributed to the inherent genetic characters associated with the genotypes. Spike length and rachis length are the important commercial traits in gladiolus which decide the rate in the market and their acceptability [12].

Earliest days to complete opening of first floret was exhibited by genotypes Rose Supreme (63.40 days) followed by Punjab Dawn (64.20 days) and genotype MPUAT Glad-1 (78.53 days) took maximum days to complete opening of first floret followed by White Prosperity (79.73 days). The results indicated that there is a particular set of flowering periods for each genotype. Results are similar to the findings of Singh et al. [11] and Solanki et al. [13]. This trait is important in gladiolus, which designates early or late flowering of genotypes and will be helpful in the availability of flowers.

Number of florets per spike was found maximum in genotype White Prosperity (13.73) followed by MPUAT Glad-1 (13.27) and genotype Green Star

Table 2. Performance of gladiolus genotypes for growth and flowering under Gangetic region of Bihar

Genotypes	Days to complete opening of first floret	Number of florets per spike	Number of spikes per plant	Number of corms	Weight of corms per hill (gm)
Applouse	93.87	9.07	1.13	2.20	69.93
Pink Lady	92.53	11.33	1.00	1.40	48.10
Summer	91.67	12.20	1.20	1.20	57.60
Sunshine					
White	96.60	13.73	1.40	1.60	66.50
Prosperity					
Rose Supreme	83.53	11.53	1.40	1.07	54.70
Red Majesty	92.00	12.13	1.47	1.80	76.30
American	90.73	11.73	1.47	1.87	60.63
Beauty					
Nova Lux	92.80	9.80	1.67	2.80	59.43
Green Spire	95.20	10.53	1.53	2.20	48.17
Pusa Subham	95.93	10.80	1.87	3.33	98.73
Darshan	96.00	9.47	1.53	2.00	32.18
Gunjan	92.67	10.20	1.40	1.67	41.70
Algarve	90.07	9.40	1.07	1.20	62.03
Arka Poonam	91.40	10.13	1.40	2.67	24.45
Green Star	92.87	8.47	1.40	1.60	46.70
Punjab Dawn	83.60	10.47	2.07	2.93	96.74
Traderhorn	87.20	12.00	1.00	1.87	42.83
Arka Gold	94.40	9.73	1.27	1.20	39.80
Kum Kum	94.80	8.73	2.73	2.93	45.20
MPUAT Glad-1	97.07	13.27	1.47	2.47	99.80
S. Em . ±	1.175	0.523	0.072	0.079	1.19
CD. at 5%	3.365	1.497	0.207	0.226	5.49
CV. %	2.200	8.440	8.519	6.863	5.67

(8.47) was observed to minimum number of florets per spike. Number of florets differs from genotypes to genotypes, this might be due to hereditary traits of the cultivar of the gladiolus, which is governed by genetic makeup of the plants. Similar results on number of florets per spike have been reported by Rani and Singh [14] and Solanki et al. [13] in gladiolus.

Among the genotypes maximum number of spikes per plant was recorded in Kum Kum (2.73) whereas, minimum was recorded in Pink Lady (1.00) and Traderhorn (1.00). Variation in yield of spikes is mainly attributed due to the variation in sprouting percentage of corms and sprouts per plant which are the genetically controlled character Ainarkar et al. [15]. Similar findings were also observed by Geeta et al. [16] and Pattanaik et al. [17].

Number of corms was found maximum in genotype Pusa Subham (3.33) followed by genotype Kum Kum (2.93) and genotype Rose Supreme (1.07) had minimum number of corms.

The variation observed in corm production amongst the genotypes studied might be due to differences in genetic as well as environmental factors. Similar result was also obtained by Singh et al. [11].

The results disclosed highly significant variations in weight of corms/plant among the genotypes. Maximum Weight of corms per hill was observed in genotype MPUAT Glad-1 (99.80 gm) followed by genotype Pusa Subham (98.73 gm), Punjab Dawn (96.74 gm) and minimum weight of corms per hill was observed in genotype Arka Poonam (24.45 gm). The variation observed in corm production amongst the genotypes studied might be due to differences in genetic as well as environmental factors. The weight of corm appeared to be associated with diameter of corm and number of cormels as evident from the results. It could be due to the fact the larger corms might have deposited more food resulting into their correspondingly heavier weight and vice versa. Similar findings were reported by Subhendu et al. [18] and Swetha et al. [19].

4. CONCLUSION

The research findings revealed that the different genotypes of gladiolus had the potential to influence the vegetative, flowering and yield traits of the plant. Among the twenty genotypes evaluated, Rose Supreme, White Prosperity, Pusa Shubham and Kum Kum were suitable for cultivation due to their flower quality as well as yield potential over other genotypes under Gangetic region of Bihar.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Bhattacharjee S, De LC. Post-harvest technology of flowers and ornamental plants. Aavishkar Publishers, Jaipur, India. 2005:11-19.
2. Farid UM, Rahman MM, Rabbani MG, Mannan MA. Effect of corm size and depth of planting on the growth and flowering of gladiolus. Pak. J. Biol. Sci. 2002;5(5):553–555.
3. Pragya JK, Ranjan BL, Attri B, Das HK, Ahmed, N. Performance of gladiolus genotypes for cut flower and corm production under high altitude of Uttarakh. Indian Journal of Horticulture. 2010;67: 386-390.
4. Singh AK, Sisodia A, Sisodia V, Ray P. Performance of Indian and exotic varieties of gladiolus under Eastern UP conditions. Journal of Ornamental Horticulture. 2017a; 20(3&4):153-157.
5. Singh D, Mishra A, Singh J, Meena B. Evaluation of morphological characters of Gladiolus (*Gladiolus hybridus* Hort.) genotypes under sub humid condition of Rajasthan. International Journal of Agriculture Sciences. 2017b;9(7):3846-3848.
6. Mushtaq S, Hafiz IA, Arif M, Anwar A. Performance evaluation of elite gladiolus cultivars under agro climatic conditions of Rawalpindi. Asian Journal of Advances in Agricultural Research. 2018;5(3):1-6.
7. Saleem M, Ahmad I, Khan MA. Cultivar effects on growth, yield and cormels production of gladiolus. Journal of Ornamental and Horticultural Plants. 2013; 3(1):39- 4.
8. Swaroop K, Krishan P, Singh KP. Performance of gladiolus under Delhi conditions. Journal of Ornamental Horticulture. 2005;8(1):32-35.
9. Susila T. Evaluation of gladiolus varieties for Visakhapatnam district of Andhra Pradesh, India. Agricultural Science Digest. 2013;33(3):237–238.
10. Azimi MH. Evaluation yield and genetically factors in different cultivars of Gladiolus. Ornamental Horticulture. 2019;26(1):8-17.
11. Singh AK, Kumar A, Ranjan R, Sinha BM, Singh MP. Performance of gladiolus genotypes under north Bihar Agro climatic conditions. International Journal of Chemical Studies. 2020;8(5):1733-1735.
12. Swaroop K, Singh KP, Kumar A. *Gladiolus* hybrids for commercial traits and their evaluation. International Journal of Current Microbiology and Applied Sciences. 2022; 11(06):335-343.
13. Solanki NS, Sen P, Singh L, Pawak R. Evaluation of different gladiolus (*Gladiolus grandiflorus*) cultivars in terms of spike quality under Allahabad agro climatic condition. International Journal of Current Microbiology and Applied Sciences. 2021; 10(09):384-390.
14. Rani R, Singh C. Evaluation of different gladiolus cultivars for quality flower production. Journal of Researches Birsa Agriculture University. 2005;17(2):227-230.
15. Ainarkar AA, Khanvilkar MH, Kulkarni MM, Dalvi NV, Shinde BD, Ainarkar AA, Mhatre MS. Evaluation of gladiolus (*Gladiolus spp.*) varieties for flowering characters and yield under Konkan ago climatic conditions of Maharashtra. The Pharma Innovation Journal. 2023;12(2):3159-3161.
16. Geeta SV, Shirol AM, Kulkarni BS, Omem T, Parvati P. Performance of gladiolus (*gladiolus hybridus* hort.) Varieties for growth, yield and flower quality characters. Plant Archives. 2014;14(2): 1147-1149.
17. Pattanaik S, Paul A, Lenka PC. Genotypic and phenotypic variability and correlation studies in gladiolus. Journal of Crop and Weed. 2015;11(1):113-119.
18. Subhendu GS, Suresh MK, Jayoti M. Genetic variability, character association and path coefficient analysis in gladiolus for various quantitative traits. Indian

- Journal of Horticulture. 2016;73(4):564-569.
19. Swetha S, Kulkarni BS, Shiragur M, Kulkarni MS, Mulge R, Naik LHMB. Genetic variability studies in gladiolus (*Gladiolus hybridus* Hort.). Journal of Pharmacognosy and Phytochemistry. 2020;9(1):726-731.

© 2023 Kumari et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/105357>