



# Effects of Salinity Stress on Growth Parameters of Soybeans (*Glycine max*) Cultivated in Obubra Local Government Area, of Cross River State Nigeria

Andrew Ashieta Aboh <sup>a\*</sup> and Eyong, Oduba Ikwa <sup>b</sup>

<sup>a</sup> Department of Agronomy, Cross River University of Technology, P.M.B. 1123, Cross River State, Nigeria.

<sup>b</sup> Department of Plant Science and Biotechnology, Cross River University of Technology, P.M.B. 1123, Calabar, Cross River, Nigeria.

## Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/AJRCS/2023/v8i4227

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/106734>

**Original Research Article**

**Received: 28/08/2023**

**Accepted: 02/11/2023**

**Published: 23/11/2023**

## ABSTRACT

Soybean (*Glycine max*) is a strategic crop plant grown to obtain edible oil and forage. High sensitivity is one of the biggest problems with soybean crop. A pot experiment was conducted in the screen house at the Cross River University of Technology, Obubra Campus to investigate the effects of different salt concentrations on the growth parameters of three different varieties of soybeans (*Glycine max*). The experiment was laid in complete randomized design (CRD) with twenty replicates for each variety. Growth parameters which include germination percentage, plant height, number of leaves and number of branches per plant were measured, and data collected was subjected to statistical analysis using Analysis of variance (ANOVA) and means were compared using Duncan's New Multiple Range Test (DNMRT). The results showed that salinity stress led to a significant reduction in percentage of germination, plant height, number of leaves, and number of branches per plant over control. At 30mM NaCl which was the highest level of saline water TGX

\*Corresponding author: E-mail: andyaboh@unicross.edu.ng;

1987-10F had a better germination percentage as compared to other varieties, which might be because of some salt tolerant genes. Plant height reduced significantly under salinity stress as compared with the control, but TGX 1835-10F at 30mM which was the highest saline treatment showed a better salt tolerance level as compared with other varieties. Number of leaves per plant, as the salinity levels increases it affected the number of leaves. There was a drastic reduction in the number of leaves with increasing salinity levels as compared with the control. TGX 1987-10F had the highest number of leaves and it was closely followed by TGX 1835-10E while 1448-2E was the least and there were significantly different from the ones at different salinity levels. The result revealed highly significant differences ( $p=0.05$ ) in the varieties, with TGX 1987-10F as the best variety appropriate for saline regions. It is therefore recommended that, TGX 1987-10F should be cultivated in saline regions since it showed a better tolerance level as compared to other varieties.

**Keywords:** Global scarcity of water resources; environmental pollution; agricultural sustainability; soybeans.

## 1. INTRODUCTION

“The beginning of 21<sup>st</sup> century is marked by global scarcity of water resources, environmental pollution and increased salinization of soil and water” [1]. “Increasing human population and reduction in land available for cultivation are two threats for agricultural sustainability” [2]. “Various environmental stresses viz. high winds, extreme temperatures, soil salinity, drought and flood have affected the production and cultivation of agricultural crops, among these, soil salinity is one of the most devastating environmental stresses, which causes major reductions in cultivated land area, crop productivity and quality” [2-4]. “A saline soil is generally defined as one in which the electrical conductivity (EC) of the saturation extract (EC<sub>e</sub>) in the root zone exceeds 4 dSm<sub>-1</sub> (approximately 40 mM NaCl) at 25 °C and has an exchangeable sodium of 15%. The yield of most crop plants is reduced at this EC<sub>e</sub>, though many crops exhibit yield reduction at lower EC<sub>e</sub>s” [5-9]. “It has been estimated that worldwide 20% of total cultivated and 33% of irrigated agricultural lands are afflicted by high salinity. Furthermore, the salinized areas are increasing at a rate of 10% annually for various reasons, including low precipitation, high surface evaporation, weathering of native rocks, irrigation with saline water, and poor cultural practices. It has been estimated that more than 50% of the arable land would be salinized by the year 2050” [10].

“Water and soil management practices have facilitated agricultural production on soil marginalized by salinity but an additional gain from these approaches seems problematic” [2]. “Impacted soils are a major limiting production factor worldwide for every major crop” [11,12].

“Soybean (*Glycine max* L.) is a strategic crop plant grown to obtain edible oil and forage” [13].

“High sensitivity to soil and water salinity is one of the biggest problems with soybean crop. Results have indicated that salinity affects growth and development of plants through osmotic and ionic stresses. Because of accumulated salts in soil under salt stress condition plant wilts apparently while soil salts such as Na<sup>+</sup> and Cl<sup>-</sup> disrupt normal growth and development of plant” [14,15]. “The agronomic traits of soybean could be severely affected by high salinity, including reduction in height, leaf size, biomass, number of internodes, number of branches, number of pods, weight per plant, and weight of 100 seeds” [16,17].

## 2. MATERIALS AND METHODS

A pot experiment was conducted in the screen house at the Cross River University of Technology, Obubra Campus to evaluate the effects of different salt concentrations on some growth parameters of three varieties of soybeans. Obubra is in central Senatorial area of Cross River State, it lies within latitude 6.0767200° and longitude 8.3324100°. It has an annual rainfall of about 2000- 2500mm with maximum temperature of about 21°-30°C [18]. The soil samples were collected from different experimental units at a depth of 0-30cm, they were bulked, air dried, sieved with 2mm mesh sieve and used for laboratory analysis for the pot experiment.

“The soils were analysed for particle size distribution using the hydrometer method” [19]. “Soil PH was determined in 1:2:5 soil to water ratio on a direct reading PH meter using the glass electrode with saturated potassium chloride calomel reference electrode. Organic carbon was determined by the wet oxidation method” as described by [20]. “The value of organic matter was obtained by multiplying the values for

organic carbon by 1.724. Exchangeable basic cations were extracted with K and Na leachate determined using EFL flame photometer and Ca and Mg by EDTA titration method, Exchangeable acidity" was by Mclean method [21]. "Effective cation exchange capacity (ECEC) was done by summation method. Available P was done by Bray P-2 method" [22].

The pot experiment was laid out in a Randomized Complete Design (CRD) with twenty replications. The treatments comprised of four levels of salt (0, 10, 20 and 30mM NaCl).

The seeds of the *Glycine max* varieties TGX1835-10E, TGX1987-10F and TGX 1448-2E were obtained from the International Institute of Tropical Agriculture (IITA), Ibadan Nigeria. 240 plastics pots of equal sizes were filled with 12kg of soil, four seeds of each variety were sown at the depth of 2cm and were later thinned down to two seedlings per pot at two weeks after planting (WAP). After planting, tap water (0.043mM) was applied to control pots and 2 dS/m NaCl-salinity irrigation water was added to the rest of the pots. When the first new leaf appeared, i.e., ten days after emergence (DAE), irrigation water with selected NaCl salinity (0, 10, 20 and 30mM) was applied, except for the control pots. Plants in the control groups were irrigated with tap water. The salt solution was applied daily until harvesting.

**Data collection:** The following parameters were measured.

Germination percentage = total number of plants germinated / total number of seeds sown

**Plant height:** The heights of the tagged plants was measured with measuring tape from the base to the top of the main axis and the mean values were recorded.

**Number of leaves:** The number of leaves of the tagged plants was determined by counting the numbers of fully expanded leaves and the mean values were recorded.

**Number of branches:** The number of branches of the tagged plants was determined by counting the branches and the mean values were recorded.

**Data analysis:** The data were analysed with SPSS version 20 software and all data collected were subjected to Analysis of Variance (ANOVA) according to Gomez and Gomez [23] and

treatments were compared using Duncan Multiple Test Range (DMRT) 1995.

### 3. RESULTS AND DISCUSSION

Table 1 showed a decrease in percentages of germination at different salinity levels as compared with the control. TGX 1448-2E had the highest germination percentage followed by TGX1987-10F and the least was from TGX 1835-10E in the control plots. At 30mM which was the highest level of treatment TGX 1987-10F showed better tolerance level than other varieties. It showed that TGX 1987-10F had a better tolerance level than other varieties. This agrees with the findings of [24,25] who reported that increasing salinity levels reduces the rate of seed germination.

Number of leaves, as the salinity levels increases it affected the number of leaves. There was a drastic reduction in the number of leaves with increasing salinity levels as compared with the control. TGX 1987-10F had the highest number of leaves and it was closely followed by TGX 1835-10E while 1448-2E was the least and there were significantly different from the ones at different salinity levels. This might be that salinity caused deficiency of potassium, which led to reduction in the number of leaves in soybean plants. This finding corroborates with that of [26,27] who reported that salinity induced reduction in the number of leaves in plants.

The result for plant height, revealed a reduction in plant height at every level of salinity as compared with the control. TGX 1448-2E had the highest plant height, followed by TGX 1987-10F while the least was TGX 1835-10E in the control plots. But at 30mM which was the highest level of saline water TGX 1835-10F had the highest plant height and it was closely followed by TGX 1987-10F, this showed that there is presence of salt tolerance genes in the varieties. This result agrees with [28] who reported "a reduction of plant growth under saline conditions which may either be due to osmotic reduction in water availability which resulted in increasing stomatal resistance or to excessive ions, Na and Cl accumulation in the plant tissues" (Machado and Serralheiro, 2017). "It has been reported that salinity stress significantly reduced net photosynthetic rates, increased energy losses for salt exclusion mechanism, largely decreased nutrient uptake and finally reduced plant growth" [29]. "Salinity inhibits the growth of plants by affecting both water absorption and biochemical

**Table 1. Effects of different levels of salinity on growth parameters of three varieties of soybeans**

Varieties	Treatments	Germination percentage (%)	Plant height (cm)	Number of leaves	Number of branches
TGX 1835-10E	Control (0)	85c	30.5c	11.1b	7.1b
	10mM (Nacl)	70e	25.3d	8.0e	4.0d
	20mM (Nacl)	60g	15.1g	5.1h	2.1f
	30mM (Nacl)	50i	10.0h	2.0j	1.0g
TGX 1987-10F	Control (0)	90b	34.3b	12.2a	8.4a
	10mM (Nacl)	80d	20.3e	9.0d	5.1c
	20mM (Nacl)	65f	13.1g	6.4g	3.1e
	30mM (Nacl)	55h	7.3i	3.2i	1.5g
TGX 1448-2E	Control (0)	92a	38.1a	4.8g	9.5a
	10mM (Nacl)	80d	24.1d	10.3c	6.2c
	20mM (Nacl)	70e	16.2f	7.0f	4.1d
	30mM (Nacl)	50i	5.3j	3.3i	2.0f

mM=Millilitres, Nacl= Sodium chloride. Means with the same superscript letter (s) in the same column are not significantly different ( $p \leq 0.05$ ) following separation by Duncan Multiple Range Test

processes, such as nitrogen and carbon dioxide assimilation and protein biosynthesis” [30]. “Under saline conditions, plants fail to maintain the required balance of organic and inorganic constituents leading to suppress growth and yield” [31]. Mshelmbula et al. [24] also reported a reduction of plant height in Cowpea with an increasing level of salinity.

Salinity stress led to a significant reduction in number of branches per plant over control and TGX 1448-2E had the highest number of branches per plant and it was closely followed by TGX 1987-10F while TGX 1835-10E had the least. At 10mM, 20mM and 30mM TGX 1448-2E showed a better level of salt tolerance than both TGX 1835-10E and TGX 1987-10F. Similar result was reported by Azene et al. [32], who reported decrement in number of branches of lentil under different salinity levels. Salinity inhibits the formation of new branch and facilitate the aging of old branch at various degrees [33].

#### 4. CONCLUSION

Growth parameters of soybeans irrespective of variety were influenced by different levels of salinity. As soil salinity increased all the traits exhibited decreasing trends. The findings of this showed that salinity stress significantly affected all the studied parameters as compared to the control. It can be summarized that TGX 1987-10F showed relatively salt tolerant ability as compared with other varieties, therefore it is recommended for cultivation in saline prone area of Cross River State, Nigeria.

#### ACKNOWLEDGEMENT

We sincerely acknowledge Tertiary Education Trust Fund (TETFUND) for funding this research work.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

- Ghulam M, Mohd SA, Rabia A. Global concern for salinity on various agro-ecosystems. In book: Salt Stress, Microbes, and Plant Interactions: Causes and Solution. 2019;1-9.
- Pooja S, Rajesh K. Soil salinity: A serious environmental issue and plant growth promoting bacteria as one of the tools for its alleviation. Saudi Journal of Biological Sciences. 2015;22:123-131.
- Kekeletso HC, Morena MT, Lizelle AP, Ian AD, Fidele T. Soil salinity, a serious environmental issue and plant responses: A metabolomics perspective. Metabolites. 2021;11(11):724.
- Rui MAM, Ricardo PS. Soil Salinity: Effect on vegetable crop growth. management practices to prevent and mitigate soil salinization. Horticulture. 2017;3(2):30.
- Kumari A, Parida AK. Metabolomics and network analysis reveal the potential metabolites and biological pathways involved in salinity tolerance of the

- halophyte *Salvadora persica*. Environ. Exp. Bot. 2018;148:85-99.
6. Singh M, Singh VP, Prasad SM. Nitrogen alleviates salinity toxicity in *Solanum lycopersicum* seedlings by regulating ROS homeostasis. Plant Physiol. Biochem. 2019;141:466-476.
  7. Ahmad P, Ahanger MA, Alyemeni MN, Wijaya L, Alam P, Ashraf M. Mitigation of sodium chloride toxicity *Solanum lycopersicum* L. by supplementation of jasmonic acid and nitric oxide. J Plant. Interact. 2018;13:64-72.
  8. Muhammad K, Aasma P, Sunny A, Zaffar M, Sajid H, Muhammad SC, Muhammad HS, Muhammad A, Parviz H, Jen-Tsung C. Hazardous impacts of soil salinity in crops, tolerance mechanisms, and amelioration through selenium supplementation. Int. J Mol Sci. 2020;21(1):148
  9. Abdalla H, Adarosy MH, Hegazy HS, Abdelhameed RE. Potential of green synthesized titanium dioxide nanoparticles for enhancing seedling emergence, vigor and tolerance indices and DPPH free radical scavenging in two varieties of soybean under salinity stress. BMC Plant Biology. 2022;22(1):1-8.
  10. Dennis LC. Climate change impacts on soil salinity in agricultural areas. European Journal of Soil Science. 2020;(72):842-862.
  11. Bashir N, Khaled RBM, Hamdi AZ, Mukhtar ME. Major limiting factors affecting agriculture use and production. 2021;1:65-75.
  12. Chartes CJ, Noble A. Sustainable intensification: Overcoming land and water constraints on food production. Food security. 2015;7(2):235-245.
  13. Ileya G, Halil B, Halis A. Oil content and composition of soybean genotypes grown in different growing seasons under mediterranean conditions. 2018;39 (2):211-215.
  14. Vaishnav A, Shukla AK, Sharma A, Kumar R, Choudhary DK. "PGPR-Mediated amelioration of crops under salt stress, in plant microbe interaction: An Approach to Sustainable Agriculture Eds. Choudhary, D., Varma, A., Tuteja, N. 2016;10:205-226.
  15. Zewdu S, Suryabhadgavan KV, Batakrisnan M. Geo-spatial approach for soil salinity mapping in Sege Irrigation Farm, South Ethiopia. J. Saudi Soc Agric. Sci. 2017;16:16-24.
  16. Otie V, Udo I, Shao Y, Itam M, Okamoto H, Ping A, Eneji E. Salinity effects on morpho-physiological and yield traits of soybean (*Glycine max* L) as mediated by foliar spray with brassinolide. Plants. 2021;10(3):541.
  17. Farooq M, Hussain M, Wakeel A, Siddique KHM. Salt stress in maize: Effects, resistance mechanisms and management. A review. Agronomy of Sustainable Development. 2015;35:461-481.
  18. Nigeria meteorological services (NIMET) Ikom; 1996.
  19. Saketu H, Gezahagn K. Measurement of soil particle size distribution using hydrometer analysis. American-Eurasian Journal of Agriculture and Environmental Sciences. 2020;20(4):243-254.
  20. Berger VY, Mityaev MV, Sukhotin AA. Practical wet oxidation experiment to determine concentrations of particulate organic matter in seawater. Oceanology. 2016;56:307-310.
  21. Mclean EO. Exchangeable aluminium in C. A. Black (ed.). Methods of Soil Analysis. Plant II Agronomy. 1965;9:985-990.
  22. Bray RH, Kurtz LT. Determination of total organic and forms of phosphorus in soils. 1945;59:39-45.
  23. Gomez KA, Gomez AA. Statistical procedures for agricultural research. John Wiley and Sons, New York; 1984.
  24. Mshelmbula BP, Zakariya R, Mensah JK, Ikhajiagbe B. Effect of Salinity on Germination, Growth and Yield Performance of Cowpea (*Vigna unguiculata* L. Walp.) in Mubi, Nigeria. Nigerian Annals of Natural Sciences. 2015;15 (1):018–023.
  25. Kumar A. Germination behaviour of soybean varieties under different salinity stress. International Journal of Applied Agricultural Research. 2017;12 (1):69-76.
  26. Smurutishree S, Bhagawati B, Nitesh. Salinity tolerance in wheat. Marumegh. 2016;3(1).
  27. Amirhossein H, Adisa A, Nima S. Global predictions of primary soil salinization under changing climate in the 21<sup>st</sup> century. 2021;12:6663.
  28. Davood S, Nasser D, Behnam AL, Manour G. An overview on improvement of crop productivity in saline soils by halotolerant and halophilic PGPRs. Biotech. 2019;9(7):261.
  29. Machado RMA, Ricardo PS. Soil salinity: Effect on vegetable crop growth.

- management practices to prevent and mitigate soil salinization. *Horticulturae*. 2017;3(2):13.
30. Mohammad AA, Nisha ST, Megha T, Surendra A, Agarwal RM. *Physiol Mol Biol Plants*. 2017;23(4):731-744.
31. Maximillian J, Brusseau ML, Glenn EP, Mathias AD. Pollution and environmental perturbations in the global system. *Environmental and Pollution Science*. 2019;3:457-476.
32. Azene T, Yohannes P, Habtamu Z. Effect of salinity on yield and yield related traits of some accessions of ethiopian lentil (*Lens culinaris* M.) under Greenhouse Conditions. *International Journal of Technology Enhancements and Emerging Engineering Research*. 2014;2(10): 2347-4289.
33. Duncan BD. Multiple range and multiple F. Test. *Biometrics*. 1955;2: 1-42.

---

© 2023 Aboh and Ikwa; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/106734>