



# Effect of Organic Nutrients and Bio Fertilizers on Soil Parameters and Nutritional Content of Amaranth

Swati Sulagna Sarangi <sup>a++</sup>, Ira Sarma <sup>b#\*</sup>, Sailen Gogoi <sup>bt</sup>  
and Aradhana Barooah <sup>ct</sup>

<sup>a</sup> Odisha University of Agriculture and Technology, Bhubaneswar, Odisha, India.

<sup>b</sup> Department of Horticulture, Assam Agricultural University, Jorhat, 785013, Assam, India.

<sup>c</sup> College of Horticulture and Farming System Research, Nalbari, Assam Agricultural University, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJECC/2023/v13i113613

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/108604>

Original Research Article

Received: 24/09/2023

Accepted: 30/11/2023

Published: 05/12/2023

## ABSTRACT

A field experiment was conducted to study the effect of organic nutrients and bio fertilizers on soil parameters and nutritional content of amaranth. Eleven treatments were taken in three replications and the experiment was conducted in a randomized block design (RBD). The treatments were T<sub>1</sub>:RDF (60:30:20 kg NPK/ha), T<sub>2</sub>:Vermicompost @ 2 t/ha, T<sub>3</sub>:Vermicompost @ 5 t/ha, T<sub>4</sub>:FYM 10 t/ha, T<sub>5</sub>:FYM 20t/ha, T<sub>6</sub>:Poultry manure 3t/ha, T<sub>7</sub>:Poultry manure 5 t/ha, T<sub>8</sub>:Vermicompost @ 2 t/ha+PSB+Azospirillum@ 5 kg/ha each, T<sub>9</sub>:FYM @ 10t/ha+ PSB+Azospirillum@ 5 kg/ha each, T<sub>10</sub>:Poultry manure 3 t/ha+ PSB+Azospirillum@ 5 kg/ha each and T<sub>11</sub>: Untreated control. The highest vitamin C (147.8 mg/100 gm) and crude fibre content (1.48%) was observed in the

<sup>++</sup> Ph.D Scholar;

<sup>#</sup> Assistant Professor;

<sup>†</sup> Principal Scientist;

<sup>‡</sup> Associate Professor

\*Corresponding author: E-mail: ira.sarma@aau.ac.in;

treatment T<sub>8</sub> and T<sub>4</sub>, respectively. Maximum calcium (8.36%) and iron (0.21%) content was observed in the treatment T<sub>8</sub>. The treatment T<sub>10</sub> recorded the highest mean performance for P<sup>H</sup>, organic carbon, available nitrogen and potash content in the soil analyzed after harvest of the crop.

**Keywords:** Vermicompost; FYM; poultry manure and amaranth.

## 1. INTRODUCTION

Amaranth is an important leafy vegetable and a good source of protein, minerals, vitamin A and vitamin C. Depending on the colour of the leaves, green and red coloured varieties are available. One of the most primary factors for increasing the production as well as the quality of amaranth leaves is the health of the soil. Soil should have ideal physical, chemical and biological conditions. Organic fertilizers positively influence the microbial population of the soil, which increases the soil's biomass, carbon, nitrogen content and dehydrogenase enzyme activity. Combining organic manure with bio-fertilizers increases crop yield and also helps to improve the physical health of the soil and serves as a reservoir for nutrients [1]. Organic fertilizers supply a variety of carbon compounds with diverse chemical compositions, ranging from readily degradable to stable, which can be utilized by soil microorganisms during mineralization to increase their growth rates and biomass [2]. Therefore, organic fertilizers have strong sustainable effects on the soil microorganisms and are the foundation for improving soil health by rising microbial activity, microbial interactions and nutrient cycling [3,4]. The long-term application of chemical fertilizers increase soil acidity and organic fertilizers can increase the value of soil pH and soil organic carbon and reduce soil acidification [5,6]. Due to rising cost of inorganic fertilizers the use of available local resources such as farmyard manure, vermicompost, poultry manure, etc. are very much essential to build up the scarcity. The use of organic manures not only increase crop production but also considerably improves the physical, chemical and biological qualities of soil as well as boosts the effectiveness of fertilizers. Vermicompost improves the physical properties by reducing the bulk density of soil and it also increases the water holding capacity of soil [7]. Well-decomposed FYM and vermicompost alone or in conjunction with bio-fertilizers aid in supplying the right amount of nutrients and preserve the health of the soil. It supplies necessary plant nutrients, which enhance the physical and chemical characteristics, increase water holding capacity and promote microbial

activity in the soil. Poultry manures are very nutrient-rich organic manure and N, P, K content are 0.5-0.9, 0.4-0.5 and 1.2-1.7 percent, respectively [8]. Biofertilizers increase soil productivity by fixing atmospheric nitrogen or by solubilizing the soil phosphate and boost plant growth by providing growth-promoting substances. Considering the importance of organic nutrients, the investigation was carried out to study the impact of vermicompost, FYM, poultry manure and bio fertilizers on soil parameters and nutrient content of amaranth.

## 2. MATERIALS AND METHODS

The experiment was conducted in a randomized block design (RBD) comprising of eleven treatments in three replications. The treatments used in the experiment were T<sub>1</sub>: RDF (60:30:20 Kg NPK/ha), T<sub>2</sub>:Vermicompost @ 2 t/ha, T<sub>3</sub>:Vermicompost @ 5 t/ha, T<sub>4</sub>:FYM 10 t/ha, T<sub>5</sub>:FYM 20 t/ha, T<sub>6</sub>:Poultry manure 3 t/ha, T<sub>7</sub>:Poultry manure 5 t/ha, T<sub>8</sub>:Vermicompost @ 2 t/ha+ PSB+ *Azospirillum*@ 5 kg/ha each, T<sub>9</sub>:FYM @ 10 t/ha+ PSB+*Azospirillum*@ 5 kg/ha each, T<sub>10</sub>:Poultry manure 3 t/ha+ PSB+*Azospirillum*@ 5 kg/ha each and T<sub>11</sub>: Untreated control. The experimental area was ploughed, leveled and harrowed to a fine tilth. The plot was divided into equal plots, each having a size of 4sqm and each plot was separated by 50 cm. The recommended dose of inorganic fertilizers for amaranth is 60:30:20 kg NPK/ha.

After proper land preparation, biofertilizers were applied to the individual plots according to the treatments three days before sowing of seeds. According to the treatment plan, vermicompost, FYM and poultry manure were also applied to the plots. The organic manures and biofertilizers were carefully mixed into the soil. Seeds were sown in well-prepared plot. At the time of sowing, seeds were mixed with fine sand for even distribution of the seeds as the seed size is very small. The seeds were sown at 1cm depth and covered evenly with thin layer of fine soil. Extra seedlings were thinned off after the seedlings were well-established to maintain a proper spacing of 10cm between plants. The quality parameters viz., vitamin C, crude fibre, calcium,

iron and soil parameters were analyzed after harvesting of crop.

### 3. RESULTS AND DISCUSSION

The treatment T<sub>8</sub> exhibited significantly the highest (147.8mg) mean performance for vitamin C content of leaves whereas, the treatment T<sub>11</sub> (Control) showed the lowest (75.3mg) value for vitamin C (Table1). The ability of *Azospirillum* to fix atmospheric nitrogen and the production of hormones that promote plant development by both vermicompost and *Azospirillum* might be responsible for the rise in vitamin C content. These hormones assisted to speed up the process of carbohydrate synthesis. Because vitamin C is a sugar acid, its synthesis is encouraged by the application of organic fertilizers [9]. Inoculation with phosphate solubilizing biofertilizer helpful in assimilation of carbohydrates and which in turn increased the synthesis of ascorbic acid [10]. The treatment T<sub>4</sub> (FYM 10 t/ha) recorded the highest (1.48%) value for crude fibre content while the lowest (1.10%) was exhibited by the treatment T<sub>11</sub> (Control). The mean value (1.47%) of treatment T<sub>5</sub> (FYM 20 t/ha) was at par with T<sub>4</sub>. This might be due to lesser amount of nitrogen in FYM as compared to vermicompost because decreased in nitrogen content enhance the fiber content of leaves. The percentage of crude fiber content is reduced with enhancement of phosphorus.

The treatment T<sub>8</sub> recorded significantly the highest mean performance for calcium content with a value of 8.36% and it was at par with the treatment T<sub>10</sub> (8.28). The increased calcium content of amaranth leaves might be due to increased absorption of micro and macro nutrients by the crop. Shankar et al. [11] reported

that more microorganisms are present in organically managed soil and these microbes create a variety of chemicals that encourage plants to take up more micronutrients from the soil. Organic acids present in vermicompost had great role in releasing of calcium from exchange sites and thereby increased the availability of calcium. According to Kumar [12], the organic acids in vermicompost have a significant role in releasing calcium from exchange sites, increasing calcium availability.

The treatment T<sub>8</sub> showed significantly the highest mean performance for iron content with a value of 0.21%. The high iron content in the leaves of amaranthus might be due to vermicompost which increased the micronutrients in soil and it also acted as a chelating agent. It plays important role to make available the other metallic micronutrients to the plants. The abundance of microorganisms due to addition of vermicompost increases siderophores production and enhances the availability of Fe to plants. This is essential for iron in particular since many soils have substantial iron content but in inaccessible form. Similar observations were also reported by Shankar et al. [11]. Vermicompost is chemically as well as biologically enriched compound as compared to compost and the pH of the vermicompost exerts a strong influence on the availability of micronutrients like iron [13].

The treatment T<sub>10</sub> (Poultry manure 3 t/ha + PSB +*Azospirillum*@ 5kg/ha each) recorded the highest mean performance for P<sup>H</sup> (5.34), organic carbon (1.24%), available nitrogen (237kg) and potash (141.23kg) in the present investigation (Table 2). Combining bio-fertilizers with organic nutrients had a positive impact on soil health by

**Table 1. Effect of organic nutrients and bio fertilizers on nutritional content of amaranth leaves**

Treatments	Vitamin C (mg/100g)	Crude fibre (%)	Calcium (%)	Iron (%)
T <sub>1</sub>	76.2	1.23	7.46	0.17
T <sub>2</sub>	102.1	1.29	6.25	0.18
T <sub>3</sub>	103.1	1.29	6.71	0.19
T <sub>4</sub>	82.2	1.48	6.22	0.17
T <sub>5</sub>	84.2	1.47	6.25	0.19
T <sub>6</sub>	92.2	1.30	5.32	0.18
T <sub>7</sub>	94.1	1.32	6.34	0.19
T <sub>8</sub>	147.8	1.14	8.36	0.21
T <sub>9</sub>	141.9	1.23	8.28	0.20
T <sub>10</sub>	145.9	1.27	7.39	0.20
T <sub>11</sub>	75.3	1.10	5.27	0.15
SEd (±)	0.4	0.03	0.21	0.01
CD (0.05)	0.9	0.06	0.45	0.02

**Table 2. Effect of organic nutrients and bio fertilizers on soil parameters after harvest**

Treatments	pH	Organic carbon (%)	Available nitrogen (kg/ha)	Available phosphorus (kg/ha)	Available potash (kg/ha)
T <sub>1</sub>	4.91	0.71	228.00	38.98	125.22
T <sub>2</sub>	5.16	0.91	215.33	45.76	134.15
T <sub>3</sub>	5.26	0.95	229.00	47.25	139.51
T <sub>4</sub>	4.96	0.75	206.00	41.12	128.06
T <sub>5</sub>	5.16	0.82	212.67	45.12	132.55
T <sub>6</sub>	5.19	0.77	215.00	46.56	135.15
T <sub>7</sub>	5.23	0.85	226.00	49.41	137.76
T <sub>8</sub>	5.28	1.19	235.00	51.15	141.15
T <sub>9</sub>	5.22	1.11	221.00	49.15	136.77
T <sub>10</sub>	5.34	1.24	237.00	52.23	141.23
T <sub>11</sub>	4.86	0.69	202.00	36.12	118.60
SEd (±)	0.03	0.01	1.34	0.15	0.06
CD (0.05)	0.05	0.03	2.81	0.32	0.13

enhancing the soil structure, texture, ability to retain water, capacity as a buffer, nutritional status and microbial population. The present study showed that organic nutrients alone and in conjunction with bio-fertilizers boost soil fertility and nutritional status. According to experimental findings, the addition of *Azospirillum* and PSB to organic nutrients greatly improved the availability of nitrogen, phosphorus and potassium as compared to the treatments without bio-fertilizers. The increase in nitrogen availability might be attributable to increased percentage of nitrogen in vermicompost and *Azospirillum* population, which may have fixed atmospheric nitrogen non-symbiotically and increased nitrogen solubilization in soil.

The increase in the amount of phosphorus that is readily available in soil following the addition of vermicompost and biofertilizer may be attributable to the more effective and efficient mineralization and solubilization of phosphorus. Phosphorus present in soil is unavailable to plants and it is made available by the action of bacteria that produce phosphatase and other phosphorus-solubilizing acids and enzymes. Similar findings were also reported by Han et al. [14], Narayanamma et al. [15] and Parthasarathi et al. [16].

#### 4. CONCLUSION

Based on the result obtained from the investigations it can be concluded that different treatments exhibited significant differences in nutritional content of amaranth leaves. The highest vitamin C, crude fibre, calcium and iron content were observed in the treatment T<sub>8</sub>. The

highest values for P<sup>H</sup>, organic carbon, available nitrogen and potash content in the soil were exhibited by the treatment T<sub>10</sub>.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Yadav S, Asati KP, Barche S. Effect of integrated nutrient management strategies on growth and yield of Amaranthus (*Amaranthus tricolor* L.). J.Pharm.Innov. 2022;11:2545-2549.
2. Lazcano C, Zhu-Barker X, Decock C. Effects of organic fertilizers on the soil microorganisms responsible for N<sub>2</sub>O emissions: a review. Microorganisms. 2021;9:983. DOI:10.3390/microorganisms9050983.
3. Lazcano C, Gomez-Brandon M, Revilla P, Domínguez J. Short-term effects of organic and inorganic fertilizers on soil microbial community structure and function. Biol. Fertil. Soils. 2013;49:723–733. DOI: 10.1007/s00374-012-0761-7.
4. Ling N, Zhu C, Xue C, Chen H, Duan Y, Peng C. Insight into how organic amendments can shape the soil microbiome in long-term field experiments as revealed by network analysis. Soil Biol. Biochem. 2016;99:137–149. DOI: 10.1016/j.soilbio.2016.05.005.
5. Wang H, Xu J, Liu X, Zhang D, Li L, Li W. Effects of long-term application of organic fertilizer on improving organic matter

- content and retarding acidity in red soil from China. *Soil Tillage Res.* 2019; 195: 104382.  
DOI: 10.1016/j.still.2019.104382.
6. Liu J, Shu A, Song W, Shi W, Li M, Zhang W. Long-term organic fertilizer substitution increases rice yield by improving soil properties and regulating soil bacteria. *Geoderma.* 2021; 404:115287.  
DOI: 10.1016/j.geoderma.2021.115287.
  7. Ravimycin T. Effects of vermicompost (VC) and farmyard manure (FYM) on the germination percentage growth biochemical and nutrient content of Coriander (*Coriandrum sativum* L.). *Int. J. Adv. Res. Biol. Sci.* 2016;3: 91-98.
  8. Adekiya AO, Aboyeji CM, Dunsin O, Asamu F, Owolabi AO, Aremu CO, Owolabi IO. Poultry manure addition affects production, plant nutritional status and heavy metals accumulation in green Amaranth (*Amaranthus hybridus*). *Int. J. Agric. Biol.* 2019;22:993-1000.
  9. Vlahova V and Popov V. Influence of the bio-fertilizer season on yield of pepper (*Capsicum annuum* L.) cultivated under organic agriculture conditions. *J. Org. Syst.* 2013;8(2):6-17.
  10. Upadhyay AK, Bahadur A, Singh J. Effect of organic manures and biofertilizers on yield, dry matter partitioning and quality traits of cabbage (*Brassica oleracea* var. *capitata*). *Indian J. Agric. Sci.* 2012; 82:31–34.
  11. Shankar KS, Sumathi S, Shankar M, Rani KU. Effect of organic farming on nutritional profile, quality characteristics and toxic parameters of amaranthus. *Indian J. Hortic.* 2013;3:378-382.
  12. Kumar PT, Binoo P, Bonny R, John MJ, Divya MR and Roch CV. Effect of organic and inorganic nutrient sources on the yield of selected tropical vegetables. *Sci. Hortic.* 2017;224:84-92.
  13. Wright AL, Provin TL, Hons FM, Zuberer DA, White RH. Soil micronutrient availability aftercompost addition to St. Augustine Grass. *Compost Science & Utilization.* 2007;15(2):127-134
  14. Han HS, Supanjani KD and Lee. Effect of co-inoculation with Phosphate and Potassium Solubilizing bacteria on mineral uptake and growth of pepper and cucumber. *Plant Soil Environ.* 2006;52: 130-136.
  15. Narayanamma M, Chiranjeevi CH, Reddy IP, Ahmed SR. Integrated nutrient management in cauliflower (*Brassica oleracea* var. *botrytis* L.). *Veg. Sci.* 2005; 32(1):62-64.
  16. Parthasarathi K, Balamurugan M and Ranganathan LS. Influence of vermicompost on the physico-chemical and biological properties in different types of soils along with yield and quality of pulse crop blackgram. *Iran. J. Environ. Health. Sci. Eng.* 2008;5(1):51-58.

© 2023 Sarangi et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/108604>