



Productive Performance and Fruit Quality of Short Days Italian Genetic-based Strawberry Cultivars and Genotypes in Southern Brazil

Paola Daiane Welter ^a, Juliana Martins de Lima ^a,
Antônio Felipe Fagherazzi ^a, Daiana Petry Rufato ^b,
Leo Rufato ^a, Aike Anneliese Kretzschmar ^a,
Geanlucca Baruzzi ^c and Amauri Bogo ^{a*}

^a Crop Production Graduate Program, Santa Catarina State University/UFSC. Avenida Luis de Camões, 2090, CEP 88520-000, Lages-SC, Brazil.

^b Department of Environmental Engineering, University of Santa Catarina State. Avenida Luis de Camões, 2090, CEP 88520-000, Lages-SC, Brazil.

^c Istituto Sperimentale per la Frutticoltura, Sezione di Forlì, Via Punta di Ferro 2, 47100 Forlì, Italy.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JEAI/2023/v45i122277

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/111023>

Original Research Article

Received: 17/10/2023

Accepted: 21/12/2023

Published: 30/12/2023

ABSTRACT

When grown under different climatic conditions, short days Italian genetic-based strawberry (IGBS) cultivars and genotypes present variations in productive performance and fruit quality. The objective of this study was to assess different productive performance and fruit quality of IGBS

*Corresponding author: E-mail: amauri.bogo@udesc.br;

cultivars and genotypes grown in highland regions of southern Brazil. The experiment were conducted in randomized block design using three cultivars and 21 genotypes during the 2019, 2020 and 2021 growth seasons. The data were subjected to multivariate analysis using principal component analysis (PCA). IGBS cultivars and genotypes differed significantly in terms of productive and fruit quality parameters, according to PCA data. The first and second main components of PCA, respectively, were responsible for 61.9% and 72.4% of the productive performance and fruit quality parameters. The Pircinque cultivar and FRF PIR 256.04, ITA 13.097.05, ITA 12.103.22, ITA 10.107.07, and ITA 10.133.02 genotypes showed the highest indices of productive parameters of total production per plant, commercial production per plant, total fruit number per plant and commercial fruits number per plant. Among these, the Pircinque cultivar and the FRF PIR 256.04 genotypes demonstrated the best fruit quality parameters of fruit color (skin luminosity, chroma, and hue angle), flavor (soluble solids/ titratable acidity ratio), and fresh commercial fruit mass in highland regions of southern Brazil during the 2018, 2019 and 2020 growth seasons.

Keywords: *Fragaria x ananassa* Duch.; agronomic performance; adaptability; yield; productivity.

1. INTRODUCTION

In the last 20 years, global strawberry (*Fragaria x ananassa* Duch.) production has more than doubled to over 3.6 million tons. The Northern Hemisphere accounts for (98%) of production, with genetic and climatic barriers preventing expansion to the south hemisphere [1]. Strawberry crop development in Brazil began in the mid-1960s, with the launch of the Campinas cultivar [2], and production of the crop has increased in the country since then. Currently, production is around 165,000 tons in about 4,500 hectares of cultivated land, with an average productivity of 36.6 tons per hectare [1].

Strawberry cultivation progress is linked to the introduction of new cultivars to the market, improvement of cultivation techniques and systems, as well as climatic adaptation. Strawberry cultivation has been concentrated primarily on family properties in southern Brazil, particularly in temperate and subtropical regions [3].

Currently, national strawberry production is almost entirely dependent on imported cultivars from the University of California, United States of America (USA) strawberry breeding programs. Many of these cultivars, however, are grown without any prior research into climatic adaptation and productive requirements for different producing regions in southern Brazil. Brazil has a diverse climate and can produce strawberries almost every month of the year. However, due to the country's edaphoclimatic diversity, the small number of available cultivars has been one of the main barriers to the development of the strawberry crop. National breeding programs are essential for encouraging

the development and introduction of new cultivars developed in other countries [4,5].

A cultivar's climatic adaptation to a specific producing region is expressed through the interaction among genotype, environment, and genetic feature, which influences production, productivity, and fruit quality. Thus, the responses of cultivars to fruit yield and quality are primarily determined by the interaction of two factors, photoperiod and temperature [4].

Adaptability studies of cultivars and genotypes are extremely important and necessary, not only in new producing regions, but also in traditional ones, which often, for cultural reasons, do not seek new cultivars that enable better agronomic performance, which is essential for producers' decision-making regarding the best material to be grown in different regions [6].

As a result, the adaptability study with short days Italian genetic-based strawberry cultivars and genotypes in southern Brazil regions is important for obtaining new materials, increasing the cultivars offer in the producing market, and materials that are more productive and have better fruit quality. Thus, the objective of this study was to assess the productive performance and fruit quality of short days Italian genetic-based strawberry cultivars and advanced genotypes grown under edaphoclimatic conditions in highland regions of southern Brazil during the 2019, 2020, and 2021 growth seasons.

2. MATERIALS AND METHODS

The experiments were conducted in Lages Municipality (27°47'05" S, 50°18'08" W), at an altitude of 922 m above sea level, in Santa

Catarina State, Southern Brazil, from May to December of the 2019, 2020, and 2021 growing seasons. According to Köppen classification, the climate of the region is subtropical humid mesothermal Cfb, and soil type is Clay Auric Cambisol [7]. The average annual temperature and rainfall are around 15.6 °C and 1,500 mm, respectively. Daily rainfall, relative humidity, and hourly temperatures were recorded from May to December of the 2019, 2020, and 2021 growth seasons by the Santa Catarina Hydrology and Environmental Resources Center (Epagri). The weather station (Onset Computer Corp., ProCasset, MA, USA; Davis Pro2-6153) was 500 m away from the vineyard.

The experiments included 24 treatments involving three Spanish and Italian cultivars and 21 Italian short days genetic-based genotypes from Consiglio per la Ricerca in Agricoltura-Centro di Olivicoltura, Frutticoltura e Agrumicoltura – CREA-OFA/Italy and Santa Catarina State University (UDESC). Strawberry seedlings used in all experiments were grown field-collected stolon's and rooted in 72 cells trays in a commercial nursery in the Farroupilha Municipality of Rio Grande do Sul State, southern Brazil. The cultivars Jonica, Sabrina, and Pircinque, and the genotypes FRF LAM 269.18, FRF PIR 075.08, FRF 256.04, FRF LAM 263.01, FRF PA 109.02, FRF PIR 079.06, ITA 10.133.02, ITA 13.079.01, ITA 13.079.02, ITA 13.097.05, ITA 12.190.02, ITA 10.107.12, ITA 10.107.07, ITA 10.107.06, ITA 12.103.04, ITA 12.103.06, ITA 12.103.12, ITA 12.103.15, ITA 10.103.22, ITA 10.128.06, and ITA 10.128.09 were evaluated.

A conventional soil cultivation system in low tunnels was used in an experimental randomized blocks design with four replications and eleven plants in each experimental unit. Strawberry seedlings were planted by hand beginning in May of each year of 2019, 2020, and 2021. Soil correction and fertilization were carried out in accordance with soil analysis [8]. Fertilization was done weekly with fertigation, which included: calcium nitrate (9.92 g/100 plants⁻¹); monoamonic phosphate (38.74 g/100 plants⁻¹), P51 (liquid fertilizer containing 51% phosphorus) (7.3 mL/100 plants⁻¹); magnesium sulfate (4.55 g/100 plants⁻¹), and potassium phosphate (31.25 g/100 plants⁻¹).

The experimental beds were 1.00 × 27 × 0.20 m in size for each cultivar and genotype (L × C × H). The beds were covered with 30-micron thick

black polyethylene plastic ('mulching'). Strawberry plants were grown in three rows in a "V" or quinconic system, with 30 cm between plants and 60,000 plants per ha. The irrigation system was drip, with 0.15 m spacing between drippers and a flow rate of 1.5 liters per hour, with three hoses per bed.

When necessary, pest and disease control was carried out using crop-specific products. Acaricide abamectin, insecticide thiametoxan, and fungicides methyl thiophanate (only before harvest), fluazinam, iprodione, pyrimethanil, and azoxystrobin + diphenconazole were also used. Weeds were controlled by weeding between the beds and manual tillage in the planting pits, and old and diseased leaves were removed with pruning shears.

Beginning at the end of August and lasting until the middle of February the following year, harvest were carried out every three days when the fruits displayed epidermis with approximately 70%–80% of the red color. Strawberry fruits were counted, weighed on a semi-analytical scale, and classified using the following criteria:

- 1) Commercial fruits are those that weight more than or equal to 10 grams and are free of injuries, deformations, or rots;
- 2) Small fruits weighing less than 10 grams and free of injuries, deformations, or rots;
- 3) Discarded fruits are those with injuries, deformations, or rots.

2.1 Productive Parameters

The following productive parameters were quantified using the number and weight of analyzed fruits.

a) total plant production (TPP) (g/plant⁻¹); b) commercial production per plant (CPP) (g/plant⁻¹); c) total productivity (TPT) (t/ha⁻¹); d) commercial productivity (CPT) (t/ha⁻¹); e) number of total fruits per plant (NTF) (fruits/plant⁻¹); f) number of commercial fruits per plant (NCF) (fruit/plant⁻¹); g) percentage of commercial production (PCP) (%); h) percentage of small fruit per plant (PSF) (%); and, i) percentage of discard fruits per plant (PDF) (%).

2.2 Fruit Quality Parameters

The following fruit quality parameters were quantified using the number and weight of analyzed fruits. a) fruit pulp firmness (FPF) (g/fruit⁻¹); b) epidermis luminosity (L*); c)

epidermis chroma (C*); d) hue angle (°hue); e) soluble solids (SS) (°Brix); (f) titratable acidity (TA) (g. 100g citric acid); g) soluble solids/titratable acidity ratio (SS/TA); h) average fresh mass of commercial fruits (AMCF) (g/fruit¹).

The physicochemical analysis was carried out using uniform samples of ten fruits per repetition, at each new plant flowering, for a total of four analyses per plant productive cycle. The physicochemical analyses were carried out in accordance with the following methodologies:

- a) Texture Analyzer TA texturometer (Stable Micro Systems Ltd., Vienna Court, UK) was used to measure the L*, C*, °hue (color angle), and FPF parameters expressed in grams;
- b) SS are measured using a digital refractometer and expressed as the concentration of sugars in fruits (°Brix); TA is measured using an automatic TITRONIC titrator® with 5 mL of juice in 45 mL of distilled water and titration in sodium hydroxide solution 0.1N at pH 8.1; and the SS/TA ratio was calculated as the ratio of SS content and TA.

2.3 Data Analysis

The analyses were carried out with data from harvest in 2019, 2020, and 2021, and were carried out by the Shapiro-Wilk normality test, using the Software R [9], via the Action Stat interface. The univariate analysis was carried out using analysis of variance and a scott knott test with a 5% probability of error using the statistical program SISVAR [10]. Fitopac 2.1's statistical program used the PCA for the multivariate analysis [11].

3. RESULTS

The Pircinque cultivar and FRF PIR 256.04, ITA 10.133.02, ITA 13,097.05, ITA 10.107.07, and ITA 12,103.22 genotypes were significantly more productive and had higher fruit quality. The Pircinque cultivar and the FRF PIR 256.04, ITA 10.133.02, ITA 13,097.05, ITA 10.107.07, and ITA 12,103.22 genotypes had a significant influence on the productive parameters of CPP, NTF, NCF and, AMCF, as well as the fruits quality parameters of L, C*, °hue, SS (°Brix), and SS/TA ratio. The Pircinque cultivar and the FRF PIR 256.04, ITA 10.133.02, ITA 13.097.05, ITA 10.107.07, and ITA 12,103.22 genotypes did not

show significant differences in fruit quality parameters, which were related to the sweet taste of the fruits and considered very promising cultivar and genotypes for the Brazilian market (Table 1).

The Pircinque and Sabrina cultivars and the FRF PIR 256.04 and ITA 12.103.22 genotypes, produced significant fruits with high FPF and SS and were found to be positively related to the fruits quality parameters L, C, °hue, and SS/TA, indicating fruits with higher sugar content (Table 1). The Pircinque and Sabrina cultivars and the FRF PIR 256.04, ITA 10.107.07, and ITA 12,103.22 genotypes, produced significantly higher-quality fruits with SS/TA ratios that were 25% higher on average than the other cultivars and genotypes, indicating high tasty fruits (Table 1).

Figs. 1 and 2 depicted the PCA data of the cultivars and genotypes productive and fruit quality parameters, respectively. The PCA data of all harvests occurring during the 2019, 2020, and 2021 growth seasons revealed, via vectors, the formation of two major groups of cultivars and genotypes with a positive correlation with productive and fruit quality parameters.

In Fig. 1, both main components were responsible for 85.4% (61.9% plus 23.5%) of the cultivars and genotypes differentiation, according to productive parameters. According to TPP, CP, NTF, and NCF productive parameters, the first main component was responsible for 61.9% of the differentiation of cultivars and genotypes. Thus, the Pircinque cultivar and the FRF PIR 256.04, ITA 13.097.05, ITA 12.103.22, ITA 10.107.07, and ITA 10.133.02 genotypes, highlighted in Fig. 1, were differentiated by showing the highest productive parameters and consequently, more productive plants. The second main component was responsible for 23.5% in the cultivars and genotypes differentiation, according to PSF and PDF productive parameters. Thus, the Pircinque cultivar and the FRF PIR 256.04, ITA 13.097.05, ITA 12.103.22, ITA 10.107.07, and ITA 10.133.02 genotypes, highlighted in Fig. 1, were differentiated by the fruit classification, considered non-commercial fruits because they are either small or discard fruits. The Pircinque cultivar and the FRF PIR 256.04 genotype produced more discarded fruits than small fruits per plant, whereas the ITA 10.107.07 and ITA 10.133.02 genotypes produced more small fruits than discarded fruits per plant.

Table 1. Analysis of variance (ANOVA) of productive parameters of a) total production per plant (TPP) (g plant⁻¹); b) commercial production per plant (CPP) (g plant⁻¹); c) total productivity (TPT) (t ha⁻¹); d) commercial productivity (CPT) (t ha⁻¹); e) number of total fruits per plant (NTF) (fruits/plant⁻¹); f) number of commercial fruits per plant (NCF) (fruit/ plant⁻¹); g) percentage of commercial production (PCP) (%); h) percentage of small fruit production (PSF) (%); i) percentage of discard fruits (PDF) (%), and fruit quality parameters of a) fruit pulp firmness (FPF) (g fruit⁻¹); b) epidermis luminosity (L*); c) epidermis chroma (C*); d) hue angle (°hue); e) soluble solids (SS) (°Brix); (f) titratable acidity (TA) (g/100g citric acid) and, g) soluble solids/titratable acidity ratio (SS/TA); h) average fresh mass of commercial fruits (AMCF) (g fruit⁻¹), of experiments conducted with short days Italian genetic-based strawberry cultivars and genotypes, during the 2019, 2020 and 2021 growth seasons, at the Lages/SC Municipality, southern Brazil.

Cultivars/genotypes	TPP	CPP	TPT	CPT	NTF	NCF	AMCF	FPF	PSF(%)	PDF(%)	PD(%)	L*	C*	°hue	SS	TA	SS/TA
Jonica	392,58* a	208,32 b	23,55 a	12,50 b	28,05 b	13,30 b	15,65 b	222,56 b	52,57 e	11,87 c	35,57 d	35,24 b	41,77 d	32,47 d	6,6 b	0,48 c	13,80 b
Pircinque	398,99 a	263,43 a	23,94 a	15,80 a	31,45 a	14,95 a	19,00 a	293,63 a	75,32 a	6,57 a	18,07 b	38,11 a	47,70 a	22,13 a	7,8 a	0,41 a	16,87 a
Sabrina	296,04 c	122,17 c	11,76 d	7,33 d	13,15 d	6,71 d	18,57 a	321,72 a	62,55 c	6,62 a	30,80 d	35,14 b	42,08 c	27,70 a	7,9 a	0,47 c	16,46 a
FRF LAM 269.18	337,50 b	219,94 b	20,25 b	13,19 b	22,53 c	11,45 b	19,32 a	193,22 c	65,15 c	5,35 a	29,55 d	33,86 c	41,16 d	29,95 b	7,2 b	0,54 d	13,27 b
FRF PIR 075.08	331,33 b	218,49 b	19,87 b	13,11 b	25,31 b	13,18 b	16,52 b	241,08 b	65,70 c	8,07 b	26,22 c	34,83 b	43,63 c	31,06 c	7,8 a	0,58 e	13,65 b
FRF PIR 256.04	412,66 a	279,43 a	24,76 a	16,79 a	32,88 a	15,11 a	18,95 a	282,30 a	67,95 c	5,17 a	26,90 c	37,16 a	46,18 a	21,73 a	7,9 a	0,38 a	16,07 a
FRF LAM 263.01	264,90 c	170,12 c	15,89 c	10,20 c	23,63 c	11,39 b	15,27 b	220,29 c	63,82 c	12,70 c	23,47 c	34,57 c	39,69 e	32,70 d	5,8 c	0,51 c	11,47 c
FRF PA 109.02	366,25 b	216,35 b	21,97 b	12,97 b	28,14 b	12,02 b	18,12 a	218,82 c	58,97 d	9,32 b	31,72 d	33,56 c	39,71 e	31,33 c	6,5 b	0,44 b	13,84 b
FRF PIR 079.06	276,19 c	164,54 c	16,57 c	9,87 c	20,37 c	9,32 c	17,42 a	197,06 c	58,77 d	8,15 b	33,07 d	38,60 a	46,87 a	34,87 e	8,2 a	0,67 e	13,38 b
ITA 10.133.02	377,73 a	273,61 a	19,06 b	13,41 b	28,67 a	14,87 a	17,41 a	228,79 c	70,35 b	13,60 c	16,05 b	34,34 c	40,86 d	30,80 c	6,8 b	0,70 e	9,54 c
ITA 13.079.01	179,59 d	81,95 d	10,77 d	4,91 e	16,67 d	5,04 e	16,21 b	247,90 b	45,67 f	14,95 c	39,35 d	35,13 b	43,09 c	33,92 e	6,8 b	0,67 e	10,06 c
ITA 13.079.02	228,35 c	143,49 c	13,69 c	8,61 c	21,63 c	9,25 c	15,56 b	263,93 b	63,00 c	25,40 d	11,60 a	35,58 b	41,00 d	33,74 e	5,4 c	0,57 d	9,39 c
ITA 13.097.05	388,22 a	262,10 a	22,09 b	13,92 b	29,86 a	15,07 a	17,87 a	164,41 d	63,10 c	7,72 b	29,20 d	34,03 c	42,23 c	32,15 d	4,9 d	0,45 b	10,74 c
ITA 12.190.02	75,49 e	49,05 e	4,53 e	2,94 f	6,16 e	3,14 f	15,84 b	98,40 e	64,90 c	9,97 b	25,15 c	39,30 a	44,39 b	34,90 e	5,8 c	0,54 d	10,81 c
ITA 10.107.12	250,47 c	147,60 c	15,02 c	8,85 c	20,84 c	7,83 d	18,66 a	249,62 b	58,42 d	19,10 d	22,50 c	34,02 c	40,10 e	31,18 c	4,6 d	0,39 a	11,60 c
ITA 10.107.07	391,36 a	284,30 a	21,68 b	13,46 b	29,55 a	13,99 a	18,78 a	153,04 d	61,95 c	15,47 c	22,55 c	39,27 a	47,80 a	20,85 a	4,8 d	0,35 a	15,82 a
ITA 10.107.06	242,83 c	157,09 c	14,57 c	9,42 c	18,01 d	8,85 c	17,56 a	153,93 d	63,97 c	12,07 c	23,90 c	32,06 d	38,42 e	29,63 b	5,7 c	0,45 b	12,87 b
ITA 12.103.04	252,19 c	143,78 c	15,13 c	8,62 c	20,31 c	8,84 c	16,17 b	212,86 c	56,82 d	9,00 b	34,17 d	30,94 d	40,47 d	27,66 a	5,7 c	0,52 c	11,07 c
ITA 12.103.06	209,75 c	168,33 c	12,58 c	10,09 c	13,75 d	9,19 c	18,23 a	235,67 b	79,67 a	9,47 b	10,85 a	30,64 d	38,33 e	26,62 a	5,4 c	0,49 c	10,86 c
ITA 12.103.12	174,98 d	109,91 d	10,50 d	6,59 d	14,35 d	6,33 d	17,39 a	266,44 b	63,00 c	15,80 c	21,25 c	33,66 c	43,35 c	30,85 c	4,6 d	0,43 b	10,63 c
ITA 12.103.15	315,13 b	201,20 b	18,90 b	12,07 b	20,25 c	10,27 b	19,68 a	192,92 c	63,85 c	6,45 a	29,67 d	33,23 c	42,62 c	30,93 c	5,4 c	0,47 c	11,46 c
ITA 12.103.22	419,13 a	292,15 a	25,14 a	17,53 a	30,75 a	18,35 a	17,97 a	304,23 a	69,52 b	11,85 c	18,62 b	36,34 a	46,19 a	25,79 a	7,6 a	0,40 a	16,62 a
ITA 10.128.06	258,77 c	188,78 c	15,52 c	11,32 c	24,33 c	11,65 b	12,44 c	251,65 b	72,90 b	11,22 c	15,87 b	33,96 c	40,60 d	32,09 d	5,9 c	0,54 d	10,78 c
ITA 10.128.09	267,68 c	169,83 c	16,06 c	10,18 c	19,60 c	9,08 c	18,87 a	214,73 c	63,35 c	8,62 b	28,07 c	33,84 c	39,46 e	31,93 d	6,2 c	0,52 c	11,75 c
Average	284,98	181,78	17,09	17,44	21,82	10,61	17,31	220,33	63,46	10,89	25,64	34,23	41,76	31,24	6,2	0,52	12,22
C.V***, %	13,17	3,45**	13,17	7,64**	6,86**	7,78**	3,73**	9,93	6,81	9,36**	5,24**	3,14	2,99	3,21	7,97	2,82**	8,53

* Means followed by the same letter in the column are not significantly different (Scott-knott test, P < 0.05);

** Data transformed by the formula Y = log (x);

*** Coefficient of variation.




Fig. 1. Principal Component Analysis (PCA) of the productive parameters of a) total production per plant (TPP) (g/plant^{-1}); b) commercial production per plant (CPP) (g/plant^{-1}); c) number of total fruits per plant (NTF) (fruits/plant^{-1}); d) number of commercial fruits per plant (NCF) (fruit/plant^{-1}); e) percentage of small fruit production per plant (PSF) (%), and, f) percentage of discard fruits per plant (PDF) (%) of experiments conducted with short days Italian genetic-based strawberry cultivars and genotypes, during the 2019, 2020 and 2021 growth seasons, at the Lages/SC Municipality, southern Brazil




Fig. 2. Principal Component Analysis (PCA) of the fruit quality parameters of a) fruit pulp firmness (FPF) (g/fruit^{-1}); b) epidermis luminosity (L^*); c) epidermis chroma (C^*); d) hue angle ($^\circ\text{hue}$); e) soluble solids/titratable acidity ratio (SS/TA), and f) average fresh mass of commercial fruits (AMCF) (g/fruit^{-1}) of experiments conducted with short days Italian genetic-based strawberry cultivars and genotypes, during the 2019, 2020 and 2021 growth seasons, at the Lages/SC Municipality, southern Brazil

In Fig. 2, both main components were responsible for 72.5% (46.4% plus 26.1%) of the cultivars and genotypes differentiation, according to fruit quality parameters. According to fruit color L^* , C^* , and $^{\circ}$ hue, the first main component was responsible for 46.4% of cultivars and genotypes differentiation. According to fruit flavor SS/TA and AMCF parameters, the second main component was responsible for 26.1% of the differentiation between cultivars and genotypes. The FPF did not show significance for any of the main components (first and second) and was not regarded as a good and reliable parameter in the differentiation of cultivars and genotypes.

The Pircinque cultivar and FRF PIR 256.04, ITA 13,097.05, ITA 12.103.22, ITA 10.107.07, and ITA 10.133.02 genotypes showed the highest productive parameters of CPP, TPP, NCF and NTF (Fig. 1). Among them, the Pircinque cultivar and FRF PIR 256.04 and ITA 12.103.22 genotypes, highlighted in Fig. 2, demonstrated the highest color, flavor, and AMCF parameters, with the highest value for the Pircinque cultivar and FRF PIR 256.04 genotype. Despite having high productive parameters, the ITA 13.097.05, ITA 12.103.22, and ITA 10.133.02 genotypes showed significantly lower fruit color parameters when compared to the Pircinque cultivar and FRF PIR 256.04 and ITA 12.103.22 genotypes that differed by the SS/TA and AMCF parameters. However, the ITA 13.097.05 genotype showed the highest SS/TA and AMCF, indicating a high sweet taste than ITA 12.103.22 and ITA 10.133.02 genotypes (Fig. 2).

4. DISCUSSION

The Pircinque cultivar and FRF PIR 256.04, ITA 10.133.02, ITA 13.097.05, ITA 10.107.07, and ITA 12.103.22 genotypes showed a positive correlation between the productive and fruit quality parameters. This correlation is highly desirable because cultivars and genotypes that combine high production and sweet fruit quality are difficult to find. These findings are consistent with Zanin et al. [12], who identified these cultivars and some of these genotypes for production in some regions of southern Brazil and other climate areas. The Pircinque cultivar and FRF PIR 256.04, ITA 10.133.02, ITA 13.097.05, ITA 10.107.07, and ITA 12.103.22 genotypes demonstrated high NTF and AMCF, which are important parameters for strawberry yield [4, 5].

Strawberry fruit size stability is an important criterion in the plant breeding selection process

for new commercial cultivars. Thus, the AMCF parameter is directly related to fruit size, because large fruits facilitate harvesting and packaging, lowering production and general service costs [3].

The TPT average of the Jonica and Pircinque cultivars, as well as the FRF PIR 256.04 and ITA 12.103.22 genotypes, was 24.35 t ha⁻¹, which was higher than the world TPT average of 22.7 t ha⁻¹ (Table 1) (Fao, 2018). The ITA 12.103.22 genotype progeny (parents) is the cultivar "Florida Fortuna" which has high production, productivity and fruit quality [13]. Zanin et al. [12] obtained better results with these same materials in a conventional system in the Santa Catarina State highland region of southern Brazil. The Jonica cultivar progeny is the "Kilo", a high productive cultivar that was also chosen for its early production and productivity when compared to the other cultivars and genotypes, as these characteristics can increase its productive potentiality and agronomic performance [14,5].

The Italian Pircinque cultivar, introduced in Brazil by the Santa Catarina State University (UDESC), had previously been evaluated and demonstrated a high degree of climate adaptability to the regions of southern Brazil [15,12]. The FRF PIR 256.04 genotype was developed as part of a strawberry breeding project developed at CREA-OFA/Italy with the goal of creating a series of strawberry cultivars with high fruit yield and quality. The FRF PIR 256.04 genotype was previously evaluated in Brazil by Faedi et al. [15], and its high productive performance, good environmental response, and visual fruit quality are supported by the current data.

The Pircinque cultivar and the FRF PIR 256.04, ITA 10.133.02, ITA 13.097.05, ITA 10.107.07, and ITA 12.103.22 genotypes demonstrated a high NTF and AMCF of 18.47 grams. Gallace et al. [16] defines an excellent commercial fruit quality as an AMCF parameter of around 12.0 grams per fruit. Additionally, this former cultivar and genotypes group (Pircinque cultivar and FRF PIR 256.04, ITA 10.133.02, ITA 13.097.05, ITA 10.107.07, and ITA 12.103.22 genotypes), showed the highest fruit red color intensity, especially by the L^* parameter, because L^* parameter expresses the degree of color luminosity ($L^* = 100 =$ white; $L^* = 0 =$ black). In this case, the red values showed by this group, indicated the tendency of the fruits to light red color. The reduction of C^* values indicate a

reduction in the saturation of fruit red pigments, which is less vivid, while the color intensity values ($^{\circ}$ hue) indicated that the red color of the fruits remained similar among the cultivar and genotypes. According to Ávila et al. (2012), lower relative of $^{\circ}$ hue values, indicate darker coloration. Thus, in the present study, the Pircinque cultivar and the FRF PIR 256.04, ITA 10.133.02, ITA 13.097.05, ITA 10.107.07 and ITA 12.103.22 genotypes showed high luminosity and consequently, light epidermis color. The fruit red external color is one of the first factors that consumers take into account in the acquisition of strawberry fruits. Fruits with intense bright red color, but not darkest, are preferable to the public consumers [17]. The epidermis luminosity (L^*) can range from 0 (dark) to 100 (light), instead in the case of C^* or also color purity, the lower of C^* value, higher is the degree color impurity and lower the pigments saturation, while high C^* values increase pigment saturation, generating a pure color [18].

The Pircinque and Sabrina cultivars, as well as the FRF PIR 079.06, FRF PIR 256.04, and FRF PIR 075.08 ITA 12.103.22 genotypes, had the highest fruit SS concentrations, a quantitative feature that may be influenced by climate conditions [19,20]. Plant photosynthesis generates SS with at least 85% sugars in their composition. The higher the SS content, the sweeter are the fruits [14]. The SS content is determined by genetic characteristics [21,19] and is influenced by climate conditions, which explains the differences observed among the cultivars and genotypes studied. Strawberry SS contents variation range from 4.0°–11.0°Brix [19], and all data presented in this study fell within this range.

Pircinque and Sabrina cultivars and the FRF PIR 075.08, ITA 10.107.07 and ITA 12.103.22 FRF PIR 075.08, ITA 10.107.07, and ITA 12.103.22 genotypes, demonstrated the highest SS/TA ratio and, as a result, high fruits sweetness, correlating with the data obtained by et al. [22] and Ahmadi et al. [23], particularly with Pircinque and Sabrina cultivars conducted under different nutritional management. A high SS/TA ratio reflects the quality of fruit flavor/aroma, which is the result of the fruit's sugar content and acidity being balanced. As a result, the higher the fruit SS/TA ratio, the more balanced the flavor/aroma and the greater the sensory fruit acceptance Barankevicz et al., 2015). The fruit SS/TA ratio is more representative than sugars and acidity

alone, allowing for a more accurate assessment of fruit flavor/aroma [24].

Pircinque cultivar and the FRF PIR 256.04, ITA 10.133.02, ITA 13.097.05, ITA 10.107.07, and ITA 12.103.22 genotypes produced not only a large number of fruits, but also a large number of marketable fruits with ideal size and visual quality, making these cultivar and genotypes potential options for commercial fresh fruits. When it comes to purchasing decisions, most consumers prioritize fruit size and appearance over flavor [12]. When developing new cultivars in breeding programs, genotypes that combine fruit quality characteristics with high productivity are always sought. However, this combination is not always easy to achieve, because many of these productive and fruit quality parameters are inversely related depending on the germplasm material available [15,25,26].

5. CONCLUSION

The Pircinque cultivar and FRF PIR 256.04, ITA 10.133.02, ITA 13.097.05, ITA 10.107.07, and ITA 12.103.22 genotypes demonstrated the highest productive and fruit quality parameters in conventional soil systems at low tunnels in edaphoclimatic conditions of highland regions of southern Brazil.

The Pircinque cultivar and FRF PIR 256.04, ITA 10.133.02, ITA 13.097.05, ITA 10.107.07, and ITA 12.103.22 genotypes are potential options for producing cultivars with a high productive parameters, fruit flavor/aroma quality, and fruits SS/TA ratio for the *in natura* market.

The FRF PIR 256.04, ITA 10.133.02, ITA 13.097.05, ITA 10.107.07, and ITA 12.103.22 genotypes were the most promising in terms of fruit production and fruit quality features and may become cultivars recommended for cultivation in southern Brazil, along with Pircinque cultivar.

ACKNOWLEDGEMENTS

The authors thank CREA-OFA/Italy, Capes (Coordination for higher Education Staff Development), Fapesc (Research and Innovation Foundation of Santa Catarina State), Cnpq (National Council for Scientific and Technological Development) for funds and fellowships.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. FAO STAT. Food and Agriculture Organization of United Nation. *Statistical of Yearbook*. Statistical of strawberry production in world; 2021.
2. Heide OM, Stavang JA, Sønsteby A. Physiology and genetics of flowering in cultivated and wild strawberries—a review. *The Journal of Horticultural Science and Biotechnology*. 2013 Jan 1;88(1): 1-8.
Available: <https://doi.org/10.1080/14620316.2013.11512930>
3. Fagherazzi AF, Cocco C, Antunes LE, de Souza JA, Rufato L. La fragolicoltura brasileira guarda avanti. *Rivista di Frutticoltura e di Ortofrutticoltura*. 2014;76(6):20-5.
4. Rivero R, Remberg SF, Heide OM, Sønsteby A. Effect of temperature and photoperiod preconditioning on flowering and yield performance of three everbearing strawberry cultivars. *Horticulturae*. 2022 Jun 6;8(6):504.
Available: <https://doi.org/10.3390/horticulturae8060504>
5. Sønsteby A, Sadojevic M, Heide OM. Production methods for high yielding plants of everbearing strawberry in the Nordic climate. *Horticulturae*. 2022 Mar 15;8(3):249.
Available: <https://doi.org/10.3390/horticulturae8030249>
6. Guimarães AG, Andrade Júnior VC, Elsayed A, YAM, Fernandes JSC, Ferreira MAM. Productive potential of strawberry cultivars. *Revista Brasileira de Fruticultura*. 2015;37(1):112-120.
Available: <https://doi.org/10.1590/0100-2945-400/13>
7. Santos HG. Sistema Brasileiro de Classificação de Solos. 5. ed. Revisão e Amplificação. Brasília, DF: Embrapa; 2018.
8. SBCS. Sociedade Brasileira de Ciência do Solo. Manual de adubação e calagem para os estados de Rio Grande do Sul e Santa Catarina. Xanxerê, SC: Núcleo Regional Sul, Comissão de Química e Fertilidade do Solo – RS/SC; 2016.
9. R Core Team. A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria; 2020.
10. Ferreira DF. Sisvar: a computer statistical analysis system. *Ciência e Agrotecnologia*. 2011;35 (6):1039-1042.
Available: <https://doi.org/10.1590/S1413-70542011000600001>
11. Shepherd GJ. Fitopac. Versão 2.1. Campinas, SP: Departamento de Botânica, Universidade Estadual de Campinas; 2010.
12. Zanin DS, Fagherazzi AF, Santos AM, Martins R, Kretschmar AA, Rufato L. Agronomic performance of cultivars and advanced selections of strawberry in the South Plateau of Santa Catarina State. *Revista Ceres*. 2019 Aug 12;66:159-67.
Available: <https://doi.org/10.1590/0034-737X201966030001>
13. Whitaker VM, Chandler CK, Peres N, do Nascimento Nunes MC, Plotto A, Sims CA. Sensation™ 'Florida127' strawberry. *HortScience*. 2015 Jul 1;50(7):1088-91.
Available: <https://doi.org/10.21273/HORTSCI.50.7.1088>
14. Ariza MT, Soria C, Medina-Mínguez JJ, Martínez-Ferri E. Incidence of misshapen fruits in strawberry plants grown under tunnels is affected by cultivar, planting date, pollination, and low temperatures. *HortScience*. 2012 Nov 1;47(11):1569-73.
Available: <https://doi.org/10.21273/HORTSCI.47.11.1569>
15. Faedi W, Baruzzi G, Lucchi P, Magnani S, Carullo A, Maltoni ML, Migani M, Sbrighi P. The new 'Pircinque' strawberry cultivar released under Italy's PIR Project. *Acta Horticulturae*. 2014;1049(1):961-1966.
Available: <https://doi.org/10.17660/ActaHortic.2014.1049.157>
16. Gallace N, Lieten P, Boonen M, Bylemans D. Reduced winter chill as a means to improve the production potential of late day-neutral strawberry cultivars. *Eur. J. Hortic. Sci*. 2019 Feb 1;84(1):20-3.
Available: <https://doi.org/10.17660/eJHS.2019/84.1.3>
17. Carpenedo S, Antunes LE, Treptow RO. Sensory characterization of strawberry grown in the region of Pelotas (Brazil). *Horticultura Brasileira*. 2016 Oct;34:565-70.
Available: <https://doi.org/10.1590/s0102-053620160417>
18. Castricini A, Dias MS, Martins RN, Santos LO. Strawberries produced in the semi-arid region of Minas Gerais, Brazil: quality of

- the frozen fruit and pulp. Brazilian Journal of Food Technology. 2017;20:1-7.
Available: <https://doi.org/10.1590/1981-6723.14916>
19. Pineli LD, Moretti CL, dos Santos MS, Campos AB, Brasileiro AV, Córdova AC, Chiarello MD. Antioxidants and other chemical and physical characteristics of two strawberry cultivars at different ripeness stages. Journal of Food Composition and Analysis. 2011 Feb 1;24(1):11-6.
Available:<https://doi.org/10.1016/j.jfca.2010.05.004>
20. Costa SI, Ferreira LV, Benati JA, Cantillano RF, Antunes LE. Qualitative parameters of neutral-day strawberries produced in soilless cultivation. Engenharia na Agricultura. 2019;27(6):481-9.
Available:<https://doi.org/10.13083/reveng.v27i6.952>
21. Cocco C, Magnani S, Maltoni ML, Quacquarelli I, Cacchi M, Antunes LE, D'Antuono LF, Faedi W, Baruzzi G. Effects of site and genotype on strawberry fruits quality traits and bioactive compounds. Journal of Berry Research. 2015 Jan 1;5(3):145-55.
Available: <https://doi.org/10.3233/JBR-150098>
22. Fagherazzi AF, Grimaldi F, Kretschmar AA, Molina AR, Gonçalves MA, Antunes LE, Baruzzi G, Rufato L. Strawberry production progress in Brazil. InVIII International Strawberry Symposium 1156 2016; 937-940.
Available:<https://doi.org/10.17660/ActaHort.2017.1156.138>
23. Ahmadi E, Alur AS, Venkat R. Impact of integrated nutrient management on yield and quality parameters of strawberry (*Fragaria× ananassa* Duch.) Cv." Sabrina" under polyhouse. International Journal of Current Microbiology and Applied Sciences. 2017;6(9):3481-7.
Available:<https://doi.org/10.20546/ijcmas.2017.609.427>
24. de Souza DC, Ossani PC, Resende LV, Cirillo MÂ, Lima e Silva LF, Xavier JB. Genetic variability between commercial cultivars and experimental strawberry hybrids with emphasis on multi-factor analysis. Magistra. 2019;30:48-59.
25. Barankevicz GB, Novello D, Resende JT, Schwarz K, Santos EF. Physical and chemical characteristics of tomato hybrids pulp during frozen storage. Horticultura Brasileira. 2015 Jan;33:7-11.
Available: <https://doi.org/10.1590/S0102-053620150000100002>
26. Costa AF, Leal NR, Ventura JA, Gonçalves LS, Amaral Júnior AT, Costa H. Adaptability and stability of strawberry cultivars using a mixed model. Acta Scientiarum. Agronomy. 2015 Oct;37:435-40.
Available:<https://doi.org/10.4025/actasciagron.v37i4.18251>

© 2023 Welter et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/111023>