



Knowledge, Attitudes, and Behavior of Students in Combating Mosquito Nest: A Case Study in High School

Sulistiyawati ^a, Herman Yuliansyah ^b,
Tri Wahyuni Sukesni ^{a*}, Fanani Arief Ghozali ^c,
Surahma Asti Mulasari ^a, Fatwa Tentama ^d,
Bambang Sudarsono ^d, Lu'lu' Nafiati ^e and Nuri Hastuti ^f

^a Faculty of Public Health, Universitas Ahmad Dahlan, Yogyakarta, Indonesia.

^b Faculty of Industrial Technology, Universitas Ahmad Dahlan, Yogyakarta, Indonesia.

^c Faculty of Teacher Training and Education, Universitas Ahmad Dahlan, Yogyakarta, Indonesia.

^d Faculty of Psychology, Universitas Ahmad Dahlan, Yogyakarta, Indonesia.

^e Faculty of Economics and Business, Universitas Ahmad Dahlan, Yogyakarta, Indonesia.

^f Binatama Vocational High School, Sleman, Indonesia.

Authors' contributions

This work was carried out in collaboration among all authors. Author SS was designed this study. Author TWS helped the analysis. And authors HY, FAG, SAM, FT, LN and BB were reviewing the protocol and the instrument. NH assisting the data collection. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJRID/2024/v15i2328

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/113203>

Original Research Article

Received: 14/12/2023

Accepted: 19/02/2024

Published: 24/02/2024

ABSTRACT

Aim: Assessing the relationship between high school student knowledge, attitudes, and behavior of students in eliminating mosquitoes.

Study Design: This research employed a cross-sectional design with participant all students in

*Corresponding author: E-mail: sulistiyawati.suyanto@ikm.uad.ac.id;

Binatama Vocational Health School, Sleman, Yogyakarta.

Place and Duration of Study: This research was done at Binatama Health Vocational School, Sleman, Yogyakarta, in November 2023.

Methodology: Quantitative research with a cross-sectional design was employed. The population and sample in this study were all from Binatama Vocational High School (311 students). An online questionnaire was used to collect the students' responses.

Results: A relationship exists between students' knowledge and behavior in eliminating mosquito nests.

Conclusion: The results of this study indicate no significant relationship exists between knowledge and high school students' attitudes regarding the elimination of mosquito nests. There is a relationship between the knowledge and behavior of high schools in eliminating mosquito nests.

Keywords: knowledge, attitude, behavior, mosquito nests.

1. INTRODUCTION

Mosquitoes are considered the world's most dangerous organisms, causing approximately 750,000 global deaths in 2018 through the spread of mosquito-borne diseases [1]. Mosquitoes are vectors of many infectious diseases caused by parasitic and viral agents such as lymphatic filariasis, chikungunya, and Zika, which cause essential pathologies in humans and animals [2]. The most common viral infection caused by the Aedes mosquito is dengue fever. More than 3.9 billion individuals in more than 129 countries are at risk of contracting dengue fever, with an estimated 96 million symptomatic cases and approximately 40,000 deaths annually [1]. In addition, around 2,400 million people worldwide are at risk of malaria, while dengue fever is estimated to cause 100-400 million infections each [3,4]. Diseases transmitted by mosquitos are endemic in many areas, including Southeast Asia. This disease is a significant cause of death and morbidity in many underdeveloped countries.

Previous research also explains that several factors can influence the incidence of dengue fever, including environmental conditions around the house, such as distance from residential areas, house design, type of water storage container, altitude, and climate. Apart from that, community behavior has a significant impact because it can provide support or create an environment that supports the development of mosquitoes that cause dengue fever. Social factors such as people's habits can be detrimental to health, for example, low compliance with clean and healthy living behavior, such as lack of activity in cleaning water reservoirs and lack of concern for yard cleanliness, as well as low [5].

Indonesia is one country that suffers from dengue disease. This country tries to control dengue spreading through mosquito control, meaning preventing and eradicating dengue fever by breaking the chain of transmission by controlling the vector through the eradication of dengue hemorrhagic fever mosquito nests in Indonesia, called PSN-DBD. Based on previous research, teenagers' role in preventing mosquito-borne diseases such as dengue hemorrhagic fever still needs to be improved [6]. This can be seen from teenagers' lack of participation or involvement in various environmental activities, including cooperation activities, ecological cleaning, and implementation of 3M actions (Drying, Covering, Burying potential breeding places) and Mosquito Nest Eradication (called PSN). Even though teenagers are considered the nation's next generation and pioneers of the reform movement who have significant duties and responsibilities towards their families, environment, and society, their involvement in dengue prevention activities still needs to be increased [7]. To carry out these efforts, teenagers must have good knowledge and understanding. Based on previous research, community behavior, which includes knowledge, attitudes, and actions, is one of the critical factors in successfully eradicating mosquito nests [8]. On the other hand, education and learning about mosquito control can improve youth attitudes and help them adopt protective practices. Previous knowledge, attitudes, and practices (KAP) studies conducted in various regions of Bangladesh regarding malaria and dengue fever have shown knowledge gaps among communities regarding transmission, prevention, or treatment and their reluctance to take preventative measures for mosquito-borne diseases [9].

Binatama Health Vocational School is one of the schools that specializes in the health curriculum. This school is located in Sleman District, Yogyakarta, and is an urban location, making it prone to dengue transmission. With these issues in mind, researchers wish to observe about the knowledge, attitudes, and practices of Binatama Health Vocational School students. This study has significance for offering input to schools and the larger community on KAP that needs to be improved among teenagers.

2. MATERIALS AND METHODS

This research used a cross-sectional design carried out at the Binatama Health Vocational School, Sleman, Yogyakarta. Population in this study was 311 students in this school from first to third grade. During data collection, we included all students as our sample, except students who were absent during data collection for reasoning such as six.

Data was collected via an online form with a set of questionnaires that developed by research group under discussion with the expert. Respondents were given a total of 31 questions, which included 9 knowledge questions, 11 attitude questions, and 11 practice questions. They completed the questionnaire at the same time in their classes, supervised by their teachers. Data were calculated and scored, in knowledge a right and wrong answer were

scored with 1 and 0 respectively. In attitude, a five-Likert scale (very agree, agree, disagree, very disagree and don't know) was posted. Favorable questionnaire was scored as 4,3,2,1,0 respectively – and the vice versa for the unfavorable. In behavior, we provided respondent with agree and disagree questionnaire, agree was scored as 1 and disagree scored as 0. For each knowledge, attitude and behavior were calculated for the total score. Knowledge was categorized as good and poor by using cut off score 75 [10] ; while attitude and practice were using the median. Median for attitude and behavior were 32 and 10 - respectively. The data then analyze using chi-square.

3. RESULTS AND DISCUSSION

In total we included 293 students on this study. Table 1 provides a complete description of the characteristics of teenagers involved in the mosquito nest eradication initiative. The majority of respondents were early adolescents (51.2%). The majority of the respondents were women (93.2%). Most respondents live at home (95.9%), and most have family members of ≤5 people (77.5%). The level of knowledge of respondents is generally high (92.8%). The average attitude of respondents tends to be in the poor category, namely around 59.7%. Apart from that, it is known that most respondents showed poor behavior, as many as 52.9%.

Table 1. Characteristics of respondents

Variable	n	%
Age		
Early teens (14-16 years)	150	51.2
Late teens (17-20 years)	143	48.8
Gender		
Man	20	6.8
Woman	274	93.2
Residence		
House	281	95.9
Boarding house or others	12	4.1
Number of Family Members		
≤5 people	227	77.5
≥6 people	66	22.5
Knowledge level		
Poor	21	7.2
Good	272	92.8
Attitude		
Poor	175	59.7
Good	118	40.3
Behavior		
Poor	155	52.9

Good	138	47.1
------	-----	------

3.1 The Relationship between Knowledge and Attitudes and Behavior in Eradicating Mosquito Nests

Tables 2 and 3 provide valuable insights into the correlation between knowledge attitudes and behavior of adolescents in the context of effort eradication of nest mosquitoes. This table demonstrates the level of understanding teenagers about nest mosquitoes' ties to attitudes toward prevention and tangible conduct that they adopt to reduce the potency of breeding site potential.

Table 2 shows several systematic conclusions that can be drawn. As many as 7.2% of respondents needed to show adequate knowledge regarding eradicating mosquito nests. In this group, the majority, namely 3.8%, showed a hostile attitude regarding these prevention efforts. In contrast, respondents with good knowledge reached 92.8%. However, the majority, namely 56%, showed a less positive attitude towards eradicating mosquito nests. The results of the analysis show exciting findings regarding the relationship between teenagers' knowledge and attitudes. The results of statistical tests with a p-value of 0.630 imply no significant relationship between the level of expertise and teenagers' attitudes toward eradicating mosquito nests.

The results of this study indicate no significant correlation between the knowledge and attitudes

of adolescents regarding the removal of mosquito nests. This finding is in line with the previous research, which also did not find a significant relationship between knowledge and attitudes [11]. This research differs from another one, which revealed a p-value of 0.002 between knowledge and attitudes toward eradicating mosquito nests [12]. Attitudes are identified as significant predictors of behavior, reflecting individual actions, and can be used to predict responses to challenging situations or certain conditions [13].

Other variables beyond knowledge also influence attitudes, such as exposure to information from various sources, including the media, social media, instructors, and parents [14]. Motivation, previous experience, and interactions with family members also form attitudes [15]. Personal experiences, the opinions of essential others, and cultural factors all play a role in creating attitudes [16]. Assume a person is genuinely free from obstacles that could inhibit the presentation of their attitude. In that case, the behavior that appears can be considered a more authentic form of expression. An expression of will or desire can manifest a person's knowledge and awareness of an object, especially in the dengue fever mosquito nest eradication. An individual's desire can be identified as a tendency to carry out an activity [17]. Participatory activities to eradicate mosquito nests significantly impact efforts to prevent and control diseases caused by mosquitoes [16].

Table 2. Relationship between knowledge and adolescents' attitudes in eradicating mosquito nests

Knowledge	Attitude			P-value	PR CI (95%)
	Negative n (%)	Positive n (%)	Total n (%)		
Poor	11 (3.8)	10 (3.4)	21 (7.2)	0.630	0.869 (0.571-1.321)
Good	164 (56)	108 (36.9)	272 (92.8)		
Total	175 (59.7)	118 (40.3)	293 (100)		

Table 3. Relationship between knowledge and adolescent behavior in eradicating mosquito nests

Knowledge	Behavior			P-value	PR CI (95%)
	Poor n (%)	Good n (%)	Total n (%)		
Poor	16 (5.5)	5 (1.7)	21 (7.2)	0.046	1.491 (1.143-1.945)
Good	139 (47.4)	133 (45.4)	272 (92.8)		
Total	155 (52.9)	138 (47.1)	293 (100)		

From Table 3, as many as 7.2% of respondents needed better knowledge regarding eradicating mosquito nests. The majority of them, as many as 5.5%, showed poor behavior in efforts to exterminate mosquito nests. On the other hand, respondents with a good level of knowledge reached 92.8%. However, the majority of them had poor behavior at 47.4%. The results of further analysis show a p-value of 0.046. This value indicates a significant relationship between adolescents' knowledge and behavior in the context of eliminating mosquito nests. There is a relationship between teenagers' knowledge and behavior in eliminating mosquito nests. Based on previous research's findings, the p-value is 0.011, indicating that there is a significant relationship between the level of knowledge and the behavior of Eradicating Mosquito Nests (PSN) in Dengue Fever (DHF) students [18]. Knowledge is identified as a risk factor that influences a person's behavior. This is because a person with a high level of knowledge may take actions outside their understanding [19]. Influencing someone's behavior requires a lot of knowledge. Experience shows that knowledge-based behavior lasts longer than behavior that is not knowledge-based [20].

A greater understanding of mosquito nest elimination can result in more effective and convenient behavior in dealing with this problem, such as mosquito breeding. Assume a person is genuinely free of obstacles that could impede the presentation of their mindset [21]. Therefore, it will impact a better understanding of the importance of controlling mosquito populations [22]. Someone with comprehensive knowledge will understand the need to take preventive steps to reduce the danger of disease transmission by mosquitos. One of the recommended actions is to dry the water reservoir at least once every week. In addition, methods such as carefully sealing water reservoirs after use, recycling used objects, or tossing them away are implemented [23]. In addition, competent people will be better aware of natural and chemical insect-repellent tactics. Utilizing natural mosquito repellents involves maintaining larvae-eating fish, growing mosquito-repelling plants, and using mosquito netting. Meanwhile, chemical prevention includes spreading abate powder, applying insect repellent, and fogging [24].

A person's intention to prevent dengue fever, such as cleaning up puddles of water, sleeping with a mosquito net, and installing mosquito nets

on home vents, is characterized as a response to well-received information [25]. Community behavior related to Mosquito Nest Eradication (PSN) comprises the complete range of activities and attitudes adopted by the community to overcome the problem of mosquito nests, including attempts to prevent and manage mosquito populations [25]. Many factors influence the number of mosquito nests, including gender and age of family members living in the same house. This is because male and female mosquitoes may have different breeding preferences or parenting patterns, which in turn affects the presence of mosquito nests. Therefore, understanding these factors is crucial in designing prevention strategies [26]. Mosquito breeding sites inside and outside the home have been identified as a trigger for an increase in the number of disease vectors carried by mosquitoes [27].

In this context, the 3M Movement, accompanied by the Eradication of Mosquito Nests (PSN), can effectively eradicate the dengue fever virus vector. The PSN 3M plus movement involves various actions, such as drying out stagnant water, closing water storage containers, recycling or reusing used items, spraying larvae-killing larvicide, cultivating larvae-eating fish, changing the water in pots or flower vases, and other efforts [28].

4. CONCLUSION

Based on our finding, it can be concluded that there is no significant relationship between knowledge and attitudes of teenagers regarding eradicating mosquito nests. There is a relationship between knowledge and behavior of teenagers in eradicating mosquito nests.

CONSENT

All authors declare that written informed consent was obtained from the respondent before they participated in this study.

ETHICAL APPROVAL

The study was approved by the Ethical review board of Universitas Ahmad Dahlan, Yogyakarta, Indonesia (ethical approval code: 012311285). All authors hereby declare that all experiments have been examined and approved by the appropriate ethics committee and have therefore

been performed by the ethical standards laid down in the 1964 declaration of Helsinki.

ACKNOWLEDGEMENTS

We thank Universitas Ahmad Dahlan, who funded this research with Grand Number PT-277/SP3/LPPM-UAD/VIII/2023.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. World Health Organization, Vector-Borne Diseases. Accessed: Jan. 31, 2024. [Online]. Available: <https://www.who.int/newsroom/factsheets/detail/vector-borne-diseases>
2. Weaver SC, Charlier C, Vasilakis N, Lecuit M. Zika, chikungunya, and other emerging vector-borne viral diseases. *Annu Rev Med.* 2018;69(1):395-408. DOI: 10.1146/annurev-med-050715-105122.
3. WHO. WHO expert committee on malaria. Web. Accessed: Feb. 02, 2024. [Online]. Available: <https://pubmed.ncbi.nlm.nih.gov/10892307/>
4. [WHO. Dengue and severe dengue, Web.
5. Mawaddah N, Rachmah S, June MSA. The Implementation of eradication of mosquito nest PSN 3M plus program in Jember Region. *Jurnal Kesehatan Komunitas Indonesia.* 2023;3(2):151-161. DOI: 10.58545/jkki.v3i2.128
6. Kosasih CE, Lukman M, Solehati T, Mediani HS. Effect of dengue hemorrhagic fever health education on knowledge and attitudes, in elementary school children in West Java, Indonesia. *Linguistics and Culture Review.* 2021;5(S1):191-200. DOI: 10.21744/lingcure.v5ns1.1349
7. Prastiana Dewi N, Azam M, Factors associated with family PSN-DBD practices in Mulyoharjo Village. *Public Health Perspective Journal.* 2017;2(1):80-88. Available: <https://journal.unnes.ac.id/nju/index.php/php/article/view/11000> (In Bahasa)
8. MRH Farid, MFD Lusno, Maramis MM, Sulistiyawati B, Utomo, Farid AF. Knowledge and attitude relationship with 3M Plus mosquito nest eradication actions in Surabaya: Literature review. *Medical and Health Science Journal.* 2023;7(01):09-19. DOI: 10.33086/mhsj.v7i01.3639
9. ZBM SN, AA RN, Zamri HM. Knowledge, attitude, and practice regarding dengue among students in a public university in Malaysia. *Bangladesh Journal of Medical Science.* 2020;19(2).
10. Arikunto S, *Prosedur penelitian suatu pendekatan praktik (Research Procedures A Practical Approach).* Jakarta: Rineka Cipta; 2013.
11. Sapanova N, Cessna S, DeChano-Cook LM, Childibaev D, Balta N. Kazakhstani high school students' environmental knowledge, attitudes, awareness and concern. *International Research in Geographical and Environmental Education.* 2023;1-18. DOI: 10.1080/10382046.2023.2281697
12. Susanti RDD, Hefniy H, Agustin YD, Nugroho SA. The relationship between knowledge level and the attitude of families of dengue hemorrhagic fever sufferers regarding the eradication of mosquito nests (PSN) in the working area of the curahdam community health center. *Jurnal Keperawatan Profesional.* 2021;9(1):18-35. DOI: 10.33650/jkp.v9i1.2035. (In Bahasa)
13. Kurniawan W, Agustini A. Relationship between Knowledge and Family Attitudes towards Eradicating Dengue Hemorrhagic Fever Mosquito Nests. *Jurnal Health Sains.* 2021;3(2):420-431. (In Bahasa)
14. SIZ Sharif Ishak, Chin YS, MN Mohd Taib, Chan YM, Mohd Shariff Z. Effectiveness of a school-based intervention on knowledge, attitude and practice on healthy lifestyle and body composition in Malaysian adolescents. *BMC Pediatr.* 2020;20(1):1-12. DOI: 10.1186/s12887-020-02023-x
15. VC da C. Rodrigues et al., Factors associated with the knowledge and attitude of adolescents regarding male condom use. *Rev bras enferm. Suppl.* 2021;74(4):1-6. DOI: 10.1590/0034-7167-2019-0452
16. Espiana I, Lestari RM, Ningsih F. The relationship between knowledge and attitudes and community behavior regarding eradicating dengue hemorrhagic fever (DHF) mosquito nests. *Jurnal Surya Medika.* 2022;8(1):129-135. DOI: 10.33084/jsm.v8i1.3454. (In Bahasa)

17. Sartiwi W, Apriyeni E, Sari IK. The relationship between knowledge and attitudes and family behavior regarding the eradication of dengue fever mosquito nests in korong sarang gagak, six lingkungan health center working area, padang pariaman regency. *Jurnal Kesehatan Medika Saintika*. 2019;9(2): 1-48. DOI: 10.30633/jkms.v9i2.232. (In Bahasa)
18. Nursanty O, Bestari RS, Ichsan B, Nurhayani. Kejobong students' level of knowledge, attitudes and behavior in eradicating ddb mosquito nests. The 13th University Research Colloquium Sekolah Tinggi Kesehatan Muhammadiyah Klaten. 2021;460-466, 2021. (In Bahasa)
19. Taniansyah DS et al., Factors associated with the mosquito nest eradication behavior of boarding house cleaning officers in tembalang subdistrict. *Jurnal Kesehatan Masyarakat, Universitas Diponegoro*. 2020;8:707-713. (In Bahasa)
20. Subhi M, Joegijantoro R, Fioner Pulupina F, Widyagama S, Malang H. The relationship between knowledge and community behavior towards filariasis. *Media Husada Journal of Environmental Health*. 2022;2(1):120-127. (In Bahasa)
21. Daryaswanti PI, GAASF Devi, Pendet NMDP, Widyana KAJ. Relationship between knowledge level and community behavior about eradication of mosquito nest. *Babali Nursing Research*. 2021;2(2): 62-68. DOI: 10.37363/bnr.2021.2253
22. Ernawati K et al., Risk factor analysis of community behavior in the eradication of mosquito nests in urban and rural areas. *Ishr*; 2019. 2020;22:49-55. DOI: 10.2991/ahsr.k.200215.010
23. Sutriyawan A, Nofianti N, Halim RD. Factors associated with the incidence of pulmonary tuberculosis. *Health Scientific Journal (JIKa)*. 2022;4(1):98-105, 2022, DOI: 10.36590/jika.v4i1.228
24. Sutriyawan A, Darmawan W, Akbar H, Habibi J, Fibrianti F. Factors that influence the eradication of mosquito nests (PSN) Through 3M plus in efforts to prevent dengue hemorrhagic fever (DBD). *Journal of Science Public Health*. 2022;11(01): 23-32. DOI: 10.33221/jikm.v11i01.936
25. Prasetyo DW, Djohan R, Riani SN, Kunci K. The relationship of knowledge with mosquito nest eradication behavior (PSN) in the community in RW 03, Cempaka Putih Timur Subdistrict, Central Jakarta and review according to islamic views. *The Relationship of Knowledge with Mosquito Nest Eradication Behavior*. 2023;2(1): 67-79.
26. Hamid A, Hamdin H. Relationship between mosquito nest eradication behavior and the habit of using mosquito repellent against dengue hemorrhagic fever in Pekat Village, Sumbawa District, Sumbawa Regency. *Journal of Science Education Research*. 2023;9(1):87-92. DOI: 10.29303/jppipa.v9i1.2035
27. Indarjo S, Suwandono A, Sulistiyani S, Shaluhyah Z. Characteristics of the home environment elementary school students in the filariasis endemic area, Pekalongan City; 2021. DOI: 10.4108/eai.28-4 -2021.2312210
28. Saptaningtyas R, Rahajoe T, Ruri wulan hidayatul maghfiroh, and chyntia rahmawati. Application of larvae hunter applications in efforts to eradicate mosquito nests. *UPI YPTK Scientific Magazine*. 2022;29(2):109-114. DOI: 10.35134/jmi.v29i2.124.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/113203>