



Assessment of Variability for Yield Attributing Traits and Seed Quality Parameters in Mung Bean (*Vigna radiata* L. Wilczek)

Gangalapuram Harini^a, Jai Prakash Gupta^{a*} and Priyanka Gupta^a

^a Department of Genetics and Plant Breeding, School of Agriculture, ITM University, Gwalior, (M.P.)-474002, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2022/v12i1131275

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/92017>

Original Research Article

Received 15 July 2022
Accepted 22 September 2022
Published 28 September 2022

ABSTRACT

The present investigation was undertaken to evaluate ten germplasm of mung bean, used to estimate genetic variability, heritability and genetic advance for 16 characters. The results showed significant differences among the mung bean genotypes in all characters. All the characters had a larger PCV, showing the influence of the environmental variable on their expression. In the present study, PCV is higher than GCV for all the characters under study. High heritability coupled with high genetic advance indicates additive gene action. Heritability for broad sense was estimated for all the studied characters observed range from 33.6 to 98.175%. Genetic advance as percentage of mean is observed range from 2.484% to 26.315%. All these characters coupled with high heritability coupled with high genetic advance can be used for the selection of lines as a parents for further breeding programme, where as in breeding programme these traits were more important for selection.

Keywords: Mung bean; green gram; genetic variability; heritability; genetic advance.

1. INTRODUCTION

Mung bean or green gram is a legume crop with chromosome number $2n=2x=22$. It is belonging to the Fabaceae family, genus *Vigna*, species

radiata, which is mostly grown for its edible seeds and sprouts. It is the third most significant pulse crop farmed in India, behind only red gram and chick pea.

*Corresponding author: E-mail: jaiprakash.soag@itmuniversity.ac.in, jaiprakashgpt28@gmail.com;

India, which accounts for over 70% of the world's manual production of green grams, is the world's largest producer, followed by China and Myanmar. According to the source Ministry of agriculture and farmers welfare, govt of India, (2021-2022), the area of production of green gram in both kharif and rabi season were 51.3 million hectares, and the production was 30.85 million tonnes, with the productivity of 601kg per hectare in 2020-2021. In Madhya Pradesh, total area under green gram production was 5.4 million hectares with the production of 6.41 million tonnes and with the productivity of 1179 kg per hectare in the crop year 2021-2022. Its seed contains 23-28 per cent protein, 3.5-4.5 per cent fiber, 60-65 per cent carbohydrate and 1.0-1.5 per cent fat. Green gram is rich in essential amino acids like lysine, leucine, phenylalanine and isoleucine [1].

Green gram is a legume crop and fixes atmospheric nitrogen with the symbiotic association with *Rhizobium* bacteria and improves soil texture and fertility. It is a short duration crop and it is an outstanding crop to use as an inter crop with major crops like cereals. It can also be grown as green manure crop to enrich the soil. Inter cropping green gram in rice-wheat and rice-rice systems raises the production of these cereal crops and decreases pest incidence.

Genetic variability in a cultivar is the key factor for the further improvement in the yield and also other traits. In green gram large amount of genetic variability has been reported [2], which indicates the potential for genetic improvement. The knowledge of genetic variability present in a breeding material is very important for the selection of superior parents for the initiation of very effective breeding programme [3]. Yield depends on many characters. It is a result of combined effect of many quantitative characters. The evaluation of association between the yield and these quantitative characters gives the minimum knowledge for the selection of characters which influence the yield. Heritability and genetic advance as per cent of mean is an effective method in the selection programme to determine the extent to which various characters can be improved through the selection. The identification of Mung bean genotypes with high yield helps breeder in identifying donors for future breeding programme. With this in view, the current study was conducted with the goal of determining the level of genetic variability in 10 Mung bean genotypes.

2. MATERIALS AND METHODS

The present research on green gram was conducted at the Crop Research Centre, School of Agriculture, Sithouli Campus, ITM University Gwalior, Madhya Pradesh during the *Zaid* season 2022, following the detailed plan of work. The coordinates are 26.140° N, 78.196° E, and the average elevation is 197 meters above sea level. It is part of the Grid Region of Madhya Pradesh.

The materials used for this experimental trial consists of 10 genetically diverse genotypes obtained from the source ICAR - Indian Institute of Pluses Research, Kanpur (U.P.), India viz., IPM-2-3, IPM-2-14, Meha, Soorya, Virat, Varsha, Kanika, Varshudha, Heena and Shikha. The germplasms accessions and checking varieties were sown during *Zaid* 2022. The field experiment was conducted in Randomized Block Design with three replications. Every genotype was space planted with row to row spacing of 30 cm and plant to plant spacing of 5 cm in a plot size of 30x10 (350 cm). Observations were made from the ten competitive plants from each replication that had been marked with tags at various growth phases of the crop plants. 13 characters were used to record the following information: days to 50% flowering, days to maturity, number of primary and secondary branches per plant, height of plant, number of pods per plant, number of clusters per plant, length of pods, number of seeds per pod, 1000 seed weight (g), biological yield per plant, harvest index, and seed yield per plant (g). The lab experiment was conducted in Complete Randomized Block with three replications. Three characters were used to record data viz., seed germination, vigour index length & vigour index mass accordance to Abdul-Baki and Anderson [4].

By using the formula proposed by Burton and De Vane [5], the levels of genotypic and phenotypic genetic variability were estimated for each feature. With the use of percentage values derived by dividing genotypic variation by total phenotypic variance, broad sense heritability was determined [6]. Based on Johnson's [7] formula, the predicted genetic advance for the chosen characters was estimated.

3. RESULTS AND DISCUSSION

The results indicated significant differences among the genotypes for 16 characters studied

in the present investigation. For each of the 16 characters under study, estimates of Analysis of Variance, Genotypic Coefficient of Variation, Phenotypic Coefficient of Variation, Heritability in a broad sense and Genetic advance as a percentage of mean have been provided.

3.1 Analysis of Variance

The analysis of variance was carried out to partition the total variance into all characters show variance due to genotype and other sources. For the following traits: number of primary branches plant⁻¹, number of secondary branches plant⁻¹, number of pods plant⁻¹, number of seeds pod⁻¹, 1000 seed weight, harvest index, vigour index length, and vigour index mass, the mean sum of squares due to genotypes were found to be highly significant ($P > 0.05$ and $P > 0.01$). Days to 50% flowering, days to maturity, plant height, number of clusters plant⁻¹, pod length, biological yield plant⁻¹, and seed germination, on the other hand, were only significant ($P > 0.05$) and showed that the genotypes had a lot of variation (Table 1). There is no significant variation for replication which shows that error due to environmental error was low. These findings are consistent with those reported by Kate et al., [8]; Mwangi et al., [9].

3.2 Phenotypic and Genotypic Coefficients of Variability

Table 2 & Fig. 1 lists the phenotypic and genotypic coefficients of variation for the yield and characteristics under consideration. For all the qualities included in each analysis, the phenotypic coefficient of variation was noticeably larger in size than the genotypic coefficient of variance. To make inferences regarding these characteristics, PCV and GCV were categorized as low (below 10%), medium (10%–20%), and high (beyond 20%).

The estimations of genotypic (GCV) and phenotypic (PCV) coefficients of variation revealed that PCV was greater than GCV for all characteristics, perhaps owing to genotype-environment combination or other environmental variables impacting the appearance of these features. Narrow differences in all variables identified between PCV and GCV showed a larger importance for hereditary components and less effect from the environment. Similar results were obtained by Garg et al., [10]; Yadav et al., [11]; Ramakrishnan et al., [12] for seed yield and its components in green gram.

Medium value of GCV and PCV were recorded for number of primary branches plant⁻¹ (12.795, 18.58), number of secondary branches plant⁻¹ (14.926, 17.418) and harvest index (11.576, 15.22). Low GCV and Moderate PCV recorded for plant height (9.876, 14.918), 1000 seed weight (8.615, 11.806), number of seeds pod⁻¹ (9.303, 12.59), and seed yield plant⁻¹ (6.947, 11.984). Low values of GCV and PCV were recorded for days to 50% flowering (2.395, 3.389), days to maturity (1.822, 2.755), number of pods plant⁻¹ (6.069, 7.284), number of clusters plant⁻¹ (6.532, 9.79), pod length (5.274, 8.407), biological yield plant⁻¹ (5.066, 8.018), seed germination (1.532, 1.825), vigour index length (8.782, 8.863) and vigour index mass (6.099, 6.863). Similar findings were reported by Sana et al., [13] and Shiv et al., [14].

3.3 Heritability

The extent of phenotypic variation relied on the additive gene action is measured by heritability. Heritability for broad sense was estimated for all the characters under study and presented in Table 2 and Fig. 2, with observed range from 33.6 to 98.175 %. High heritability (above 60%) was observed for number of secondary branches plant⁻¹ (73.44), number of pods plant⁻¹ (69.422), seed germination (70.69), vigour index length (98.175) and vigour index mass (78.991). Moderate heritability (30-60%) was observed for traits like days to 50% flowering (38.924), days to maturity (43.765), plant height (43.826), number of primary branches plant⁻¹ (47.422), number of clusters plant⁻¹ (44.514), pod length (39.355), number of seeds pod⁻¹ (54.605), 1000 seed weight (53.245), biological yield plant⁻¹ (39.913), harvest index (57.846) and seed yield plant⁻¹ (33.6). Low heritability (below 30%) haven't observed for any of the traits in the study. Similar results were obtained by Mwangi et al., [9].

3.4 Genetic Advance as Percentage of Mean

Table 2 and Fig. 2 exhibit genetic progress as a percentage of the mean, with the observed range being between 2.484% and 26.315%. The number of secondary branches per plant showed the largest genetic advance as a percentage of mean (above 20%) at 5% selection intensity (26.351). Moderate genetic advance as percentage of mean (10-20%) was recorded for plant height (13.468), number of primary branches per plant (18.151), number of pods per plant (10.416), number of seeds per pod

(14.162), 1000 seed weight (12.95), harvest index (18.136), vigour index length (17.925) and vigour index mass (11.167). Low genetic advance as percentage of mean (below 10%) was recorded for days to 50% flowering (3.078), days to maturity (2.484), number of clusters per

plant (8.978), pod length (6.816), biological yield plant⁻¹ (6.593), seed germination % (2.658) and seed yield per plant (8.296). These results are in agreement with those reported by Garg et al., [9]; Yadav et al., [11]; Ramakrishnan et al., [12]; Mwangi et al., [9].

Table 1. Analysis of variance for seed yield and its components in green gram

Sl. No.	Characters	Mean sum of squares		
		Replication	Genotypes	Error
	Degree of freedom	2	9	18
1	Days to 50% flowering	0.10	5.15*	1.77
2	Days to maturity	0.70	5.79*	1.74
3	Plant height (cm)	17.96	76.21*	22.81
4	No. of primary branches plant ⁻¹	0.15	0.68**	0.18
5	No. of secondary branches plant ⁻¹	0.24	0.67**	0.07
6	No. of pods plant ⁻¹	0.15	3.06**	0.39
7	No. of clusters plant ⁻¹	0.00	0.23*	0.07
8	Pod length	0.07	0.56*	0.19
9	Number of seeds pod ⁻¹	0.23	1.37**	0.30
10	1000 seed weight (g)	17.59	81.98**	18.56
11	Biological yield plant ⁻¹ (g)	0.23	8.83*	2.95
12	Harvest index (%)	1.77	39.89**	7.80
13	Seed germination (%)	0.40	7.26**	0.88
14	Vigour index length	540.15	56424.63**	347.55
15	Vigour index mass	0.10	56.30**	4.58
16	Seed yield plant ⁻¹ (g)	0.019	2.00*	0.79

*=significant at 5% and **=significant at 1% level of probability

Table 2. Estimates of variability components for 16 quantitative traits of 10 green gram genotypes

Characters	Coefficient of Variation		Heritability (bs) (%)	Genetic Advance 5%	Gen. Adv As % of Mean 5%
	GCV	PCV			
Days to 50% flowering	2.395	3.839	38.924	1.364	3.078
Days to maturity	1.822	2.755	43.765	1.585	2.484
Plant height (cm)	9.876	14.918	43.826	5.754	13.468
No. of primary branches plant ⁻¹	12.795	18.58	47.422	0.579	18.151
No. of secondary branches plant ⁻¹	14.926	17.418	73.44	0.791	26.351
No. of pods plant ⁻¹	6.069	7.284	69.422	1.619	10.416
No. of clusters plant ⁻¹	6.532	9.79	44.514	0.323	8.978
Pod length	5.274	8.407	39.355	0.452	6.816
Number of seeds pod ⁻¹	9.303	12.59	54.605	0.909	14.162
1000 seed weight (g)	8.615	11.806	53.245	6.911	12.95
Biological yield plant ⁻¹ (g)	5.066	8.018	39.913	1.822	6.593
Harvest index (%)	11.576	15.22	57.846	5.125	18.136
Seed germination (%)	1.535	1.825	70.69	2.525	2.658
Vigour index length	8.782	8.863	98.175	9.061	17.925
Vigour index mass	6.099	6.863	78.991	7.601	11.167
Seed yield plant ⁻¹ (g)	6.947	11.984	33.6	0.75	8.296

GCV: Genotypic Coefficient of Variation; PCV: Phenotypic Coefficient of Variation




Fig. 1. Genotypic and Phenotypic coefficient of variation estimates of 10 genotypes of Mung bean




Fig. 2. Heritability, genetic advance and genetic advance as per cent mean estimates of 10 genotypes of Mung bean

4. CONCLUSION

Analysis of variability (PCV, GCV) revealed comparable differences between phenotypic and genotypic coefficients of variation indicating less influence of environment on the phenotypic expression of the character. Higher value of GCV and PCV were recorded for number of primary branches per plant, number of secondary branches per plant and harvest index. Moderate PCV and Low GCV recorded for plant height, 1000 seed weight, number of seeds per pod, and seed yield per plant. Low values of GCV and PCV were recorded for days to 50% flowering, days to maturity, number of pods per plant, number of clusters per plant, pod length, biological yield per plant, seed germination, vigour index length and vigour index mass. High heritability coupled with high genetic advance indicates additive gene action, which is beneficial for crop improvement. The characters, no. of secondary branches per plant, no. of pods per plant, seed germination, vigour index length, and vigour index mass show high heritability coupled with high genetic advance. Meha and shikha varieties are high in heritability and genetic advance, can be select as a parents for further breeding program.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Lambrides CJ, Godwin I. Mung bean in: Genome mapping and molecular breeding in plants: Pulses, sugar and tuber crops (Edited by Kole C), Springer, Berlin and Heidelberg. 2007;3:69-90.
- Moulick D, Samanta S, Saha B, Mazumder MK, Dogra S, Panigrahi K, Banerjee S, Ghosh D, Santra SC. Salinity stress responses in three popular field crops belonging to fabaceae family: Current status and future prospect in the plant family fabaceae. Springer, Singapore. 2020;519-541.
- Sakthivel G, Jeberson S, Singh NB, Sharma PR, Kumar S, Jalaj VK, Sinha B, Singh NO. Genetic variability, correlation and path analysis in lentil germplasm (*Lens culinaris* Medik.). The pharma innovation journal. 2019;8(6):417-420.
- Abdul Baki AA, Anderson JD. Vigor determination in soybean seed by multiple criteria. Crop science. 1973;13(6):630-633.
- Burton GW, De Vane EH. Estimating heritability in tall fescusce from Replicated clone natural materials. Agronomy Journal. 1953;45:171-181.
- Hanson GH, Robinson HF, Comstock RE. Biometrical studies of yield in segregating populations of Korean Lespedeza. Agronomy Journal. 1956;48:268-272.
- Johnson HW, Robinson HF, Comstock RE. Estimates of genetic and environmental variability in soybean. Agronomy Journal. 1955;47:314-318.
- Kate AM, Dahat DV, Chavan BH. Genetic variability, heritability, correlation and path analysis studies in green gram [*Vigna radiata* (L.) Wilczek]. International Journal of Development Research. 2017;7(11):16704-16707.
- Mwangi JW, Okoth OR, Kariuki MP, Piero NM. Genetic and phenotypic diversity of selected Kenyan mung bean (*Vigna radiata* L. Wilczek) genotypes. J Genet Eng Biotechnol. 2021;19:142.
- Garg G, Verma PK, Hari Kesh. Genetic variability, correlation and path analysis in mungbean [*Vigna radiata* (L.) Wilczek]. International Journal of Current Microbiology and Applied Sciences. 2017; 6(11):2166-2173.
- Yadav R, Ram BJ, Suresh BG, Lavanya GR. Studies on genetic variability, character association for yield and yield components in green gram [*Vigna radiata* (L.) Wilczek]. Journal of Pharmacognosy and Phytochemistry. 2017;6(4):1988-1991.
- Ramakrishnan CKD, Savithramma DL, Vijayabharathi A. Studies on genetic variability, correlation and path analysis for yield and yield related traits in Greengram [*Vigna radiata* (L.) Wilczek]. International Journal of Current Microbiology and Applied Sciences. 2018; 7(3):2753-2761.
- Sana AY, Pithia MS, Raval LJ, Vora ZN. Genetic variability, heritability & genetic advance for seed yield and its components in F2 generations of mung bean [*Vigna radiata* (L.) Wilczek]. International Journal of Pure and Applied Bioscience. 2017;5(2):532-535.

14. Shiv A, Ramtekey V, Vadodariya GD, Modha KG, Patel RK. Genetic variability, heritability & genetic advance in F3 progenies of mung bean [*Vigna radiata* (L.) Wilczek]. International journal of pure and applied bioscience. 2017;6(12):3086-3094.

© 2022 Harini et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/92017>