

Modelling the Relationship between Weather Variables and Rice Yellow Stem Borer Population: A Count Data Modelling Approach

Bojjareddy Nanda Kumar Reddy ^a, Santosha Rathod ^{b*}, Supriya Kallakuri ^b,
Yerram Sridhar ^b, Meena Admala ^a, Seetalam Malathi ^c, Prमित Pandit ^d
and Bellamkonda Jyostna ^a

^a College of Agriculture, Professor Jayashankar Telangana State Agricultural University,
Hyderabad 500030, India.

^b ICAR-Indian Institute of Rice Research (IIRR), Hyderabad, 500030, India.

^c PJTSAU-Regional Agricultural Research Station, Warangal 506007, India.

^d Bidhan Chandra Krishi Viswavidyalaya, Nadia 741252, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2022/v12i111410

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/93244>

Original Research Article

Received 05 September 2022

Accepted 01 November 2022

Published 04 November 2022

ABSTRACT

Aim: This study was conducted to model the relationship between discrete dependent variable (yellow stem borer population) and continuous weather variables.

Data Description: The yellow stem borer (YSB) population and standard meteorological week (SMW) wise weather variables (temperature, relative humidity, rainfall and sunshine hours) data of Warangal centre (Telangana state) generated under All India Co-Ordinated Rice Improvement Project (AICRIP) from 2013-2021 were considered for the study. The YSB population were recorded daily using light trap with an incandescent bulb and are counted as weekly cumulative catches.

Methodology: The weekly cumulative trapped YSB populations and weekly averages of climatological data were considered as inputs to the models under consideration. In this study the classical linear regression i.e. step-wise multiple linear regression and count regression models such as Poisson, negative binomial, zero inflated Poisson and zero inflated negative binomial regression models were employed.

Result: The empirical results revealed that the zero inflated count regression models viz., zero inflated Poisson regression and zero inflated negative binomial regression models performed better compared to the classical linear regression, Poisson and negative binomial regression models, further the negative binomial regression model outperformed all models as it yielded lowest mean square error (MSE) and highest R^2 values. The average percentage reduction in accuracy of zero-inflated negative binomial regression model over classical model was around 4 percent.

Conclusion: Based on the results obtained in this study, it is concluded that the zero inflated models performs better compared to classical models as they are unable to handle the presence of excess zeroes, as a result provides more prediction error and lower R^2 values. Further, the models developed in this study will be of great assistance in identifying the factors influencing occurrence of YSB population in rice.

Keywords: Yellow stem borer; light trap catches; weather variables; linear regression, count regression models; zero inflated count regression models.

1. INTRODUCTION

“Rice (*Oryza sativa*) is the most staple food crop of India and Asian continent. India has the largest rice-growing area in the world, covering an area of 44 million hectares and producing 112 million tons of rice” [1]. The insect pest complex of rice crop has undergone a drastic change during the last three decades following green revolution. The losses in rice yield are due to moderate to severe incidence of stem borer, gundhi bugs, hoppers and other insect pests in India. The intensity of insect pest damage varies in different seasons, years and agro-climatic zones due to variability in weather parameters and biotic factors. Yellow Stem Borer (YSB) *Scirpophaga incertulas* Walker has emerged as one of the most important pests of rice during post green revolution years throughout the country and damage of YSB can lead to about 20 % yield loss in early planted rice crops and 70% in late planted crops. The YSB larva lives inside the rice stem and grows for a period of 40 days and pupates inside the stem in a white silken cocoon and later becomes adult YSB. The YSB larvae bore at the base of the plants during the vegetative stage. On older plants, they bore

through the upper nodes and feed toward the base. The larva YSB, adult YSB and damage symptoms of yellow stem borer infected rice are depicted in Fig. 1.

“Both biotic and abiotic factors are believed to be responsible for pest population dynamics” [2]. “Besides inherent biotic potential of the insect to a large extent abiotic factors like temperature, rainfall, relative humidity, sunshine hours etc. determine the abundance of insect pests in a crop ecosystem. An efficient forewarning system based on robust statistical model to predict YSB population dynamics is of great importance in designing and implementation of effective location specific pest management strategies to avoid yield losses” [3]. Count regression modelling is a popular approach employing in understanding the relationship between the discrete dependent count variables and continuous or discrete exogenous variables. Crop pest modelling is one of the major areas of count data regression modelling wherein daily or weekly counts of insects (pests), are considered as dependent variables and corresponding weather variables are considered as exogenous variables.




Fig. 1. (a) Larva yellow stem borer (b) Adult yellow stem borer (c) Symptoms of yellow stem borer infected rice

Though the count regression data modelling approach is applied in many areas; Identifying defects in manufacturing [4], In domestic violence data [5], caries research [6], spontaneous abortion [7], modelling of death due to covid-19 in Ghana [8] and zoological data [9] but their applications are limited in insect pest modeling. Contrary to count regression modelling, the continuous linear regression model was applied in many insect pest modelling studies and as well as in agricultural data sets; weather based regression modelling of white rust disease of Indian mustard [10], regression modelling factors influencing agricultural diversification of Karnataka [11], regression modelling of biometrical characters in Sesame [12], regression analysis of growth patterns of root and shoot morphological traits in Sesame [13], weather based YSB regression model [14], modelling of YSB population under changing climatic scenarios [15].

Crop pest modelling is one of the popular areas of discrete regression modelling wherein daily or weekly counts of insects or pests which are considered dependent variable. As the YSB population will not occur regularly on daily basis there use to be many zeroes in weekly count data under such condition the classical regression and count regression models may not yield better results alternatively zero excess count regression models can be used to model the data with excess zeroes. With this background, this study was undertaken to develop zero excess count regression models using data driven approaches. Further, the paper is arranged in different sections; the data description and statistical models used viz., stepwise regression model, Poisson regression, Negative binomial regression model and their counter parts such as zero inflated poisson regression model and zero inflated negative binomial regression models are explained in methodology section. The results obtained under each section are depicted and discussed in results and discussion section and finally the outcome of the work are highlighted in conclusion section.

2. MATERIAL AND METHODS

2.1 Data Description

The weekly cumulative catches of YSB population was recorded from light trap installed at All India Coordinated Rice Improvement Project (AICRIP), PJTSAU-Regional Agricultural

Research Station, Warangal centre. Corresponding climatological data on maximum temperature (MAXT), minimum temperature (MINT), total rainfall (RF), morning relative humidity (RHM), evening relative humidity (RHE) and sunshine hours (SSH) were also collected from weather stations installed at Warangal centre. Standard meteorological week (SMW)-wise cumulative catches of YSB population and weekly averages of climatological parameters from 2013-2021 were considered for this study. The required data for this study were taken from AICRIP ICAR-Indian Institute of Rice Research Hyderabad. The study area is depicted in Fig. 2.

2.2 Statistical Models

In this work statistical modelling framework begin with summary statistics where data distribution and heterogeneity among the data in terms of coefficient of variation (CV) is explained. The SMW-wise data from 2013-2021 is depicted graphically with time series plots. The correlation analysis was carried out to understand the interrelationship between the YSB population and other exogenous variables. Further, the stepwise multiple linear regression (MLR) was carried out know the cause and effect relationship between YSB population and independent weather variables.

The regression equation in terms of matrix notation can be expressed as;

$$Y = X\beta + e \quad \dots (1)$$

where, Y is the dependent variable, X is the vector of exogenous variables, β is the regression coefficient vector, and e is the residuals term.

Correlation analysis and stepwise MLR analysis were carried out in SAS version 9.3 available at ICAR- Indian institute of Rice research Hyderabad [16]. The count regression models were built in R software [17].

2.2.1 Count regression models

The count data observations following generalized linear model (GLM) framework is elaborated in terms of Poisson and Negative binomial regression models as follows;

Poisson Regression Model: The Poisson regression model is derived from Poisson distribution where mean and variances are equal.

A discrete random variable Y is said to follow Poisson distribution with parameter $\lambda > 0$, if its probability mass function (pmf) is given as follows;

$$f(Y = y) = P_r(Y = y) = \frac{e^{-\lambda} \lambda^y}{y!} \quad \dots (2)$$

where the mean and variance expression of the distribution is $E(y) = \sigma^2_y = \lambda$, changes in means also influence the changes variance [4]. The Poisson regression model is expressed as follows;

$$y_i = \exp(X_i' \beta) + \varepsilon_i \quad \dots (3)$$

The parameters of Poisson regression models are estimated using maximum likelihood estimation method.

Negative Binomial Regression Model: The Negative binomial regression is a class of generalized linear model where the discrete dependent variable Y is over dispersed. This

model is applied to over dispersed count data or to the count data where conditional variance is greater than conditional mean. The negative binomial distribution has same mean structure as Poisson distribution and has additional over distribution parameter ϕ .

Let Y be a random variable follows negative binomial distribution with parameters (r, θ) , where $\theta \in (0, 1)$ and r an integer, then its pmf is given as

$$P(Y = y) = \binom{y+r-1}{y} \theta^y (1-\theta)^r, y = 0, 1, 2$$

$$y \sim \text{NegBin}(r, \theta) \quad \dots (4)$$

$$E(Y) = \frac{r\theta}{(1-\theta)} \text{ and } Var(Y) = \frac{r\theta}{(1-\theta)^2}$$

If the conditional distribution of the outcome variable is over-dispersed, then confidence interval of the negative binomial distribution is shorter than the Poisson distribution [6].




Fig. 2. Location map of Warangal centre (study area)

2.2.2 Zero inflated count regression models

When the count data observations contain excess zeroes in the data sets, then the regular count regression models unable to estimate the parameters properly, then one has to switch to the model which considers the excess zero in the data, called as zero inflated count regression models [18].

In zero inflated Poisson regression (ZIPAR) model, the counts $Y_i=0$ with probability p_i and $1-p_i$ which follows Poisson distribution with mean μ_i where $i=0,1,2,\dots,n$. The zero inflated Poisson regression model is a mixture of two components i.e. zero and no-zero components.

$$P(Y = y) = \begin{cases} p_i + (1 - p_i)e^{-\mu_i} & y_i = 0 \\ (1 - p_i) \frac{e^{-\mu_i} \mu_i^{y_i}}{y_i!} & , y_i = 1,2,3 \end{cases} \dots(5)$$

The zero inflated negative binomial regression (ZINBR) model is a mixture distribution which assigns p to 'extra' zeros and a mass of $(1 - p)$ to a negative binomial distribution, where $0 \leq p \leq 1$. The probability mass function for ZINB is expressed as follows;

$$P(Y = y) = \begin{cases} p + (1 - p) \left(\frac{\phi}{\mu + \phi}\right)^\phi & y = 0 \\ (1 - p) \frac{\tau(y + \phi) \left(\frac{\phi}{\mu + \phi}\right)^\phi \left(\frac{\mu}{\mu + \phi}\right)^y}{\tau(y + 1) \Gamma(\phi)} & , y = 1,2,3 \end{cases} \quad (6)$$

Where, ϕ^{-1} is dispersion, μ is mean and $\Gamma(\cdot)$ is gamma function respectively. The ZIPAR and ZINBAR models are represented in terms of logistic regression function and negative binomial regression function in following equation;

$$Logit(p_i) = x_i^T \beta \text{ and } Logit(\mu_i) = Z_i^T \gamma$$

where β are the vector coefficients X_i^T and γ are the vector coefficients Z_i^T .

3. RESULTS AND DISCUSSION

The weekly (SMW wise) counts of yellow stem borer light trap catches at Warangal centre observed during 2013–2021 period was plotted in Fig. 3. The yellow stem incidence was higher between 36th to 45th SMWs every year.

3.1 Summary Statistics

Summary statistics of the dependent variable i.e. yellow stem borer population and exogenous weather variables were calculated and presented in Table 1. The YSB population data under consideration has high percentage of CV and an abnormality of data as skewness and kurtosis are out of normal range. Summary statistics of weather variables presented in Table 1 were self-explanatory, showing that data under consideration were highly heterogeneous in nature.




Fig. 3. YSB population of Warangal centre

Table 1. Descriptive statistics of Yellow Stem Borer of Warangal Centre

Location	Statistics	Population	MAXT	MINT	RF	RHM	RHE	SSH
Warangal	Mean	28.2	32.4	20.6	8.1	97.07	56.2	6.1
	S. E	1.7	0.20	0.20	1.05	0.2	0.52	0.12
	Skewness	2.01	1.02	-0.52	5.2	-2.33	0.22	-0.67
	Kurtosis	5.1	0.24	-0.74	32.5	11.9	-0.43	-0.49
	Minimum	0	25.7	11.2	0	53.3	28	0
	Maximum	266	45.9	31	204.7	97.7	94	11.1
	CV (%)	135.8	13.6	21.1	278.4	5.1	20.27	44.3

MAXT: maximum temperature, MINT: minimum temperature, RF: rainfall RHM: morning relative humidity, RHE: evening relative humidity, SSH: sunshine hours

Table 2. Pearson correlation coefficients between YSB and weather variables

	YSB	MAXT	MINT	RF	MRH	SSH
MAXT	-0.14348 (0.0019)					
MINT	-0.19489 ($< .0001$)	0.59858 ($< .0001$)				
RF	-0.05397 (0.2444)	-0.19653 ($< .0001$)	0.14864 (0.0013)			
MRH	0.11316 (0.0144)	-0.22146 ($< .0001$)	-0.04595 (0.3218)	0.05786 (0.2120)		
ERH	-0.00137 (0.9765)	-0.22175 ($< .0001$)	0.43084 ($< .0001$)	0.32459 ($< .0001$)	0.48630 ($< .0001$)	
SSH	0.18990 ($< .0001$)	0.45273 ($< .0001$)	-0.01842 (0.6914)	-0.39359 ($< .0001$)	-0.15700 (0.0007)	-0.47375 ($< .0001$)

Values in parentheses represent probability values

Table 3. Stepwise Regression Analysis of yellow stem borer of Warangal centre

Parameter	Estimate	S.E.	F Value	Probability	R ²	Model R ²
Intercept	29.41	19.75	2.22	0.1371		
MINT	-1.86	0.69	18.36	$< .0001$	0.03	0.03
SSH	4.94	0.77	17.38	$< .0001$	0.03	0.07
ERH	0.75	0.21	25.30	$< .0001$	0.04	0.04
MAXT	-1.07	0.67	2.52	0.1128	0.004	0.12

3.2 Correlation Analysis

Pearson correlation coefficients between YSB populations and considered climatological variables are depicted in Table 2. A low negative significant correlation between YSB and MAXT & MINT was observed; a low positive significant correlation was observed between YSB population and MRH & SSH. The ERH and Rainfall data had non-significant correlation with YSB population at this centre. The bivariate correlation coefficients between weather variables are given in Table 1 are self-explanatory.

3.3 Stepwise Regression Analysis

To identify the factors influencing the incidence of YSB population and weather variables a step-

wise linear regression analysis was carried out in SAS 9.3 version available at ICAR-Indian Institute of Rice Research Hyderabad. Results of the stepwise regression analysis are depicted in Table 3. For the dependent variable YSB population, MINT, SSH and ERH are significantly contributing, further MINT have negative effect on YSB population and SSH and ERH have positive impact on YSB population for the data under consideration. Similar results were found in Prasannakumar et al. [2]. Though the listed variables had significant influence on the Yellow Stem Borer populations, the model R² value for the fitted regression in the Warangal centre is low, indicating that the model is not a strong fit, for which non-linearity and high heterogeneity in dependent variables may be responsible.

3.4 Results of Count Regression Models

As explained in methodology section, Poisson and negative binomial regression models were fitted for the YSB population and corresponding weather variables are also considered to build the model.

The model parameters for poisson regression model were given in Table 4 for the Warangal centre and all the weather parameters are significantly contributing to the YSB population incidence.

Similarly, negative binomial regression model was also fitted for the YSB population and weather variables, where MINT, RHM and SSH showed significant influence on the dependent variable where as other variables shown non-significant relationship with the YSB population.

3.5 Results of Zero Inflated Count Regression Models

An alternative way for modelling the data contains more zeroes by using zero-inflated count regression models. In this article, we have employed Zero-inflated Poisson regression and zero-inflated negative binomial regression models and their parameter estimates are given in Tables 6 and 7 respectively. Table 6 provides parameter estimation of ZIPR model, in which most of the weather parameters are significant for nonzero outcomes as probability of significance is $p < 0.0001$ and for zero outcomes MINT, RF, MRH and SSH are significant.

Table 7 provides parameter estimation of ZINBR model for both zero and non-zero outcomes. Parameters like ERH and SSH are significant for non-zero outcome model and RF, MRH and SSH are significant for zero outcome ZINMR models.

Table 4. Parameter estimation of Poisson regression model

Warangal	Estimate	Std. Error	Z value	Probability
Intercept	0.22	0.25	0.875	0.382
MAXT	-0.037	0.0035	-10.390	<0.0001
MINT	-0.065	0.003	-17.625	<0.0001
RF	0.001	0.0004	4.077	<0.0001
RHM	0.032	0.002	11.080	<0.0001
RHE	0.026	0.001	18.984	<0.0001
SSH	0.202	0.004	41.556	<0.0001

Table 5. Parameter estimation of negative binomial regression model

Parameter	Estimate	Std. Error	z value	Probability
Intercept	-3.00	1.74	-1.72	0.08
MAXT	-0.01	0.02	-0.58	0.55
MINT	-0.08	0.03	-2.78	<0.0001
RF	0.009	0.005	1.03	0.30
RHM	0.07	0.01	3.80	<0.001
RHE	0.01	0.01	1.40	0.15
SSH	0.18	0.03	5.40	<0.0001

Table 6. Parameter estimation of zero inflated poisson regression model

Parameter	Estimate	P(Y>0)			P(Y=0)			
		Std. Error	Z value	Prob.	Estimate	Std. Error	Z value	Prob.
Intercept	0.92	0.27	3.32	<0.0001	-1.92	3.38	-0.56	0.57
MAXT	-0.037	0.0035	-9.56	<0.0001	-0.04	0.06	-0.78	0.43
MINT	-0.033	0.0038	-9.80	<0.0001	0.42	0.09	4.56	<0.0001
RF	-0.0021	0.005	-4.55	<0.0001	-0.03	0.01	-2.89	<0.0001
MRH	0.02	0.0032	6.41	<0.0001	-0.07	0.03	-1.98	0.04
ERH	0.036	0.0014	22.06	<0.0001	0.01	0.02	0.60	0.54
SSH	0.14	0.0048	29.47	<0.0001	-0.29	0.06	-4.34	<0.0001

Table 7. Parameter estimation of zero inflated negative binomial regression model

Parameter	P(Y>0)				P(Y=0)			
	Estimate	Std. Error	Z value	Probability	Estimate	Std. Error	Z Value	Probability
Intercept	-0.090	1.96	-0.04	0.96	-5.20	5.38	-0.96	0.333
MAXT	-0.018	0.02	-0.72	0.47	-0.02	0.08	-0.25	0.80
MINT	-0.050	0.02	-1.72	0.08	0.55	0.16	3.418	0.005
RF	-0.001	0.004	-0.46	0.64	-0.04	0.01	-2.28	0.02
MRH	0.028	0.02	1.27	0.20	-0.11	0.05	-2.08	0.03
ERH	0.033	0.01	3.07	0.005	0.05	0.03	1.59	0.11
SSH	0.133	0.02	4.67	P<0.001	-0.32	0.09	-3.50	0.001

Table 8. Comparison of performance of different models at Warangal centre

Warangal	SRM	PRM	NBRM	ZIPRM	ZINBRM
R ²	0.125	0.145	0.113	0.172	0.176
MSE	1320.62	1256.85	1313.18	1219.39	1275.99
RMSE	36.34	35.45	36.24	34.92	35.72

SRM: Stepwise regression model, PRM: Poisson regression model, NBRM: Negative binomial regression model, ZIPRM: Zero inflated Poisson regression model, ZINBRM: Zero inflated negative binomial regression model




Fig. 4. Actual vs fitted values of models

The results of modelling and predictions of the yellow stem borer population at Warangal centre obtained by applying different models were compared in terms of R², MSE and RMSE and are presented in Table 8.

In this study it is revealed that stepwise regression model was weak (low R²) due to non-linearity and heterogeneity in the dependent variable. different models applied in this study viz., Poisson regression model, negative binomial regression model, zero inflated Poisson regression model and zero inflated negative binomial regression model. The zero inflated count regression models outperformed the classical

models as their R² is high and MSE and RMSE is low, further ZINBR model outperformed all the models as it has highest R² value and lowest MSE and RMSE values. The better performance of ZINBR model could be due to its ability to capture over dispersion and heterogeneity in the data sets. Similar results were also found in Tawiah et al. [8] where zero-inflated models performed better than classical models in modelling COVID-19 deaths in Ghana country. Further, the actual vs fitted values of all the models are depicted in Fig. 4 which is clearly indicates, ZINBR model fitted values are closer to the actual YSB population values.

4. CONCLUSION

In the present study, different regression models were developed to model the relationship between weather variables namely maximum temperature (MAXT), minimum temperature (MINT), total rainfall (RF), morning relative humidity (RHM), evening relative humidity (RHE) and sunshine hours (SSH). The results showed that stepwise linear regression model unable to capture the pattern present in the heterogeneous count data as its yielded high RMSE values whereas count regression model are better in identifying the relationship between weather variables and YSB population. Among all the models implemented, ZINBR model performed better compared to all the models. The data under consideration was over dispersed, possibly due to two facts that are excess of zeros present in the data and heterogeneous nature of data. The classical regression models unable to model such data under consideration, an alternative way for modelling such type of data is by using the zero-inflated regression model which considers the excess of zeroes.

Rice yellow stem borer is most serious pest in rice cultivation, causing significant economic losses in rice ecologies across the country. The models developed in this study utilizing regression techniques will be of great assistance in identifying the relationship between weather variables and occurrence of YSB population, so that the appropriate management measures can be engaged to minimize the yield losses. In the future, it is expected that various count regression techniques will be intensively used to model the data of other crop pests.

ACKNOWLEDGEMENT

This is part of the first author's post graduate thesis work at Professor Jayashankar Telangana State Agricultural University. The authors would like to thank ICAR-Indian Institute of Rice Research Hyderabad and PJTSAU Hyderabad for their support in this research work.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Agricultural Statistics at Glance; Ministry of Agriculture & Farmers Welfare Department of Agriculture; Cooperation & Farmers Welfare Directorate of Economics & Statistics; Government of India; 2021.

Available: [https://eands.dacnet.nic.in/PDF/Agricultural%20Statistics%20at%20a%20Glance%20-%202021%20\(English%20version\).pdf](https://eands.dacnet.nic.in/PDF/Agricultural%20Statistics%20at%20a%20Glance%20-%202021%20(English%20version).pdf)
 Accessed on 15 May 2022

2. Prasannakumar NR, Chander S, Kumar LV, Moral MT. Development of weather-based rice yellow stem borer prediction model for the Cauvery command rice areas, Karnataka, India. *Cogent Food & Agriculture*. 2015;1(1):995281. DOI: 10.1080/23311932.2014.995281
3. Ho GTT, Le CV, Nguyen TH, Ueno T. Incidence of yellow rice stem borer (*Scirpophaga incertulas*) Walker in Haiphong, Vietnam and control efficacy of egg mass removal and insecticides. *Journal- Faculty of Agriculture Kyushu University*. 2013;58(2):301-306.
4. Lambert D. Zero-inflated poisson regression, with an application to defects in manufacturing. *Technometrics*. 1992;34(1): 1–14.
5. Felix F, Singh KP. Zero-inflated generalized poisson regression model with an application to domestic violence data. *J Data Sci*. 2006;4(1):117–130.
6. Mwalili SM, Lesafre E, Declerck D. The zero-inflated negative binomial regression model with correction for misclassification: an example in caries research. *Stat Methods Med Res*. 2008;17(2):123–139.
7. Verma P, Swain PK, Singh KK. Count data regression modeling: an application to spontaneous abortion. *Reprod Health*. 2020;17:106. Available: <https://doi.org/10.1186/s12978-020-00955-2>
8. Tawiah K, Iddrisu WA, Asampana AK. Zero-Inflated time series modelling of COVID-19 deaths in Ghana. *Journal of Environmental and Public Health*; 2021.
9. Ilknur Ozmen, Felix Famoye. Count regression models with an application to zoological data containing structural zeros. *J. Data Sci*. 2022;5(4):491-502. DOI: 10.6339/JDS.2007.05(4).385
10. Basavaraj K, Rathi AS, Gaurav NP, Anil Kumar, Santosha Rathod, Ram Niwas. Influence of weather variables on progression of white rust disease of Indian mustard. *Journal of Agrometeorology*. 2020;22(3):362-366.
11. Rathod S, Surendra HS, Munirajappa R, Chandrashekar H. Statistical assessment on the factor influencing agricultural diversification in different districts of

- Karnataka. Environ. Ecol. 2011;30(3A): 790-794.
12. Narayanaswamy T, Surendra HS, Rathod S. Multiple stepwise regression analysis to estimate root length, seed yield per plant and number of capsules per plant in Sesame (*Seasamum indicum* L.). Mysore J. Agric. Sci. 2012a;46(3):581-587.
 13. Narayanaswamy T, Surendra HS, Rathod S. Fitting of statistical models for growth patterns of root and shoot morphological traits in Sesame (*Seasamum indicum* L.). Environ. Ecol. 2012b;30(4):1362-1365.
 14. Prasannakumar NR, Subhash Chander L, Vijay Kumar, Manuel Tejada Moral (Reviewing Editor). Development of weather based rice yellow stem borer prediction model for the Cauvery command rice areas, Karnataka, India. Cogent Food & Agriculture. 2015;1(1). DOI: 10.1080/23311932.2014.995281
 15. Nurhayati E, Koesmaryono Y, Impron. Predictive modelling of rice yellow stem borer population dynamics under climate change scenarios in Indramayu. IOP Conf. Ser.: Earth Environ. Sci. 2017;58: 012054.
 16. SAS Software, Version 9.3; SAS Institute: Cary, NC, USA; 2011.
 17. R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria; 2022. Available: <https://www.R-project.org/>
 18. Workie MS, Azene AG. Bayesian zero-inflated regression model with application to under-five child mortality. J Big Data. 2021;8, 4. Available: <https://doi.org/10.1186/s40537-020-00389-4>

© 2022 Reddy et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/93244>