



## Molecular Characterization of Fungi Associated with Dump Site Soil

**N. G. Ogbuji<sup>1,2\*</sup>, A. E. Ataga<sup>1,2</sup>, P. M. Tari-Ukuta<sup>2</sup> and C. J. Olisedeme<sup>1</sup>**

<sup>1</sup>Department of Plant Science and Biotechnology, University of Port Harcourt, Port Harcourt, Nigeria.

<sup>2</sup>Regional Center for Biotechnology and Bio-Resources Research, University of Port Harcourt, Port Harcourt, Nigeria.

### Authors' contributions

This work was carried out in collaboration among all authors. Authors AEA, NGO, PMTU and CJO conceived and designed the research. Material preparation and data collection were performed by authors NGO, PMTU and CJO. Data analysis was performed by authors NGO and PMTU. The first draft of manuscript was written by authors PMTU, NGO and CJO. All authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

### Article Information

DOI: 10.9734/JABB/2021/v24i930239

Editor(s):

- (1) Dr. Mohamed AliAbdel-Rahman, Al-Azhar University, Egypt.  
(2) Prof. Joana Chiang, China Medical University, Taiwan.

Reviewers:

- (1) Suraiya Binte Mostafiz, University Technology Malaysia, Malaysia.  
(2) Maduka, Ndukwu, Wellspring University, Nigeria.

Complete Peer review History, details of the editor(s), Reviewers and additional Reviewers are available here:  
<https://www.sdiarticle5.com/review-history/76248>

Received 11 September 2021

Accepted 22 November 2021

Published 04 December 2021

**Original Research Article**

### ABSTRACT

**Aims:** A study was conducted to identify fungal species isolated from dumpsite soil in University of Port Harcourt using molecular techniques.

**Methodology:** Molecular methods for determining the species of a fungus based on the amplification and sequencing of the internal transcribed spacer (ITS) region of the fungal rRNA operon using molecular markers was applied. Soil sample was collected from a dumpsite in the University of Port Harcourt, Rivers State, Nigeria. Isolation of fungi associated with the dumpsite soil was carried out using spread plate method. Fungal genomic DNA was extracted using Quick-DNA Fungal/Bacterial Miniprep kit. The ITS1-2 gene of the isolates was amplified by Polymerase Chain Reaction (PCR) using the primer pair; ITS4 and ITS5.

**Results:** The sequences of the amplified ITS region were blasted against known sequences on the National Centre for Biotechnology Information (NCBI) database. Nucleotide sequence analysis revealed the species identity of the fungal isolates to be: *Aspergillus fumigatus*, *Trichoderma*

\*Corresponding author: E-mail: nkechi.iyanyi@gmail.com;

*harzianum*, *Aspergillus felis*, *Aspergillus templicola*, *Aspergillus flavipes*, *Aspergillus fumigatus* and *Cunninghamella binariae*. Phylogenetic analysis was carried out to ascertain the relationship between the isolates and other closely-related isolates on GenBank. Isolates 2 (*Trichoderma harzianum*) and 7 (*Cunninghamella binariae*), 3 (*Aspergillus felis*) and 6 (*Aspergillus fumigatus*), and 4 (*Aspergillus templicola*) and 5 (*Aspergillus flavipes*) were found to be more closely related to each other.

**Conclusion:** The molecular techniques employed successfully identified the organisms to the species level as these techniques are based on the genetic constitution of organisms. The result obtained from this study will complement the information on the fungal organisms associated with dumpsite soil.

**Keywords:** Dumpsite; fungi; phylogeny; sequencing; soil.

## 1. INTRODUCTION

Soils are naturally occurring physical covering of the earth's surface and represent the interface of three material states: solids (geological and dead biological materials), liquid (water) and gases. There are four major categories of soil components: organic, minerals, liquid and gases; and there are three properties associated with the soil: chemical (eg: pH), physical (eg: texture) and biological (eg: microbial accumulation) [1]. Soils are the foundation of all terrestrial ecosystems and are home to a vast diversity of bacteria, archaea, fungi, insects, annelids, and other invertebrates, as well as plants and algae. These soil dwellers provide food or nutrients that support organisms that live above and below ground.

A dumpsite is a piece of land used to dispose solid wastes [2]. Dumpsites are mostly without environmental controls as individuals even dump wastes beyond the land carved out for waste disposal. In some cases, solid wastes cover some parts of major roads at some points as a consequence of this. The siting of open dumps around residential areas is associated with various environmental health hazards such as breathing difficulties, malaria etc [3]. Dumpsites act as inoculums of pathogens to the soil, which can lead to the emergence of infections [4]. Soil-borne pathogens such as *Fusarium* spp., *Rhizoctonia* spp., *Verticillium* spp., *Sclerotinia* spp., *Ralstonia* spp., *Pectobacterium* spp., *Agrobacterium* spp. and *Streptomyces* spp. cause diseases in plants with fungal pathogens being the most virulent, chiefly responsible for root rots, wilts, blight and other plant diseases that are usually difficult to control with conventional methods [5].

Identification of fungal species using morphological and microscopic techniques is

laborious and prone to errors [6]. Molecular characterization is based on the genetic composition of organisms which reveals more information than the traditional cultural technique which is based on the morphological and microscopic characteristics of organisms. The use of Polymerase Chain Reaction (PCR) to amplify the Internal Transcribed Spacer (ITS) region of fungi and subsequently sequence the amplified region provides adequate information for identification of fungal species. The use of molecular methods of identification of microorganisms can lead to successful identification of novel species and should be employed in the study of microbial diversities [7]. Characterization of the microbial population of soil is essential for the understanding of biodiversity of soil ecosystem and for the development of control strategies against pathogenic organisms. Most of the authors that have worked on the microorganisms associated with dumpsite soils in Nigeria have used the traditional cultural techniques as the basis of identification. This study was aimed at isolating and identifying the fungi associated with dumpsite soil using both traditional cultural techniques and basic molecular techniques in order to determine the most effective technique that will be able to classify the organisms to the species level. The cultural technique was used to isolate the organisms while the molecular method was basically used for the identification of the species obtained from the study.

## 2. MATERIALS AND METHODS

### 2.1 Study Area and Sampling

The study was carried out at the Regional Center for Biotechnology and Bio-resources Research, University of Port Harcourt, Rivers State, Nigeria. Sequencing of the PCR products was done at the International Institute of Tropical Agriculture

(IITA), Ibadan. The dumpsite is an open dumpsite situated in University of Port Harcourt and consisted of disposed papers, bottles, food remnants, plastics, tin cans, grasses, polyethene bags, leathers, faeces and urine. Soil samples were obtained from five different points at a distance of 6 feet apart. Samples were collected under the refuse pile at 0-15cm soil depth, using an auger borer. Fifteen grammes of soil was collected. The soil was transferred into a sterile zip bag and taken to the laboratory for fungal isolation.

## 2.2 Isolation of Fungi from Dumpsite Soil

Isolation of fungi from the dumpsite soil was carried out using the serial dilution method. One gramme of dumpsite soil was transferred into a test tube containing 10 ml of sterile physiological saline. The soil was mixed well with the solution and 1ml was aseptically transferred to another test tube containing 9 ml of physiological saline. This was repeated for three times and 0.1ml aliquot from the 1,000x dilution was aseptically transferred onto the surface of freshly prepared Potato Dextrose Agar (PDA). A sterile glass rod was used to spread the inoculums all over the surface of the PDA for even distribution of microorganisms. Ten replicates were used and the inoculated plates were incubated at room temperature ( $25^{\circ}\text{C} \pm 2^{\circ}\text{C}$ ) for 7 days. Individual colonies distinguished by their discrete colours were picked from the plates using sterile forceps and sub cultured on freshly prepared PDA plates to obtain pure cultures.

## 2.3 Fungal DNA Extraction, Gene Amplification and Sequencing

Genomic DNA was extracted following the protocol of Quick-DNA™ Fungal / Bacterial MiniPrep Kit (Zymo Research Group, California, U.S.A.) with slight modifications. The mycelium from each fungal isolate was scraped off from a 7-day fully grown culture using a sterile surgical blade and transferred into a sterilized mortar. Liquid nitrogen (-196 °C) was added to the isolate in the mortar in order to freeze the sample for easy homogenization. After the sample was completely frozen, 750 µl of bashing bead buffer (composed of Tris-hydrogen chloride, sodium chloride, sodium deoxychlorate and sodium dodecyl sulfate) was added, and the sample was immediately homogenized using pestle. The mixture was transferred into a 1.5 ml microcentrifuge tube, 200 µl of nuclease free water was added into the tube and centrifuged at

10,000 x g for 1 minute. The supernatant was discarded and the pellet was subsequently used for cell lysis, precipitation and DNA purification, following the manufacturer's protocol. Isolated DNA was stored at 4°C for further analysis.

DNA quantity and purity were measured using Nano Drop 2000c Spectrophotometer (Thermo Fisher Scientific Inc. Wilmington, Delaware, U.S.A). DNA purity was determined as the ratio of UV absorbance by DNA at 260 nm to absorbance at 280 nm. DNA quality was determined by gel electrophoresis on a 1% agarose gel.

The fungal ITS1-2 gene sequences were amplified using the primer pair, ITS4-TCCTCCGCTTATTGATATGS (forward) and ITS5- GGAAGTAAAAGTCGTAACAAGG (Reverse) through Polymerase Chain Reaction (PCR). PCR cocktail contained 3 µL of genomic DNA (10 ng/µL), 0.1 µL of 5 µg/µL Taq DNA polymerase, 2.5 µL of 10x PCR buffer, 1.0 µL of DMSO, 0.8 µL of 2.5 mM DNTPs, 1.0 µL of 25 mM MgCl<sub>2</sub> (Promega), and 1 µL each of forward primer and reverse primer (concentration of 5µM). The total reaction volume was made up to 25 µl using 13.4 µl Nuclease free water. Amplifications were performed in a thermal cycler (Eppendorf) using an initial denaturation step of 94°C for 5 minutes, followed by 36 cycles of denaturation at 94°C for 30 seconds, annealing at 54°C for 30 seconds and elongation for 45 seconds at 72°C with a final extension for 7 min at 72°C. The amplicons (PCR products) from the above reaction were separated by gel electrophoresis on 1.5% agarose gel.


Amplified products were sequenced on ABI 3500 Genetic Analyzer (Thermo Fisher Scientific, Massachusetts, United States of America) using Sanger sequencing. Single-stranded DNA template, a DNA polymerase, a DNA primer, dideoxynucleotide triphosphates (ddNTPs) and deoxynucleotide triphosphates (dNTPs) were used for sequencing. Each dideoxynucleotide (ddNTP) incorporated intermittently terminated DNA chain elongation as they lack 3'-OH group which is responsible for the formation of a phosphodiester bond between two nucleotides. Consequently, extension of the DNA stopped.

## 2.4 Statistical Analysis

Frequency of occurrence of fungi were determined based on the Score Method recommended by Ataga and Akueshi [8]; where

the numeral “1” represents absence of fungi and “2” represents presence of fungi; so that any value above “1” indicates presence of fungi. The identification of the isolated fungi by traditional cultural method was carried out with reference to Umechuruba and Elenwo [9] and Ataga et al.

[10]. For identification using molecular technique, sequence editing which comprises trimming and sequence alignment using Clustal W were performed on MEGA X. Sequences were blasted on NCBI database for identification of species.



**Fig. 1. Map of University of Port Harcourt**  
Source (Larinde and Oladele, 2014)



**Fig. 2. Dumpsite studied in University of Port Harcourt**

## 2.5 Phylogeny

The sequences of the ITS1-2 gene of the isolates were compared with the ITS sequences in GenBank. Best BLAST hits were used for the construction of neighbor-joining phylogenetic tree using the maximum composite likelihood method and evolutionary analysis was conducted on MEGA X software [11].

## 3. RESULTS

### 3.1 Fungal Isolates Obtained from Dumpsite Soil

A total of seven (7) fungi were isolated from the dumpsite soil. The pure cultures of the fungi are presented in Fig. 1. The frequency of occurrence of each isolate was determined. Isolates 2 and 7 had the highest frequency of occurrence (1.8), followed by isolate 1 (1.7), isolates 3 and 4 (1.6), isolate 5 (1.5), and isolate 6 (1.4). Table 1 shows the morphological description of the isolates after seven days on potato dextrose agar medium with their frequencies of occurrence. The Zymo Fungal / Bacteria Miniprep kit effectively


extracted the fungal DNA with 260/280 nm ratio (DNA purity) between 1.28 and 1.97.

### 3.2 Amplified Products from ITS1-2 Gene

Extracted genomic DNA from the seven fungal isolates were successfully amplified using the ITS4 and ITS5 primer pair. Amplified DNA showed bands on gel when viewed under UV light. The fragment produced by each species on 1.5 % agarose gel was compared with the different fragments of 1Kb DNA ladder to obtain the size of the amplicons as presented in Fig. 2. The expected amplicon size for ITS gene was obtained.

### 3.3 DNA Sequences and Blasting

The numbers of base pairs of the sequenced ITS1-2 region of the fungal isolates are: 583bp, 622 bp, 681 bp, 728 bp, 588 bp, 643 bp and 665 bp for samples 1 to 7 respectively. The sequence alignment of each isolate was more than 200base pairs which is ideal for the identification of microorganisms.




**Fig. 3. Fungal isolates obtained from University of Port Harcourt waste dumpsite soil and cultured on PDA**

Note: DS-1 to DS-7 represent isolate ID 1 to 7.DS-1: *Aspergillus* sp., DS-2: *Trichoderma* sp., DS-3: *Aspergillus* sp., DS-4: *Aspergillus* sp., DS-5: *Aspergillus* sp., DS-6: *Aspergillus* sp., DS-7: *Cunninghamella* sp

**Table 1. Fungi isolated from dumpsite soil at room temperature ( $25^{\circ}\text{C} \pm 2^{\circ}\text{C}$ )**

Isolate ID	Morphology	Frequency of Occurrence	Suspected Organism
DS-1	Rough, dark green mycelia	1.7	<i>Aspergillus</i> sp.
DS-2	Smooth, dark green mycelia	1.8	<i>Trichoderma</i> sp.
DS-3	Smooth, dark and hard mycelia	1.6	<i>Aspergillus</i> sp.
DS-4	Smooth, white, cotton-like mycelia	1.6	<i>Aspergillus</i> sp.
DS-5	Hard, white mycelia with yellow spores in the middle	1.5	<i>Aspergillus</i> sp.
DS-6	Smooth, hard dark green mycelia	1.4	<i>Aspergillus</i> sp.
DS-7	Grey hairy mycelia with dark base	1.8	<i>Cunninghamella</i> sp.

**Fig. 4. Gel electrophoresis profile of amplified ITS1-2 gene sequences of the fungal DNA from University of Port Harcourt waste dumpsite soil**

Note: L – 1 kb DNA ladder; Lane 1 to 7- Amplicons from fungal isolates

BLAST results revealed the species identity of the fungal isolates to be: DS 1- *Aspergillus fumigatus*; DS 2- *Trichoderma harzianum*; DS 3- *Aspergillus felis*; DS 4- *Aspergillus templicola*; DS 5- *Aspergillus flavipes*; DS 6- *Aspergillus fumigatus*; DS 7- *Cunninghamella binariae*. The taxonomic affinities of the ITS1-2 gene sequences with the most closely related sequences on GenBank are shown in Table 2. Each sequence of the fungal isolates was given a strain number, the sequences were submitted to the GenBank and accession numbers were assigned to each isolate as shown in Table 3.

### 3.4 Phylogenetic Analysis

The sequence analysis showed genetic diversity among the isolates with five isolates belonging to the genus *Aspergillus*. The phylogenetic tree constructed revealed the organisms (on GenBank) that are most closely related to the


isolates obtained from the dumpsite soil, as presented in Fig. 3. The vertical lines on the trees indicate the difference between the branches. The greater the length of the vertical line, the more the difference between the branches. The isolates were found to be closely-related to *Aspergillus fumigatus*, *Trichoderma harzianum*, *Aspergillus felis*, *Aspergillus templicola*, *Aspergillus flavipes*, *Aspergillus fumigatus* and *Cunninghamella binariae*. These organisms are in turn closely related to *Curvularia affinis*, *Aspergillus micronesiensis*, *Aspergillus parafelis*, *Botryosphaeria obtusa*, *Gongronella guangdongensis* and *Trichoderma aureoviride*. Isolates 2 (*Trichoderma harzianum*) and 7 (*Cunninghamella binariae*), 3 (*Aspergillus felis*) and 6 (*Aspergillus fumigatus*), and 4 (*Aspergillus templicola*) and 5 (*Aspergillus flavipes*) were found to be more closely related to each other.

**Table 2. Putative Taxonomic affinities of sequence types inferred from BLAST search of ITS sequences**

<b>Sample ID</b>	<b>Taxonomic affinity (Gene bank no)</b>	<b>Percentage Similarity</b>
DS-1	<i>Aspergillus fumigatus</i> (MK732091.1)	90.04
DS-2	<i>Trichoderma harzianum</i> (KR868343.1)	100
DS-3	<i>Aspergillus felis</i> (MH865509.1)	84.56
DS-4	<i>Aspergillus templicola</i> (KP987081.1)	99.20
DS-5	<i>Aspergillus flavipes</i> (MH865240.1)	99.81
DS-6	<i>Aspergillus fumigatus</i> (MH141301.1)	90.17
DS-7	<i>Cunninghamella binariae</i> (MH858865.1)	99.00

**Table 3. Accession and strain numbers of fungal isolates**

<b>Sample</b>	<b>Accession number</b>	<b>Strain number</b>
DS-1	MN602899	RCBBR_AEA1
DS-2	MN602970	RCBBR_AEA2
DS-3	MN603056	RCBBR_AEA3
DS-4	MN602898	RCBBR_AEA4
DS-5	MN602941	RCBBR_AEA5
DS-6	MN603087	RCBBR_AEA6
DS-7	MN603006	RCBBR_AEA7



**Fig. 5. Phylogenetic tree showing the relationship between the fungal isolates obtained from University of Port Harcourt waste dumpsite and other fungal species on GenBan**

#### 4. DISCUSSION

Dumpsite accommodates numerous and diverse microorganisms, some of which are pathogenic to other environmental inhabitants such as plants, animals and humans. In this study, isolates 2 and 7 had the highest frequency of occurrence (1.8), followed by isolate 1 (1.7), isolates 3 and 4 (1.6), isolate 5 (1.5), and isolate 6 (1.4).

PCR amplification of the ITS1-2 gene sequences of the fungal isolates yielded amplicons between 580 to 800 base pairs. Internal Transcribed Spacer (ITS) region has been reported for its importance in biodiversity study and is proven to be more reliable in distinguishing fungal species [12] than traditional cultural methods. The molecular techniques used in this study proved successful as they led to the identification of seven fungal species viz: *Aspergillus fumigatus*, *Trichoderma harzianum*, *Aspergillus felis*, *Aspergillus templicola*, *Aspergillus flavipes*, *Aspergillus fumigatus* and *Cunninghamella binariae*.

The genus *Aspergillus* belongs to the phylum Ascomycota, class Euromycetes, order Eurotiales and family Trichomaceae. *Aspergillus fumigatus* has been isolated from milled rice [13] and semi-desert soils [14]. It is an opportunistic filamentous fungus, one of the most common *Aspergillus* species that cause diseases in individuals with immunodeficiency which include patients receiving immunosuppressive therapy for autoimmune or neoplastic disease, organ transplant recipients, and AIDS patients [15]. *A. fumigatus* is a saprotroph, widespread in nature and is typically found in soil and decaying organic matter such as compost and heaps, where it plays an essential role in carbon and nitrogen recycling [16]. Colonies of the fungus produce thousands of minute grey-green conidia (2-3 µm) from their conidiophores and these conidia readily become airborne. *A. fumigatus* is capable of growth at 37°C and can grow at temperatures up to 50°C, with conidia surviving at 70 °C, the typical temperature of self-heating compost heaps. Its spores are ubiquitous in the atmosphere.

*Aspergillus felis* is a thermo-tolerant fungus, with optimum growing temperatures of ≥45°C and <20°C. Species of the genus *Aspergillus* can cause aspergillosis, in a diverse range of human and animal hosts. *A. felis* can be easily misidentified as the closely related fungus *A.*

*fumigatus*, which is a well-studied cause of disease in humans. However, *A. felis* is intrinsically more resistant to anti-fungal drugs than *A. fumigatus* [17].

*Aspergillus templicola* belongs to the section *Flavipedes*. Members *Flavipedes* are found in soils, rhizospheres, caves and indoor environments usually as endophytes in plants, contaminants in food and as human pathogens [18]. They have been isolated from hypersaline soils [19]. *Aspergillus flavipes* is another fungus belonging to the *Flavipedes* section in the genus *Aspergillus* and is one of the most common and widely distributed species of this section. The species was first described in 2014 [20]. *A. flavipes* has been reported to produce extrolite, a wide array of bioactive compounds such as sterigmatocystin, citrinin, and lovastatin [16]. *A. flavipes* is also an endophytic fungi capable of producing secondary metabolites such as flavipesin A, flavipesin B, guignardic acid, and dibutyl phthalate that exert antibiosis against plant and human pathogens [21].

*Trichoderma harzianum*, a filamentous soil fungus, is an effective bio-control agent of several economic important soil-borne plant pathogens and viral vectors [22]. They are present in nearly all soils, where they occur as opportunistic plant symbionts and antagonists to many phytopathogenic fungi, thereby protecting plants from diseases. They are used for foliar application, seed treatment and soil treatment for suppression of various disease-causing fungal pathogens. *Trichoderma* strains are naturally resistant to many toxic compounds including herbicides, fungicides and pesticides such as phenolic compounds.

*Cunninghamella binariae* is a filamentous fungus found in soil and plant materials, particularly at Mediterranean and subtropical zones. It has also been isolated from animal feces [23]. *Cunninghamella* is a genus in the order Mucorales covering species with distinctive morphological features. In addition to being common contaminants, *Cunninghamella* species cause infections such as Mucormycosis in immune-compromised patients with high mortality [24].

The works of other researchers, who employed the traditional cultural method of characterization, revealed similar fungi species with the ones isolated in this study, with species of the *Aspergillus* genus being predominant in the

different waste dumpsites studied. Simon-Oke and Alade [25] carried out a study on two dumpsites in Akure metropolis in Nigeria and isolated *Aspergillus flavus*, *A. fumigatus*, *A. niger* and *Mucor mucedo* among other fungal and bacterial species. The authors suggested that the low bacteria and fungi count in one of the dumpsites may be as a result of smaller size of the dumpsite and the low human activities in the area. Eleanya et al. [26] reported the isolation of *Aspergillus niger*, *Fusarium* spp., *Rhizopus* spp. and *Mucor* spp. from dumpsites in Bwari, Federal Capital Territory, Nigeria. Williams and Hakam [27] isolated *Aspergillus*, *Fusarium*, *Mucor*, *Penicillium* and *Saccharomyces* species from dumpsites in Port Harcourt. While the traditional cultural method they employed revealed similar fungi species, the molecular method of characterization employed in this study revealed that more fungi species such as, *Aspergillus flavipes*, *Aspergillus templicola*, *Trichoderma harzianum* and *Cunninghamella binariae* can also be found in dumpsite soil. Microorganisms tend to be more abundant in the soil when there is increased moisture for instance, during the rainy season [25].

The molecular method used in this study effectively characterized all the fungi isolates at the species level. This study has outlined some of the fungal organisms associated with dumpsite soil and the economic importance of the isolates have also been highlighted to give readers the insight into the beneficial or detrimental effects of the isolates. Some of the organisms obtained from this study are plant pathogens while others are of public health concern.

## 5. CONCLUSION

Five species of *Aspergillus* and one specie each of *Cunninghamella* and *Trichoderma* were obtained from this study. Though most of the isolates obtained from this study belong to the genus *Aspergillus*, other fungi genera isolated (*Trichoderma* and *Cunninghamella*) are equally of economic importance. Dumpsites are mapped out for waste disposal, so the soils are usually contaminated with certain microorganisms some of which may be pathogenic or beneficial to man. This study has shown that dumpsite soil holds diverse fungal species of public health significance. The results obtained from the current research work will complement the existing knowledge on the fungal organisms associated with dumpsite soil and also educate individuals on the dangers of living close to

dumpsites due to the adverse effects of health implications that can be caused by some of these dumpsite fungi.

## DISCLAIMER

The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Dominati E, Patterson M and Mackay A. A framework for classifying and quantifying natural capital and ecosystem services of soils. *Ecol Econ.* 2010;69: 1958-1968.
2. Tse AC, Adamu CI. Assessment of anthropogenic influence on quality of groundwater in hand-dug wells in parts of Makurdi metropolis, North central Nigeria. *I JS.* 2012;14(1):123-135.
3. Aluko OO, Obafemi TH, Obiajunwa PO, Obiajunwa CJ, Obisanya OA, Odanye OH, Odeleye AO. Solid waste management and health hazards associated with residence around open dumpsites in heterogeneous urban settlements in Southwest Nigeria. *Int. J. Environ. Health Res.* 2021;1-16.  
DOI: 10.1080/09603123.2021.1879738
4. Nyandjou JM, Yakubu SE, Abdullahi IO, Machido DA. Enteric bacteria of public health significance isolated from Zaria metropolis dumpsite soil. *Sci World J.* 2018;13(4):30-34.
5. Baysal-Gurel F, Gardener BM and Miller SA. Soil borne diseases management in organic vegetable production. *eOrganic;* 2012.

Accessed 14 October 2021.

- Available:<https://articles.extension.org/pages/64951/soilborne-diseases-management-in-organic-vegetable-production>
6. Adeniyi M, Titilawo Y, Oluduro A. Molecular identification of wild Nigerian mushrooms using internal transcribed spacer: Polymerase chain reaction. *AMB Express.* 2018;8:148-156.
  7. Elijah AI, Atanda OO, Popoola AR, Uzochukwu SV. Molecular characterization and potential of bacterial species associated with cassava waste. *Niger Food J.* 2014;32(2):56-65.
  8. Ataga AE, Akueshi CO. Fungi associated with sunflower seeds in Nigeria. *Seed Res.* 1986;1(24):64-65.
  9. Umechuruba CI and Elenwo EN. Diagnostic techniques in mycology with practical exercise. Belk Publishers. 1997;30- 42.
  10. Ataga AE, Elenwo EN and Nwachukwu EO. Laboratory exercises and series in mycology. ACOTEC Technologies, Port Harcourt, Nigeria. 2010;58-64.
  11. Kumar S, Stecher G, Li M, Knyaz C and Tamura K. MEGA X: Molecular evolutionary genetics analysis across computing platforms. *Mol Biol Evol.* 2018;35:1547-1549.
  12. Schoch CL, Seifert KA, Huhndorf S, Robert V, Spouge JL, Levesque CA, et al. Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for fungi. *PNAS USA.* 2012;09(16):1-6.
  13. Yazdani D, Zainal-Abidin MA, Tan YH, Kamaruzaman S. Molecular identification of *Aspergillus* and *Eurotium* species isolated from rice and their toxin-producing ability. *Microbiology* 2011;80(50):720-727.
  14. Giusiano GE, Piontelli E, Fernández MS, Mangiaterra ML, Cattana ME, Kocsblé S, et al. Biodiversity of species of *Aspergillus* section fumigatiin semi-desert soils in Argentina. *Rev Argent Microbiol.* 2017;49(3):247-254.
  15. Ben-Ami R, Lewis RE, Kontoyiannis DP. Enemy of the immunosuppressed state: an update on the pathogenesis of *Aspergillus fumigatus* infection. *Br J Haematol.* 2010;150(4):406-417.
  16. Fang W, Latge JP. Microbe profile: *Aspergillus fumigatus*: A saprophytic fungi and opportunistic fungal pathogen. *Microbiology.* 2018;164(8):1009-1011.
  17. Barrs VR, Van-Doorn TM, Houbraken J, Kidd SE, Martin P, Pinheiro MD, et al. *Aspergillus felis* sp. nov., an emerging agent of invasive aspergillosis in humans, cats and dogs. *PLoS One* 2013;8(6):1-11.
  18. Hubka V, Nováková A, Kolarik M, Jurjevic Z, Peterson SW. Revision of *Aspergillus* section Flavipedes: seven new species and proposal of section Jani sect. nov. *Mycologia* 2015;107(1):169-208.
  19. Arzanlou M, Samadi R, Frisvad JC, Houbraken J, Ghosta Y. Two novel *Aspergillus* species from hypersaline soils of the National Park of Lake Urmia, Iran. *Mycol Prog.* 2016;15(10-11):1081-1092.
  20. Visagie CM, Hirook, Y, Tanney T, Whitfield E, Mwange K, Meijer M, et al. *Aspergillus*, *Penicillium* and *Talaromyces* isolated from house dust samples collected around the world. *Stud Mycol.* 2014;78:63-139.
  21. Zhang H, Tang Y, Ruan C, Bai X. Bioactive secondary metabolites from the endophytic *Aspergillus* genus. *Rec Nat Prod.* 2016;10(1):1-16.
  22. Brunner K, Zeilinger S, Ciliento R, Woo SL, Lorito M, Kubicek CP, Malc RL. Improvement of the fungal biocontrol agent *Trichoderma atroviridae* to enhance both antagonism and induction of plant systemic disease resistance. *Appl Environ Microbiol.* 2005;71:3959-3965.
  23. Nguyen TT, Hyde KD, Lee HB. *Cunninghamella binariae*, *Mucor ardhlaengikitus*, *Mucor gigasporus* and *Umbelopsis changbaiensis*, Newly recorded species from amphibian feces and soil in Korea. *Phytotaxa.* 2019;425:19-34.
  24. Yu J, Walther G, Van-Diepeningen A, Van-Den-Ende A, Li RY, Moussa T, et al. DNA barcoding of clinically relevant *Cunninghamella* species. *Med Mycol J.* 2014;53(2):99-106.
  25. Simon-Oke IA and Alade T. Survey andidentification of bacteria and fungi associated with two dumpsites in Akure metropolis. *Microbiol Res Int.* 2018;24(5):1-8.

26. Eleanya EU, Oyegue TO, Rwang PG, Ozoude TO, Okey-Ndeche NF. Microbiological analysis of dumpsites in Bwari, Federal Capital Territory, Nigeria. *Microbiol Res Int.* 2017;22(3):1-6.
27. Williams JO, Hakam K. Microorganisms associated with dumpsite in Port Harcourt Metropolis, Nigeria. *J Ecol Nat Environ.* 2016;8(2):9-12.

© 2021 Ogbuji et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/76248>*