



Non-commutativity Over Canonical Suspension η for Genus $g \geq 1$ in Hypercomplex Structures for Potential ρ_ϕ

Deep Bhattacharjee ^{a,b*}

^a Theoretical Physics Research Division of AATWRI Aerospace and Defense Research Directorate, India.

^b Electro Gravitational Space Propulsion Laboratory, India.

Author's contribution

The sole author designed, analyzed, interpreted and prepared the manuscript.

Article Information

DOI: 10.9734/ARJOM/2022/v18i11607

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/92716>

Original Research Article

Received 09 August 2022

Accepted 19 October 2022

Published 25 October 2022

Abstract

Any matrix multiplication is non-commutative which has been shown here in terms of suspension n^\ddagger , annihilator, and factor as established over a ring following the parameter k over a set of elements upto n for an operator to map the ring R to its opposite R^{op} having been through a continuous representation of permutation upto $n -$ cycles being satisfied for a factor f along with its inverse f^{-1} over a denoted orbit γ on $k -$ parameterized ring justified via suspension $\eta \in \eta^0, \eta^1$ implying the same global non-commutativity for the annihilator \mathcal{A} . This will be used for the construction of the genus-alteration scenario where the suspension η^0 acting with its opponent η^1 on any topological space \mathcal{J} can alter the geometry making a change in the manifolds for taking over the Boolean $(1,0)$ satisfying the concerned operations.

Keywords: Operators; non-commutativity; annihilator; boolean.

1 Introduction

As mentioned in paper [1] the structural suspension for functor η over the canonical stabilizer points $2^2\eta$ for genus $g \geq 1$, no such higher dimensional generalizations have been considered or computed. Here, in this paper, I have taken the same suspension functor η which in a way can alter the genus g depending on the commutativity or non - commutativity as depicted over map φ^x . The non-trivial question that can arise is ‘why should one considers the commutativity and non-commutativity when it comes to suspension functor η in genus $g \geq 1$, scenario?’[2-5]. Taking the affine parameter ∂ where $\partial = \coprod_{\omega=\infty}(\eta^0, \eta^1)^\omega / \sim$ (not to be confused with

*Corresponding author: Email: deep.bh@outlook.com;

the parameter of the paper in Ref. 5) over a value of $\omega = \infty$ for an infinite hypercomplex manifold existing in intersection points of grids in space-time geometry, any operator \mathcal{O} when makes a proper orbit γ that is for the n -cycles permutations over factor $\mathfrak{f} \simeq$ effective operations in ring R with its inverse $\mathfrak{f}^{-1} \simeq$ effective operations in opposite ring R^{op} for $1,2,3,4 \ni x$ over map φ^x , the orbit $\gamma = \sum_{x=1}^4 \varphi^x \cong \prod k$ [as explained in details in Sec. II] there exists commutativity or no-suspension if the mapping through n -cycles with the denoted orbit makes a trajectory from the domain to the codomain back to the domain or otherwise back to the point of initiation then the orbit γ will be closed; there will always be a false Boolean (0) denotes no change or alteration in the geometries of the hypercomplex structures, but if the point of initiation and point where the orbit ends is not the same then the orbit will remain open implying a non-commutativity over a true Boolean (1) denotes the suspension as the gap – difference all computed for orbit $\gamma \cong \prod k$ over ring \oplus^k is sufficient to alter the genus geometries via suspension through the functor $\eta^1 \forall \eta \in \eta^0, \eta^1$ where the gap ρ denotes the non-commutativity to suffice in the following relations [6-9],

gap ρ is non-existent iff $domain \rightarrow \dots \rightarrow codomain \rightarrow \dots \rightarrow domain$ establishes for $1,2 \ni x$ over map φ^x
 gap ρ is existent iff $domain \rightarrow \dots \rightarrow codomain \rightarrow \dots \rightarrow domain \rightarrow \dots \rightarrow codomain$ establishes for $3,4 \ni x$ over map φ^x

Establishing the properties,

$$\left\{ \begin{array}{l} n^{th} \text{ order cycles in } domain \rightarrow \dots \rightarrow codomain \\ \text{Makes suspension } \eta \\ \text{For difference or open orbits of } \gamma \end{array} \right.$$

$$\left\{ \begin{array}{l} codomain - domain \simeq gap \rho \Rightarrow open \text{ orbits} \equiv non - commutativity \\ \text{Over commutative / non - commutative parameter } \mathcal{C} [\text{appearing in Sec. IV}] \\ \text{Over commutative / non - commutative parameter} \end{array} \right.$$

For construction, I will take a 4-tuple relation to justify the suspension provided all other necessary relations will be there co-existing beside these 4-parameters. This 4-tuple can be represented through [10-12],

$$(\tau(\nabla^y), \iota^\lambda, /_{\rho\lambda}, \chi^\lambda) \text{ such that } \forall \lambda = 1 \text{ or } 0 \text{ where } (0,1) \text{ is Boolean and } y \text{ in } \nabla^y \text{ takes values } 1 \text{ or } -1 [\text{as described in Sec.IV}]$$

Representing each parameter with a 0 or 1 except ∇^y taken over the variable λ where there exist 4 relations,

- A. *Tension* $\tau(\nabla^y)$ – This should be considered as the tension τ acting on ∇^y with y either 1 or -1 depending upon the orientation [as discussed in detail in Sec.IV]. This tension would build up for open orbit when operator \mathcal{O} won't commute because of the value of x as 3 or 4 for map φ^x [as described in Sec.III].
- B. *Trajectory* ι^λ – This gives the value of the orbit γ in terms of the trajectory ι with the value as appeared in λ in ι^λ gives 1 for open orbit when operator \mathcal{O} won't commute because of the value of x as 3 or 4 for map φ^x [as described in Sec.III] or gives 0 for closed orbit when operator \mathcal{O} commute because of the value of x as 1 or 2 for map φ^x [as described in Sec.III]
- C. *Loads* $/_{\rho\lambda}$ – This provides the impact on the gap ρ when the orbit is open for the load to exist as 1 representing the variable λ making it prominent for deformation to be discussed in [Point D] else its 0 as appeared in case of close orbit in $/_{\rho\lambda}$ with / denotes the closure.
- D. *Deformations* χ^λ – This relates [Point A, B, C] for the deformation in structure which is affected by the parameter χ with the deformation occurs only in case of open orbit for cumulative tension $\oint \tau(\nabla^y)$ with this in effect only when λ in χ^λ takes up 1 else in case of close orbit the deformation won't exist for $\lambda = 0$ in χ^λ .

2 Constructions

Suspension over a topological space \mathcal{J} can induce its inert geometry making it bend or create genus ≥ 1 which imparts a relation that the same space can be transformed into an algebraic category if we make a grid over that space and in each intersection of the aforesaid grid, the manifolds that are existent in hypercomplex form can be

changed from their one structure to the other resulting a change in the number of genera thereby inducing a different category of alterations. But this can only suffice if the suspension functor η^0 can be grouped with another suspension functor that opposes its own kind η^1 as while η^1 creates genus, η^0 removes or destroys the genus provided there exists two relations [1,10,11,13],

1. Two hypercomplex for concerned functors $\eta^1, \eta^0 \exists \eta \in \eta^0, \eta^1$ are non-commutative as appeared in annihilator \mathcal{A} .
2. There exists the relation of equivalence closure for the affine parameter ∂ through ring $R = \oplus^k$ where $\partial = \coprod_{\omega=\infty} (\eta^0, \eta^1)^\omega / \sim$ over a value of $\omega = \infty$ for an infinite hypercomplex manifold present in topological space \mathcal{J} but to each one of them the operation is closed for the parameter ∂ .

In space-time grid there are multiple compact Kähler with Ricci flatness or vanishing Ricci curvature with different supersymmetric phenomenology related structure that they denote. Considering the genus geometry, for a particular hypercomplex structure if the operation of suspension in Boolean over η^1, η^0 satisfies then for the taken parameterization k of ring R being defined over an affine parameter ∂ through that same ring with also its opposite denoting the alteration operation that can spread over multiple hypercomplex structures, one needs to be careful that the operation is closed via equivalence on one hypercomplex structures else the effective geometric suspension of both types of Boolean (0,1) where to categorize properly this can be considered upon as a viable representation [4, 5,14-18],

3. I'll categorize hypercomplex structure \mathcal{H} with the number of genera as its sub, viz, \mathcal{H}_1 as a hypercomplex with 1-genus, \mathcal{H}_2 as a hypercomplex with 2-genus and so on, where this example is trivial just to denote and explain the taken parameters[1,10].
4. If I denote a hypercomplex with 3-genus then I would categorize this from 1 to 3 in the set notation $\mathcal{H}_3 = \{1,2,3\}$ and another one with 4-genus as $\mathcal{H}_4 = \{1,2,3,4\}$ with so on for more genera [9-11].
5. η^0, η^1 will always act on \mathcal{H}_g with g being the number of genus where 0 and 1 in η denotes the suspension or not through the resultant orbit γ being open or closed making the order in a way as to represent the previously said affine parameter closed through ∂ for $\omega = \infty$ for an infinite hypercomplex manifold present in topological space \mathcal{J} [7,8,19,20] that would act in 2 ways via disjoint union among hypercomplex \mathcal{H}_g with the other $\mathcal{H}_g^{\wedge/}$ for \mathcal{H}_g with $g = 4$ represented via $\mathcal{H}_g^{\wedge/} = \{1,2,3,4\}$ and \mathcal{H}_g^{\wedge} with $g = 2$ represented via $\mathcal{H}_g^{\wedge} = \{1,2\}$ with $\wedge = 0$ or 1 and / associated with \mathcal{H} simply denotes the two different hypercomplex without arising any confusion η will act in the geometry alteration of genus g through the resemblance of (trivial just to denote the case) but (non-trivial in respect of operations at Planck's scale) if η_0 acts on $\mathcal{H}_g^{\wedge/}$, i.e., $\mathcal{H}_4^{0/}$ and η_1 acts on \mathcal{H}_g^{\wedge} , i.e., \mathcal{H}_2^1 then this suffice for two such hypercomplex structures where \oplus^k where $\partial = \coprod_{\omega=\infty} (\eta^0, \eta^1)^\omega / \sim$ over a value of $\omega = \infty$ closed for infinite hypercomplex manifold [6,11,13]

$$\mathcal{H}_g^{\wedge/} \coprod \mathcal{H}_g^{\wedge} = \mathcal{H}_4^{0/} \cup \mathcal{H}_2^{1*} = \{(1,0), (2,0), (3,0), (4,0), (1,1), (2,1)\}$$

$$\exists \mathcal{H}_4^{0/} = \{(1,0), (2,0), (3,0), (4,0)\} \text{ and } \mathcal{H}_2^{1*} = \{(1,1), (2,1)\}$$

$\forall \coprod_{\omega=\infty} (\eta^0, \eta^1)^\omega / \sim$ with the annihilator \mathcal{A} representing globally non-commutativity for factors $\mathfrak{f}, \mathfrak{f}^{-1}$ in matrix multiplications [5].

3 Formulations and Dependability

Considering a ring R of k – parameterized where $k = \{1,2,3 \dots n\}$ where each element denotes an orbit γ taking over the ring for an operator \mathcal{O} to move from the ring R to the opposite of ring R^{op} to establish a factor \mathfrak{f} which corresponds to the permutations upto n – cycles for the n^{th} element in set k making 4 – possible relations over a commutative $\mathcal{C} \exists c_+, c_- \in \mathcal{C}$ giving the 5th relation by sufficing 4 –relations to be considered below in stepwise formulations [10,11,13].

Taking the ring R if it's established via a movement or rather a cyclic movement through an operator \mathcal{O} from ring R to its opposite R^{op} then a chain-multiplier can be denoted for the action of the operator \mathcal{O} to denote an orbit γ as stated in 3 – relations below [1,5],

$$\left\{ \begin{array}{l} \text{ring } R \simeq \bigoplus^k \\ \text{operator } \mathcal{O} \rightarrow R \rightarrow R^{op} \\ \text{orbit } \gamma \cong \prod k \end{array} \right.$$

Considering the affine parameter ∂ through ring $R = \bigoplus^k$ where $\partial = \coprod_{\omega=\infty} (\eta^0, \eta^1)^\omega / \sim$ over a value of $\omega = \infty$ for an infinite hypercomplex manifold [For hypercomplex manifolds Ref. [11,13,15-19,21] gives a detailed account] present in topological space \mathcal{J} – The suspension functor η over the set $\eta \in \eta^0, \eta^1$ but to each one of them the operation is closed for the parameter ∂ and the reason for closing the operation is due to the control of suspension [1] from affecting one genus and then spreading to other genera though when in each genus two operations are being carried out for both commutative and non-commutative forms over the same parameter \mathcal{C} to be represented afterward for an infinite number of hypercomplex structures $\omega = \infty$, if one observes closely orbit $\gamma \in \text{proper orbit, suspension orbit}$ then the change in the geometry of the hypercomplex structures can only be effective iff the orbit moves from the domain to co-domain (here the ring R and opposite ring R^{op}) establishes 4 relations over map φ^x [1,6,9,10,13],

$$\left\{ \begin{array}{ll} \varphi^1 : R \rightarrow \dots \rightarrow R & \Rightarrow \text{no suspension as there is commutativity} \\ \varphi^2 : R^{op} \rightarrow \dots \rightarrow R^{op} & \Rightarrow \text{no suspension as there is commutativity} \\ \varphi^3 : R \rightarrow \dots \rightarrow R^{op} & \Rightarrow \text{suspension as there is non - commutativity} \\ \varphi^4 : R^{op} \rightarrow \dots \rightarrow R & \Rightarrow \text{suspension as there is non - commutativity} \end{array} \right. \left. \begin{array}{l} \text{Boolean False}(\eta^0) \\ \text{Boolean False}(\eta^0) \\ \text{Boolean True}(\eta^1) \\ \text{Boolean True}(\eta^1) \end{array} \right|$$

Establishing the orbit taken before $\gamma \cong \prod k$ where γ denotes the formulation [1],

$$\gamma = \sum_{x=1}^4 \varphi^x \cong \prod k$$

4 Structuring Factor with Its Inverse

Here I'll consider 2 specific factor where one is the inverse of the other in generating the n-cycle for consideration upon factor \mathfrak{f} with it's inverse \mathfrak{f}^{-1} where the cycles are the orbits being represented over the integral [22],

$$\nabla^1 = 2\pi i \oint_{\gamma} \mathfrak{f} d\varphi^x$$

$$\nabla^{-1} = -2\pi i \oint_{\gamma} \mathfrak{f}^{-1} d\varphi^x$$

Now, as appeared in [Point A] in [Sec. I] the tension $\tau(\nabla^y)$ should be defined for y in ∇^y provided the representation map of φ^x either takes the value 1 or 2 / or 3 or 4 the trajectory ι^λ needs to be calculated which is the $n - \text{trajectory}$ for cycle – $n \left\{ \begin{array}{l} \simeq R \\ \simeq R^{op} \end{array} \right.$ in either case through the movement from 1 – n or 1 – n in inverse cycles for factor $\mathfrak{f}, \mathfrak{f}^{-1} \supset \iota \vee \nabla^1, \nabla^{-1} \ni \nabla^y$ one can easily find 2 – orbits $\left\{ \begin{array}{l} \text{open} \\ \text{close} \end{array} \right.$ in the order 1 \rightarrow 2 \rightarrow 3 ... \rightarrow

n or $3 \rightarrow 2 \rightarrow 1 \dots n$ for its inverse only and if only map φ^x for the value of $x \begin{cases} 1,2 \\ 3,4 \end{cases} \simeq \eta, \mathcal{C}$ [C appearing below] denoting the relation [1,6-8,10,11,19,20],

$$\left\{ \begin{array}{l} 1,2 \Rightarrow \text{closed} \\ 3,4 \Rightarrow \text{open} \\ \exists \text{all exists in } \begin{cases} R \rightarrow R^{op} \\ R^{op} \rightarrow R \end{cases} \forall n - \text{elements of } k \text{ in } \bigoplus^k \end{array} \right.$$

Thus, the detailed cycles can be shown for the relation $k = \{1,2,3 \dots n\}$ in ring $R = \bigoplus^k$ with opposite R^{op} suffice [10,11,13],

$$\left\{ \begin{array}{l} \nabla^1 \Rightarrow \text{factor } \mathfrak{f} = \begin{pmatrix} 1 & 2 & 3 & \dots & n \\ 2 & 3 & 1 & \dots & n \end{pmatrix} \text{ for cycle - } n \simeq R \\ \nabla^{-1} \Rightarrow \text{factor } \mathfrak{f}^{-1} = \begin{pmatrix} 3 & 2 & 1 & \dots & n \\ 2 & 1 & 3 & \dots & n \end{pmatrix} \text{ for cycle - } n \simeq R^{op} \end{array} \right.$$

Therefore to take into account [Point C and D] viz, loads $/_{\rho\lambda}$ and deformations χ^λ for making a way to establish the non-trivial representation of commutative / non – commutative parameter $\mathcal{C} \in c_+, c_-$ such that $c_+ = \text{factor } \mathfrak{f}$ and $c_- = \text{factor } \mathfrak{f}^{-1}$ one gets a relation [2,5,23],

$$\text{iff } c_+ \times c_- = c_+ \times c_- \Rightarrow R \rightarrow R^{op} \rightarrow R \rightarrow R^{op} \xrightarrow{\sim} R \cong \text{proper orbit } \gamma$$

Else,

$$\text{iff } c_+ \times c_- \neq c_+ \times c_- \Rightarrow R \rightarrow R^{op} \rightarrow R \rightarrow R^{op} \xrightarrow{\sim} R^{op} \cong \text{suspension orbit } \gamma$$

5 Suspensions with Boolean Representation

As mentioned in [Sec II] at the first page in [Point 3] for a proper representation I'll consider a set of two hypercomplex structures as concerned for the alterations of the genera geometry. If two hypercomplex structures with $\wedge = 0$ or 1 in any values of genus g to consider $\mathcal{H}_g^{\wedge/}$ and \mathcal{H}_g^{\wedge} in a set as described over the relation [2,23,24],

$$\star = (\mathcal{H}_g^{\wedge/}, \mathcal{H}_g^{\wedge})$$

Then through proper attachment of k – parameterized ring \bigoplus^k where $\partial = \coprod_{\omega=\infty} (\eta^0, \eta^1)^\omega / \sim$ and as described over [Points 1,2,3 in Results] one gets an equivalence relation for the operation being stated there over another affine (σ, ϵ) where σ represents space needed for the operations and ϵ represents the elements associated with the operator σ thereby making a 2-tuple relation suffice [1-3,5,11,13,14,25,28,26],

$$\epsilon \simeq \star \text{ for every } \epsilon \text{ operated under } \sigma \text{ over the suspension functor } \mathbf{Sus}^{\Psi^x}(\sigma, \epsilon)^+ \exists x = 0 \text{ or } 1 \text{ representing } (\eta^0, \eta^1)$$

Establishing [1,2,24,25],

$$\mathbf{Sus}^{\Psi^x}(\sigma, \epsilon) \simeq \mathcal{H}_g^{\wedge/} \coprod \mathcal{H}_g^{\wedge} = \mathcal{H}_4^{0/*} \cup \mathcal{H}_2^{1*} = \{(1,0), (2,0), (3,0), (4,0), (1,1), (2,1)\}$$

$$\exists \mathcal{H}_4^{0/*} = \{(1,0), (2,0), (3,0), (4,0)\} \text{ and } \mathcal{H}_2^{1*} = \{(1,1), (2,1)\}$$

$\forall \prod_{\omega=\infty}(\eta^0, \eta^1)^\omega / \sim$ with the annihilator \mathcal{A} representing globally non-commutativity for factors $\mathfrak{f}, \mathfrak{f}^{-1}$ in matrix multiplications.

Therefore, considering a chain of flow or orbits for (a trivial example to denote) if $\mathcal{H}_g^{\wedge/}$ initiates the orbit $\gamma \cong \prod k$ in R and goes through R^{op} with a reverse flow from R^{op} to R this will deduce the open or closed (where open means the break in continuous flow) representing the 4-tuple relation $(\tau(\nabla^y), t^\lambda, /_{\rho^\lambda}, \chi^\lambda)$ where all the 4-parameters would establish the required relation that I'll explain if and only if this 4-tuple relation holds true for the relation of $\mathbf{Sus}^{\Psi^x}(\sigma, \epsilon)^+$ via the same suspension functor $\eta \in \eta^0, \eta^1$ such that in the case $\sigma \Rightarrow (\tau(\nabla^y), t^\lambda, /_{\rho^\lambda}, \chi^\lambda)$ one can establish a Boolean $True(\eta^1)$ or $False(\eta^0)$ if [1-4,10,13,14,16,23-26],

$$\left. \begin{array}{l} \text{proper orbit } \gamma \xrightarrow{\text{ends up to}} \text{suspension orbit } \gamma \\ \text{suspension orbit } \gamma \xrightarrow{\text{ends up to}} \text{proper orbit } \gamma \\ \\ \text{proper orbit } \gamma \xrightarrow{\text{ends up to}} \text{proper orbit } \gamma \\ \text{suspension orbit } \gamma \xrightarrow{\text{ends up to}} \text{suspension orbit } \gamma \end{array} \right\} \eta \quad \begin{array}{l} | \text{ establishes } \eta^1 \\ \\ | \text{ establishes } \eta^0 \end{array} \quad \text{eq. (A)}$$

Over a consideration,

$$\text{For } \mathbf{Sus}^{\Psi^x}(\sigma, \epsilon)^+ \left\{ \begin{array}{l} \epsilon \simeq \star \text{ for every } \epsilon \text{ operated under } \sigma \text{ over the suspension functor } \mathbf{Sus}^{\Psi^x}(\sigma, \epsilon)^+ \\ \sigma \Rightarrow (\tau(\nabla^y), t^\lambda, /_{\rho^\lambda}, \chi^\lambda) \\ \text{for } \Psi^x, x \text{ takes the value 0 or 1 establishes eq. (A)} \\ \\ \text{above three relation suffice}_{\Downarrow} \\ \text{Global non - commutativity for annihilator } \mathcal{A} \end{array} \right.$$

Explained as,

Trajectory t^λ is defined in terms of orbit γ where the orbit denotes a continuous flow from $R \rightarrow R^{op} \rightarrow R \dots$ and if this flow continues over a period without any break that is a closed orbit then the value of λ in t^λ won't break and thus having a value of 1 for open orbit when the trajectory breaks and thereby induces a tension $\tau(\nabla^y)$ with y being a trivial parameter for the orientation if the orbit starts from R or takes the start from R^{op} where if the orbit is closed then this can be taken as a smooth flows over the hypercomplex structures which would regularize the structure been compact topological space but if the orbit is open which means the trajectory is not closed then the point where the orbit discontinues induces a tension from both ends of the discontinued fibers that is justifying the flow from a movement of a ring to opposite ring and so on... The induced tension by the break of the fiber would create a suspension over the hypercomplex structure where that suspension can create a genus for the periodic discontinuation of the flows if and only if there exists non-commutativity in the open orbits when the tension will impart a load $/_{\rho^\lambda}$ where $\lambda = 0$ appears in case of close orbit but for the open orbit with $\lambda = 1$ the number of genus would depend upon the strength of the gap between two ends of the broken fibers where the gap would make a proportionality relation and inverse proportionality relation through [7-9,18-21,27],

$$\begin{array}{l} \text{strength of gap } \rho \propto \text{number of genus} \\ \text{weakness of gap } \rho \propto 1/\text{number of genus} \end{array}$$

Therefore, this will depend upon the deformation χ^λ for the value of $\lambda = 1$ in case of open orbit thus making the suspension for the concerned functor $\mathbf{Sus}^{\Psi^1}(\sigma, \epsilon)^1$ where for each deformation or suspension or even the

¹ This suspension functor \mathbf{Sus}^{Ψ^x} is first introduced in the Preprint <https://doi.org/10.21203/rs.3.rs-1798323/v1>

regularization or the eraser of the suspension provided there exists commutativity for value 0 and non-commutativity for value 1 only iff for the infinite hypercomplex manifolds and not the two hypercomplex manifolds that I grouped to established the suspension there is non-commutativity and no commutativity would be structured over the Topological space \mathcal{J} when there arrives the annihilator \mathcal{A} [5,9-11,13,21,27].

Thus, the annihilator \mathcal{A} can only be established if η^0, η^1 be active elements (whereby active means that both the relation needs to suffice) for suspension η one gets the desired relation [1,5],

$$\text{for both } \eta^1, \eta^0 \text{ following eq. (A)} \left\{ \begin{array}{l} \left(\begin{array}{cc} \dagger & \dagger^{-1} \\ \dagger^{-1} & \dagger \end{array} \right) \text{ taken as } \alpha \\ \left(\begin{array}{cc} \dagger & \dagger \\ \dagger^{-1} & \dagger^{-1} \end{array} \right) \text{ taken as } \beta \end{array} \right. \Rightarrow \alpha \times \beta$$

$\neq \beta \times \alpha$ implies non – commutativity

6 Cobordism with Ring Homeomorphism for Potential ρ_ϕ

For the Topological space \mathcal{J} , assuming it to be compact having $d -$ dimensions then for the potential ρ_ϕ taken over the previously described two hypercomplex manifolds (I'll drop here all the indices to simply the notations) \mathcal{H} and \mathcal{H}' having $(d - 1) -$ dimensions then for the associated structure $\mathcal{S} \ni \mathcal{J}$; cobordism can be justified over 3-relations [22,27,28],

$$\left\{ \begin{array}{l} \rho_\phi = \mathcal{H} \coprod \mathcal{H}' \\ \partial\mathcal{S} = \rho_\phi \\ \text{cobordism satisfies through } \partial\mathcal{S} \end{array} \right.$$

Such that for $\partial\mathcal{S}$ two cases of dimensions are categorized as described above; \mathcal{H} and \mathcal{H}' having $(d - 1) -$ dimensions with \mathcal{S} of $d -$ dimensions there exists a ring homeomorphism for function $f: R \rightarrow R^{op}$ for the potential ρ_ϕ where the non-commutativity for the factors \dagger and \dagger^{-1} if one takes two elements ϕ and ϕ^0 in ring R then that non-commutativity will give two relations which can suffice the polynomials in formal power series]denoting rational numbers \mathbb{Q} in the $4k$ cohomology group then for that potential ρ_ϕ the rational coefficients can be defined for the vector bundles \mathcal{F} over \mathcal{J} defined through rational \mathbb{Q} and potential ρ_ϕ through [7,8,18-20],

$$\left\{ \begin{array}{l} \rho_{\phi_{(k)}}(\mathcal{F}, \mathbb{Q}) \\ H^{4k}(\mathcal{J}, \mathbb{Q}) \end{array} \right.$$

Where for genus g on \mathcal{H} and \mathcal{H}' is defined for manifold $\mathcal{S} \ni \mathcal{J}$ where cobordism is satisfied through $\partial\mathcal{S}$ over the relation [22,27,28],

$$\mathcal{S} \subseteq \mathcal{J} \text{ for } \rho_\phi \ni g(\mathcal{H} \coprod \mathcal{H}') = g(\mathcal{H}) + g(\mathcal{H}')$$

Therefore for the Boolean (0,1) there is a prominently identifiable η^0, η^1 suspension-category for a constant \mathbb{C} and a parameter φ sufficing non-commutativity in global scenario for $\omega = \infty$ in $\coprod_{\omega=\infty} (\eta^0, \eta^1)^\omega / \sim$ where ring homeomorphism is satisfied for the value of the potential $\rho_\phi = 1$ with the above mentioned relation $\mathcal{S} \subseteq \mathcal{J}$ satisfying cobordism through $\mathcal{X}(\mathcal{S})$ iff $\mathbb{C}, \varphi \ni R^{op}$ then [1,5,9,12,17,21,27,28],

$$\eta^1 \Rightarrow \mathbb{C} = \varphi = 1 \text{ and } \eta^0 = \begin{cases} \varphi^2 \neq \mathbb{C} \\ \mathbb{C} \neq 0 \end{cases}$$

Thus the global structure can be defined over the parameter ω over the topological space $\mathcal{S} \subseteq \mathcal{J}$ via the coefficient \mathcal{X} established over ρ_ϕ in the relation [14],

$$x_{\rho_\phi}(\mathcal{S}) = \int_{\mathcal{S}} \prod_{\omega=2}^{\infty} \eta$$

7 Discussion and Conclusion

Higher dimensional generalization has been made by extending the Preprint <https://doi.org/10.21203/rs.3.rs-1798323/v1> where any generalization or smoothness of the genus or the induced suspension in the compact topological space has been established through the two-grouping hypercomplex for open and closed orbit all though the 4-tuple relation being considered upon. However, for the global scenario there will be no commutativity for the two hypercomplex grouping; there will be non-commutativity and this has been topologically marked via cobordism and ring homeomorphism provided there exist a peculiar potential ρ_ϕ that ultimately suffice the Boolean for the concerned suspension category in rotations and inverse rotations through the complex planes where the genera are rational numbers \mathbb{Q} in the $4k$ cohomology group.

The k appearing in $4k$ cohomology group and in ring $R = \bigoplus^k$ where $\partial = \prod_{\omega=\infty} (\eta^0, \eta^1)^\omega / \sim$ are completely different although the notation appears to be the same.

Disclaimer

This paper is an extended version of a preprint document of the same author.

The preprint document is available in this link: https://figshare.com/articles/preprint/Non_commutativity_over_canonical_suspension_for_genus_g_1_in_hypercomplex_structures_for_potential_/21265989

[As per journal policy, pre-print article can be published as a journal article, provided it is not published in any other journal]

Competing Interests

Author has declared that no competing interests exist.

References

- [1] Bhattacharjee D. Suspension of structures in two-dimensional topologies with or without the presence of $g \geq 1$ genus deformations for canonical 22η stabilizer points. Res Square. 2022a, July 1. DOI: 10.21203/rs.3.rs-1798323/v1
- [2] Eisenbud D. Commutative algebra: with a view toward algebraic geometry (graduate texts in mathematics, 150). Springer; 1995.
- [3] Greene B. The elegant universe: superstrings, hidden dimensions, and the quest for the ultimate theory [reprint]. W W Norton & Company; 2010.
- [4] Pagliani P, Chakraborty M. A geometry of approximation: rough set theory: logic, algebra and topology of conceptual patterns (trends in logic, 27). Springer; 2010, November 30 (Softcover reprint of hardcover. 1st ed; 2008); 2008.
- [5] Deep Bhattacharjee. Generators of borel measurable commutative algebra on compact hausdorff taking von neumann AW* OVER *-ISOMORPHISM. EPRA International Journal of Research & Development (IJRD). 2022;122–124. Available:<https://doi.org/10.36713/epra11269>

- [6] Bhattacharjee D. Generalization of Grothendieck duality over Serre duality in deg Cohen-Macaulay schemes representing Calabi–Yau 3-fold on Bogomolov–Tian–Todorov Theorem. Res Square. 2022b. DOI: 10.21203/rs.3.rs-1781474/v1
- [7] Itenberg I, Mikhalkin G, Shustin E, I. Tropical algebraic geometry (Oberwolfach seminars, 35). Birkhäuser; 2009.
- [8] Marino M, Thaddeus M, Vakil R, Behrend K, Manetti M 2008, August 22. Enumerative Invariants in Algebraic Geometry and String Theory: lectures given at the C.I.M.E. Summer School Held in Cetraro, Italy, June 6-11, (Lect Notes Math. 1947) (2008th ed.). Springer; 2005.
- [9] Bhattacharjee D. Generalized Poincaré conjecture via Alexander trick over C-isomorphism extension to h-cobordism on inclusion maps with associated Kan-complex. Res Square. 2022e, July 7. DOI: 10.21203/rs.3.rs-1830184/v1
- [10] Bhattacharjee D. Imaginary Cycles of Permutations for Genus $g=3$. Available from: <https://easychair.org/publications/preprint/CWFx>. In: Complex geometries. EasyChair Preprint No. 8200; 2022.
- [11] Bhattacharjee D. Establishing equivalence among hypercomplex structures via Kodaira embedding theorem for nonsingular quintic 3-fold having positively closed (1,1)-form Kähler potential $i\bar{\partial}\partial^*\rho$. Res Square. 2022a. DOI: 10.21203/rs.3.rs-1635957/v1.
- [12] Bhattacharjee D. Violating the second law of thermodynamics in a dynamical system through equivalence closure via mutual information carriers of a 5-tuple measure space. Tech Rxiv; 2021. DOI: 10.36227/techrxiv.16854772.v1
- [13] Bhattacharjee D, Roy SS, Behera AK. Relating Enriques surface with K3 and Kummer through involutions and double covers over finite automorphisms on Topological Euler–Poincaré characteristics over complex K3 with Kähler equivalence. Res Square; 2022. DOI: 10.21203/rs.3.rs-2011341/v1
- [14] Bhattacharjee D. An outlined tour of geometry and topology as perceived through physics and mathematics emphasizing geometrization, elliptization, uniformization, and projectivization for Thurston’s 8-geometries covering Riemann over Teichmüller spaces. Tech Rxiv. 2022a, June 28. DOI: 10.36227/techrxiv.20134382.v1
- [15] Bhattacharjee D. Generalization of quartic and quintic Calabi – Yau manifolds fibered by polarized K3 surfaces. Res Square. 2022, August 16. doi: 10.21203/rs.3.rs-1965255/v1
- [16] Spanier EH. Algebraic topology (0003-1981. Corr. 3rd ed.). Springer; 1994.
- [17] Bhattacharjee D. Rigorously computed enumerative norms as prescribed through quantum cohomological connectivity over Gromov – Witten invariants. Tech Rxiv; 2022a. DOI: 10.36227/techrxiv.19524214.v1
- [18] Kosinski AA. Differential manifolds (Dover books on mathematics). Illustrated. Dover Publications; 2007
- [19] Gross M, Huybrechts D, Joyce D, Ellingsrud G, Olson L, Ranestad K et al. Calabi-Yau manifolds and related geometries. Springer; 2002.
- [20] Ricolfi AT. An invitation to modern enumerative geometry (SISSA springer series, 3). Springer; 2022.
- [21] Oh Y. Symplectic topology and Floer homology: Volume 1. Symplectic geometry and pseudoholomorphic curves. Cambridge University Press; 2015.

- [22] Bhattacharjee D. The gateway to parallel universe & connected physics. Preprints; 2021a.
DOI: 10.20944/preprints202104.0350.v1.
- [23] Penrose R. The road to reality: A complete guide to the laws of the universe (illustrated). Vintage Book Company; 2007.
- [24] Munkres JR. Topology updated. 2nd Ed. Pearson Education; 2021.
- [25] Arfken GB, Weber HJ, Harris FE. Mathematical methods for physicists. 7th ed. Elsevier India; 2022.
- [26] Boas ML. Mathematical methods in the physical sciences. 3rd ed. Wiley; 2005.
- [27] Publishing, L.S.I. Algebraic COBORDISM Notebook Algebraic COBORDISM Notebook. Journal. Independently published; 2020
- [28] Stong RE. Notes on cobordism theory (Princeton legacy library, 1879). Princeton University Press; 2015.

© 2022 Bhattacharjee; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here (Please copy paste the total link in your browser address bar)

<https://www.sdiarticle5.com/review-history/92716>