



Assessment of Genetic Variability in Physiological Traits on Chickpea in Three Sown Environments

**Pavan Dhopre^{a#}, Dhanesh Kumar Raidas^{b++},
Pradeep N. Dawane^{c++}, Vaqar Malik^{d‡} and N. Vinutha^{e‡}**

^a Department of Plant Breeding and Genetics, RVSKVV-RAK College of Agriculture, Sehore-466001 (MP), India.

^b Department of Plant Physiology, RVSKVV- RAK College of Agriculture, Sehore-466001 (MP), India.

^c Department of Agricultural Entomology, Regional Fruit Research Station, Katol, Nagpur-441302 (M.H.), India.

^d Department of Plantation, Spices, Medicinal and Aromatic Crop, RVSKVV- KNK College of Horticulture, Mandsaur-458001 (MP), India.

^e Department of Agronomy, Tamil Nadu Agriculture University, Coimbatore, Tamil Nadu-641003, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/CJAST/2022/v41i474028

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/95177>

Original Research Article

Received: 17/10/2022

Accepted: 24/12/2022

Published: 24/12/2022

ABSTRACT

The present studies were conducted to the estimation of genetic variability and heritability for physiological traits in chickpea (*Cicer arietinum* L.) under Randomized Completely Block design with two replications in the field of the department of Plant Breeding and Genetics, of RAK College of Agriculture, Sehore (MP), during the crop season 2019- 2020. The high heritability were observed

P.G. Student;

++Assistant Professor;

‡ Ph.D. Student;

*Corresponding author: E-mail: pavandhopre22@rediffmail.com;

for LA at 45 DAS (99.22%, 99.61%, 93.87%) & 60 DAS (99.87%, 99.88%, 98.91%), LAI at 45 DAS (99.18%, 99.6%, 93.64%) & 60 DAS (99.86%, 99.89%, 98.92%), TDM at 45DAS (97.21%, 9.13%, 99.05%) & 60 DAS (99.2%, 99.59%, 99.55%), NAR (98.06%, 97.24%, 92.9%), CGR (98.62%, 99.28%, 99.18%), chlorophyll index (SPAD) (95.23%, 94.72%, 82.59%) in normal, late and very late sowing conditions as well as the collection of genotypes for these traits could be useful in the above-mentioned environments. As a result, most of the characters were highly heritable and less influenced by the environment. The high Genetic advance as per mean was observed for LA at 45 DAS (73.17%, 91.15%, 60.49%) & 60 DAS (85.11%, 64.40%, 68.11%), LAI at 45 DAS (73.12%, 91.06%, 60.07%) & 60 DAS (85.07%, 64.37%, 68.15%), TDM at 45DAS (70.40%, 74.89%, 53.54%) & 60 DAS (71.79%, 65.27%, 39.49%), NAR (160.61%, 53.33%, 81.64%), CGR (79.16%, 73.90%, 60.10%), chlorophyll index (SPAD) (59.37%, 52.58%, 43.00%) in E-I, E-II & E-III sown environments, which suggested that these characters can be useful for selecting higher yielding genotypes.

Keywords: Chlorophyll index; environments; genetic advance; genotypes; heritability; LAI.

1. INTRODUCTION

Pulses, generally known as food legumes, belonging to family Fabaceae are an important group among staple pulse crops, next only to cereals providing vital protein and vitamins across the world. They form a major and cheapest source of dietary protein especially for vegetarians which form a major part of our Indian population. Pulses are rich source of minerals like calcium, phosphorus, iron and also certain essential amino acids. Chickpea is a food legume grown primarily in arid and semi-arid zones in India, where it is frequently subjected to drought and high temperatures during the reproductive stage. Chickpea flowering and podding are highly sensitive to changes in the external environment, with drastic reductions in seed yields reported when plants are exposed to high temperatures [1]. Chickpeas are well tailored for growth and pod filling in a range of temperatures 30-15°C [2]. Temperature plays a major role in pheno-phases *i.e.*, took longer days in cool regimes, than in the delayed sowing in chickpea genotype [3,4]. Temperature during sowing time from early to late and extra late directly related to vegetative and reproductive seasons. Sowing very much favorable for vegetative growth during mid and end of November as well as least stress on the reproductive phase because temperature did not change drastically, but several affected by growth of chickpea crop and grain filling time in December and January and genotype response also showed variations with different temperatures on pollen fertility and stigma sensitivity during reproductive period. For these reasons, the present investigation was carried out to assess the PCV, GCV, heritability, genetics advance of physiological traits on chickpea in three different environments.

2. MATERIALS AND METHODS

Twenty-five chickpea genotypes were grown in three environments (Normal 28 Nov. mid-late 28 Dec. and very- late 28 Jan.2020) during Rabi season 2019-20 under all India coordinated research project in chickpea in the experiment field of RAK College of Agriculture, Sehore (M.P.). The observations were identified on nine physiological traits such as leaf area (cm²) at 45 Days after sowing and 60 DAS, leaf area index (LAI) at 45 Days after sowing & 60 DAS, total dry matter (g) per plant at 45 Days after sowing & 60 DAS, net assimilation rate (g/cm²/day), crop growth rate (g/m²/day), chlorophyll index (SPAD) each of the three environments these were estimated from five randomly selected plants. The genotypes were organized in randomized complete block design with two replications. Data were subjected to statistical analysis to work out genotypic (GCV) and phenotypic (PCV) coefficients of variation, heritability and genetic advance as per cent of mean as per standard methods. Standard statistical procedure was used for the analysis of variance [5], genotypic and phenotypic coefficients of variation [6], genetic advance [7].

3. RESULTS AND DISCUSSION

The present investigation aimed to assess genotype variability (GCV, PCV, Heritability, Genetic advance) of physiological characters on chickpea in three environments (normal, mid-late, very-late), respectively. The analysis of variance for nine physiological characters of twenty-five chickpea genotypes derived from three environments and evaluated under normal, mid-late, very-late sown conditions are shown in (Table 1). The mean sum of squares due to various sources of variation

for physiological characters revealed that all three environments obtained significant differences, indicating that the variability among the selected genotypes was significant.

3.1 Phenotypic Coefficient Variation and Genotypic Coefficient Variation

The estimate of PCV were higher than an estimate of GCV for all most the traits, that suggested the apparent variation is not only due to genotypes but also due to the influence of environment. The estimations of the genetic variability parameters for various physiological traits are shown in (Table 2).

3.2 LA&LAI at 45 DAS and 60 DAS

The leaf area & leaf area index expresses the functional scale of the crop stand assimilation apparatus and has been used as the primary value for the measurement of other growth determinants [8,9]. Significant variations were found in LA & LAI at 45 and 60 DAS due to different climatic conditions. A progressive pattern of LA & LAI increase was observed under normal growing condition. The highest PCV were recorded for leaf area at 45 DAS (29.68%, 44.41%, 35.69%) & 60 DAS (41.36%, 31.29%, 33.42%), LAI at 45 DAS (29.65%, 44.37%, 35.59%) & 60 DAS (41.35%, 31.28%, 33.44%), as well as The highest GCV were noted for leaf area at 45 DAS (29.51%, 44.33%, 34.88%) & 60 DAS (41.34%, 31.28%, 33.24%), LAI at 45 DAS (29.47%, 44.28%, 34.75%) & 60 DAS (41.32%, 31.26%, 33.26%).

3.3 TDM at 45 DAS and 60 DAS

The present investigation indicated that the total dry matter at 45 DAS and 60 DAS significant variances were found due to heat stress plant growth was reduced in heat stress conditions. The highest PCV showed for total dry matter at 45 DAS (35.15%, 36.67%, 26.23%) Whereas, it was also highest for total dry matter at 60 DAS in E-I (35.13%) & E-II (31.81%). Medium PCV was noted for total dry matter at 60 DAS (19.25%) in E-III as well as highest GCV were recorded for total dry matter at 45 DAS (34.66%, 36.51%, 26.11%) in all three environments (E-I, E-II, E-III) and also total dry matter at 60 DAS in E-I (34.99%) & E-II (31.75%) recorded highest values. Whereas, the medium

GCV was noted for total dry matter at 60 DAS (19.21%) in E-III.

3.4 NAR and CGR

High estimates of PCV were observed for NAR and CGR. Similar trend was identified at genotypic level also. Thus, the present investigation revealed that the existence of sufficient genetic variability in the population and there is a lot of scope for achieving desirable improvement. The difference between phenotypic coefficient of variation and genotypic coefficient of variation was high for these traits. This suggests that the expression of these traits was least affected by the environmental factors and their phenotype is the true representative of its genotype. Further, the selection on the basis of per cent performance will be effective. In normal date of sowing NAR ranged 0.006032 to 0.066939 (g/cm²/day) and in late sowing ranged 0.002621 to 0.0124 (g/cm²/day) and also the CGR ranged in E-I normal sown conditions was 8.93 to 36.24 (g/m²/day) and 3.51 to 16.2 (g/m²/day) in late sowing ranged. The highest PCV were recorded for NAR (79.5%, 26.61%, 42.65%), CGR (38.96%, 36.13%, 29.41%) in all environmental conditions (E-I, E-II, E-III) as well as the highest GCV were reported NAR (78.73%, 26.25%, 41.11%), CGR (38.69%, 36%, 29.29%) in all three environments (E-I, E-II, E-III).

3.5 Chlorophyll Index

Heat stress can also cause damage to photosynthetic apparatus such as increased photo oxidation of chlorophyll and inhibition of chlorophyll biosynthesis and damage in lamellar membrane of chloroplast. The chlorophyll index was reduced when temperature get increased due to high temperature synthesis of chlorophyll was negatively affected which resulted chlorophyll index and photosynthesis rate reduction in high temperature conditions. In normal date (E-I) of sowing chlorophyll index ranged was 20.23 to 67.26 in E-I, 23.41 to 74.05 in E-II and 18.32 to 53.13 SPAD in late date (E-III) of sowing. Reduction of mean value from E-II to E-III was 15.89 SPAD due to high temperature. The highest PCV were recorded for chlorophyll index (30.26%, 26.94%, 25.27%) in all environmental conditions (E-I, E-II, E-III) as well as the highest GCV were showed chlorophyll index (29.53%, 26.22%, 22.97%) in all three environments (E-I, E-II, E-III).

Table 1. Analysis of variance for physiological characters in chickpea genotypes over three environmental conditions

Source of variations	ENV	d.f	LA at 45 DAS	LA at 60 DAS	LAI at 45 DAS	LAI at 60 DAS	TDM at 45 DAS	TDM at 60 DAS	NAR	CGR	Chlorophyll Index
Replications	E1	1	526.46	600.88	0.006	0.006	0.001	0.031	3.49x10 ⁻⁶	0.11	0.03
	E2	1	86.06	84.68	0.001	0.001	0.036	0.29	4.21x10 ⁻⁵	0.62	7.43
	E3	1	541.75	1194.98	0.007	0.013	0.147	0.153	1.88x10 ⁻⁷	0.001	0.02
Genotypes	E1	24	5308.01**	72025.2**	0.059**	0.79**	2.22*	34.56**	3.37x10 ^{-4**}	116.28**	293.1**
	E2	24	28808.09**	14393.22**	0.32**	0.32**	2.15*	24.94**	9.53x10 ^{-3**}	88.04**	329.26**
	E3	24	7483.98**	46262.22**	0.083**	0.515**	0.697	3.14**	1.39x10 ^{-5**}	15.47**	122.22**
Error	E1	24	30.74	44.24	0.00	0.001	0.031	0.13	3.28x10 ⁻⁶	0.80	7.15
	E2	24	55.25	80.34	0.001	0.001	0.009	0.05	1.33x10 ⁻⁴	0.31	8.92
	E3	24	173.04	252.1	0.002	0.003	0.003	0.007	5.12x10 ⁻⁷	0.063	11.65

Table 2. Genetic parameters of variation for physiological traits over three environmental conditions

Characters	ENV.	Mean	Range		h ² (bs) (%)	GCA (%)	PCA (%)	Genetic advance	GA as % of mean
			Max.	Min.					
LA at 45 DAS	E-I	180.24	371.34	90.06	99.22	35.66	35.80	131.90	73.17
	E-II	270.47	527.87	116.10	99.61	44.33	44.41	246.52	91.15
	E-III	170.31	290.82	82.55	93.87	30.31	31.28	103.02	60.49
LA at 60 DAS	E-I	458.88	1027.17	193.50	99.87	41.34	41.36	390.56	85.11
	E-II	856.97	1321.76	382.43	99.88	31.28	31.29	551.90	64.40
	E-III	456.27	804.27	161.06	98.91	33.24	33.42	310.75	68.11
LAI at 45 DAS	E-I	0.60	1.23	0.30	99.18	35.64	35.78	0.43	73.12
	E-II	0.90	1.75	0.38	99.6	44.28	44.37	0.82	91.06
	E-III	0.57	0.96	0.27	93.64	30.13	31.14	0.34	60.07
LAI at 60 DAS	E-I	1.52	3.42	0.64	99.86	41.32	41.35	1.30	85.07
	E-II	2.85	4.40	1.27	99.89	31.26	31.28	1.84	64.37
	E-III	1.52	2.68	0.53	98.92	33.26	33.44	1.04	68.15
TDM at 45 DAS	E-I	3.01	4.91	1.08	97.21	34.66	35.15	2.12	70.40
	E-II	2.83	4.72	1.05	99.13	36.51	36.67	2.12	74.89
	E-III	2.25	3.61	1.07	99.05	26.11	26.23	1.21	53.54

Characters	ENV.	Mean	Range		h ² (bs) (%)	GCA (%)	PCA (%)	Genetic advance	GA as % of mean
			Max.	Min.					
TDM at 60 DAS	E-I	11.85	20.18	5.77	99.2	34.99	35.13	8.51	71.79
	E-II	11.11	18.92	4.96	99.59	31.75	31.81	7.25	65.27
	E-III	6.51	9.32	4.02	99.55	19.21	19.25	2.57	39.49
NAR	E-I	0.01641	0.06694	0.006032	98.06	78.73	79.5	0.03	160.61
	E-II	0.261225	0.40538	0.104167	97.24	26.25	26.61	0.14	53.33
	E-III	0.006301	0.0124	0.002621	92.9	41.11	42.65	0.01	81.64
CGR	E-I	19.63	36.24	8.93	98.62	38.69	38.96	15.54	79.16
	E-II	18.39	33.53	7.60	99.28	36	36.13	13.59	73.90
	E-III	9.47	16.2	3.51	99.18	29.29	29.41	5.69	60.10
Chlorophyll index (SPAD)	E-I	40.48	67.26	20.23	95.23	29.53	30.26	24.03	59.37
	E-II	48.25	74.05	23.41	94.72	26.22	26.94	25.37	52.58
	E-III	32.36	53.13	18.32	82.59	22.97	25.27	13.92	43.00

3.6 Heritability

Heritability estimated in broad sense, is the ratio of genotypic variance to the phenotypic variance and is expressed in percentage. It is an index of transmission of a character from parents to their offspring. It helps the plant breeders in the selection of superior genotypes from the genetically variable population. Heritability into broad sense as high (above 60%), medium (30-60%), and low (below 30%) [10]. The high heritability were observed for LA at 45 DAS (99.22%, 99.61%, 93.87%) & 60 DAS (99.87%, 99.88%, 98.91%), LAI at 45 DAS (99.18%, 99.6%, 93.64%) & 60 DAS (99.86%, 99.89%, 98.92%), TDM at 45DAS (97.21%, 9.13%, 99.05%) & 60 DAS (99.2%, 99.59%, 99.55%), NAR (98.06%, 97.24%, 92.9%), CGR (98.62%, 99.28%, 99.18%), chlorophyll index (SPAD) (95.23%, 94.72%, 82.59%) in normal, late and very late sowing conditions. The high heritability was also observed for all physiological parameters in normal late and very late planting conditions, as well as the collection of genotypes for these traits could be useful in the above-mentioned environments. As a result, most of the characters were highly heritable and less influenced by the environment.

3.7 Genetic Advance as % of Mean

Expected genetic advance as percent of mean is the product of selection intensity, heritability and phenotypic standard deviation. Heritability in conjunction with genetic advance would give a more reliable index for selection. More genetic advance could be expected from a population with wide variability, and high heritability estimates. The genetic advance (GA) for physiological traits as per mean in per cent was reported in Table 2. The high Genetic advance as per mean was observed for LA at 45 DAS (73.17%, 91.15%, 60.49%) & 60 DAS (85.11%, 64.40%, 68.11%), LAI at 45 DAS (73.12%, 91.06%, 60.07 %) & 60 DAS (85.07%, 64.37%, 68.15%), TDM at 45DAS (70.40%, 74.89%, 53.54%) & 60 DAS (71.79%, 65.27%, 39.49%), NAR (160.61%, 53.33%, 81.64%), CGR (79.16%, 73.90%, 60.10%), chlorophyll index (SPAD) (59.37%, 52.58%, 43.00%) in E-I, E-II & E-III planting condition scan be used for selecting higher yielding genotypes.

4. CONCLUSION

To present study, the nine physiological traits were observed that significantly superior in E-I

followed by E-II sowing conditions. the physiological traits were identified such as total dry matter at 60 DAS, CGR and chlorophyll index were best performed in E-I planting conditions while, LA & LAI at 45 & 60 DAS, total dry matter at 45 DAS were in E-II. The total dry matter was recorded least affected in E-I, E-II, E-III sowing conditions and chlorophyll index was the high affected in three planting conditions.

ACKNOWLEDGEMENT

We are thankful to the Department of Plant Breeding & Genetics and Department of Plant Plant Physiology for providing lab facilities and with thanks for the research farm of RAK College of Agriculture, Sehore (M.P.) for conduct of experiments. I sincerely thanks to my advisory members and other scientist for their valuable time for data analysis.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Bahuguna RA, Shah D, Jha J, Pandey SK, Khetarpal S, Anand A, Pal M. Effect of mild temperature stress on reproductive dynamics and yield of chickpea (*Cicer arietinum* L.). Ind. J. Plant Physiol. 2012;17:1-8.
2. Basu PS, Ali M, Chaturvedi SK. Terminal heat stress adversely affects chickpea productivity in Northern India –strategies to improve thermo tolerance in the crop under climate change. 2011:189-192.
3. Clarke HJ. Improving tolerance to low temperature in chickpea. In: 4th European Conference on Grain Le-gumes. Towards the Sustainable Production of Healthy Food, Feed and Novel Products. Cracow, Poland. 2001:34–35.
4. Kiran BA. Effect of temperature regimes on productivity of chickpea genotype. University of Agriculture Sciences, Dharwad, Karnataka. J. Agric. Sci. 2014; 28(2):168-171.
5. Fisher RA. The correlation among relatives on the supposition of mandelian inheritance. Trans. Royal Soc. of Edinburgh. 1918;52:399-433.

6. Burton GW. Quantitative inheritance of grasses. Proc. 6th Int. Grassland congress. 1952;1:277-283
7. Johnson HW, Robinson HF, Comstock RE. Estimation of genetic and environmental variability in soybeans. Agron. J. 1955;47: 314-318.
8. Watson DJ. The physiological basis of variation in yield. Adv. In Agron. 1952; 14(4):101-145.
9. Watson DJ. Comparative physiological studies on the growth of field crops variation in net assimilation of leaf area between species and varieties within and between years. Ann, Bot. 1947;11:41-76.
10. Robinson HF, Comstock RE, Harvey PH. Estimates of heritability and degree of dominance in corn. Agronomy J. 1949; 41:253-259.

© 2022 Dhopre et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

*The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/95177>*