

Asian Journal of Food Research and Nutrition

Volume 1, Issue 4, Page 37-49, 2022; Article no.AJFRN.94710

Harnessing the Potentials of Sweet Potato (*Ipomoea batatas* L.) for Enhanced Livelihood and Development

Kwame Yaw-Kan Kojo^{a*} and Naomi Adoma Fosu^a

^a Roots and Tubers Division, CSIR-Crops Research Institute, P.O. Box 3785, Kumasi, Ghana.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: https://www.sdiarticle5.com/review-history/94710

Review Article

Received: 09/10/2022 Accepted: 15/12/2022 Published: 24/12/2022

ABSTRACT

This review gives an overview of the potentials of Sweet potato as a versatile and multi-functional crop. It brings together careful reviews of some of the outstanding characteristics and revealing potentials of the crop which have been established and reported by several researches and authors. The contents of this review contains reviews on components such as the various types of Sweet potato based on the flesh color and their various constituent compounds, nutritive features, phenolic acids contained in Sweet potato, the phytochemical and pharmacological properties such as diseases' prevention and promotion of good health through its antioxidant, anticancer, antiulcer, anti-inflammation, cardiovascular properties and the enhancement of the immune system, processing of Sweet potato in to various food products such as vinegar, chips, flakes, cubes, yoghurt, cheese and some industrial utilization in the processing of alcohol and starch, the utilization potentials in animal feed; use as silage, processed to hay, as green-chop, the use of the tuber in poultry feeds and the contributions of the crop towards food security and development through the ability to fight hunger, poverty reduction, income generation, climate resilient and adaptability to vast range of climates. Various authors and researches have investigated and voted Sweet potato as one of the crops with the best production, consumption and utilization attributes

^{*}Corresponding author: E-mail: kkyawkan9@gmail.com;

which can help the smallholder farmers in the forest, tropical and subtropical zones to achieved food security and households' income stability. Sweet potato is gaining attention and registering growing demands with appreciating growth in exports and imports in some countries across the globe due to its quality and potentials. The developing countries in particular which have the comparable advantages in the cultivation of Sweet potato should develop schemes and policies that will enable them to tap in to its potentials for livelihood improvement and development.

Keywords: Anthocynin; carotene; phytochemical; pharmacological; sweet potato.

1. INTRODUCTION

Sweet potato is a root-tuber crop which is rated as the 5th and 7th most principal food crop in the less developed countries and globally respectively [1]. Sweet potato comes next after cassava among the root and tuber crops, making it the second most valuable tuber crop in the world [2]. Sweet potato is the most prominent among the members of the Ipomoea genus, and it is by far the only member in the genus with the characteristics of bearing root tubers of great nutrition and economic values [3]. Sweet potato is regarded as one of grown the important crops for man's consumption across the globe. Sweet potato, among other crops is of economic advantage and has been rated as the 15th and the 29th cultivated crop on the account of the amount of vield and the area under cultivation respectively. with the average annual yield of 92, 000000 tons [4-6].

Sweet potato is arguably an important tuber crop in most areas, cultivated in over 100 countries across the globe and has been regarded as a staple food for the people in the tropical and subtropical regions, which provides nutritional security for both the urban and rural dwellers through enhanced cultivation and utilization [7,8]. Sweet potato is grown around the world in about 114 nations, with the global average yield of 14 tons/hectare [9]. Sweet potato among other staple crops is well adaptable to wide range of climates, rapid and short maturity and promising yield capacity [9].

2. SWEET POTATO TYPES, NUTRITIONAL AND CHEMICAL CONSTITUENTS

2.1 Types of Sweet potato According to Flesh Color

There are different types of Sweet potato varieties growing across the globe, with variation in skin and flesh color such as white, cream, orange, purple, pale yellow, etc. these types of Sweet potato do not only vary in the fleshed or skin color, but also possess different levels and kinds of important chemical constituents of enviable relevance in nutrition, medicine and other industrial usage (Table 1).

2.2 Nutritional Components of Sweet Potato

Sweet potato is a rich source of various important nutrient ingredients such as protein, energy, carbohydrates, Lipids, minerals and vitamins [14] (Fig. 1).

The flour produce from the tubers of Sweet potato also contains some sugars, which contribute greatly to the dough fermentation in the process of making bread [16] (Fig. 2).

2.3 Phenolic Acids Present in Sweet Potato

The various varieties of Sweet potato generally contain a number of phenolic acids which are importantly linked to the unique sensory traits possess by the varieties (Fig. 3).

Table 1. Types of Sweet potato according the fleshed color and their major compounds

Sweet potato type	Major bioactive compound contained	Reference
Purple fleshed	Anthocyanins, phenolic acids	[10, 11]
Orange fleshed	β-5 cryptoxanthin, $β$ -carotene, $α$ -carotene, flavonoids	[8,12]
White fleshed	Starch, phenolic acids	[13, 11]
Cream fleshed	Starch, phenolic acids	[13, 11]

Kojo and Fosu; Asian J. Food Res. Nutri., vol. 1, no. 4, pp. 37-49, 2022; Article no.AJFRN.94710




Fig. 1. Nutritional composition of sweet potato [14,15]




Fig. 2. Sugars identified in sweet potato flour

Kojo and Fosu; Asian J. Food Res. Nutri., vol. 1, no. 4, pp. 37-49, 2022; Article no.AJFRN.94710




Fig. 3. Phenolic acids present in sweet potato [17,18]




Fig. 4. Main phytochemical constituents of Sweet potato [19,20]

3. BENEFITS AND POTENTIALS OF SWEET POTATO

3.1 Industrial Utilization

The tubers of Sweet potato can be used to produce starch, which is indigenously used to produce noodles, sugar, vermicelli, syrups, contributing significantly to food products (Fig. 4). Additionally the starch and the sugar obtained from it can also be used to make products such as monosodium, citric acid, microbial enzymes, glutamate, lactic acid and alcohol [9,20]. Sweet potato are also used in the manufacturing of snacks and bakery foods like chips [21]. The purple and orange fleshed varieties of Sweet potato are reported to have been extensively applied in the making of inartificial anthocyanin colorants and β -carotene in some beverage products on a large scale in Japan [9]. The purple and orange fleshed Sweet potato are used in making beer and wine, due to their characteristics of being rich sources of starch which is convertible to sugar [9]. Sweet potato provides good industrial raw materials for the producing of alcohol and starch [22].

Maintaining and scaling up the growth and expansion of the Sweet potato sub-sector through processing and formulation of food products is directly proportional to the commercialization and value addition of Sweet potato, due to its richness in polyphenols and dietary fiber, which have the potentials of reducing the digestibility of starch [23,24].

Alpha-amylase: the beta-amylase enzyme, oxidase causes the breaking down of the starch component of Sweet potato during processing [25], and the enzymatic reaction of the amylase results to maltose formation during processing leading to the high level of sweetness and sugar content of the end product formed [26].

The Mitsui and Toyota motor companies are collaboratively exploring the potential of Sweet potato in the manufacturing of plastics materials which are bio-degradable, and has also projected Sweet potato as a potential efficient energy source like hydrogen and alcohol in the day ahead of us [27] (Fig. 5).

3.2 The Pharmacological and Health Potentials of Sweet Potato

There are several therapeutic, pharmacological, phytochemicals and other health related

attributes associated with Sweet potato, which have been studied and their relevance being investigated. Sweet potato is an excellent source of several phytochemical compounds of therapeutic importance (Table 2). Nutritionally, the various parts of sweet potato have great dietary fiber, the stem and leaves have high carotene, vitamin E, vitamin C, vitamin B2 and iron [28]. The sweet potato roots are good sources of vitamin C, starch, β -carotene and sugar [29]. It is reported that the tubers of Sweet potato have high protein level than that of yam and cassava [30].

3.3 The Uses of Sweet Potato in Feed and Feed Ingredients of Animals

The vegetative parts and the tuber of Sweet potato are use as silage and fodder for livestock [49]. There was improvement in the growth of young female cattle when fed with roughage diet, supplemented with silage prepared from the vines of Sweet potato [50]. According to Mibach et al. [51] substituting of ground-corn for any amount of Sweet potato flour for sheep put on diets of multiple constituents does not interferes with the intake and digestibility of dry matter, nitrogen retention and microbial synthesis of protein. According to Murugan et al. [52] averagely 43% of the Sweet potato produced per annum globally is utilized as feed for animals; the vines and other foliar parts of Sweet potato are commonly use as protein supplementary feed and feed materials for farm animals like pigs in several countries across the globe such as Uganda, Korea, Taiwan, Vietnam, Indonesia, Papua New Guinea, China, Philippines and India.

The tubers of Sweet potato has been studied and identified as one of the nutritious and less expensive alternative feed material for broiler chickens [52,53]. The addition of about 25% and 20% of the orange fleshed Sweet potato tubermeal to layer and broiler chickens feeds respectively do not has any negative impact on the quality of eggs and growth [54]. Sweet potato-meal inclusive diet has proven to have an enhancer influence on the growth and performance of quails in Japan [55]. The most desirable feed conversion efficiency and growth can be achieved in chicken, when fed with diet containing about 10% of dried Sweet potatoleaves, and can result to more weight gain in broilers in particular compare with other diets containing no Sweet potato leaves [56]. A diet of dried leaves of Sweet potato can increase the

level of the yellow color of the egg yolk and the skin of chickens [57]. The meal prepared from the tubers of the orange fleshed Sweet potato has been suggested with proven results, to be the best substitute for corn at the rate of 12% -16% and 15% and 20% during the starter and grower stages of chicken respectively, with positive influence on chicken characteristics such as ingestion, weight increase and meat attributes [58]. Meal prepared from Sweet potato tubers can be included at the levels of 25% and 50% in the diets of layer and broiler chickens respectively to achieved optimal production [59].

3.4 The Potentials of Sweet Potato in Fostering Food Security

According to Scott and Maldonado [60] Sweet potato is largely cultivated in low and middle income earning countries, and because of that, scaling up of Sweet potato cultivation is noted as a greater tool to foster food security and improve the livelihoods and living standards of the economically disadvantaged class of the rural and urban settings.

Sweet potato plays several fundamental functions, contributing significantly to the "food

basket" of the globe, which has a viable potential of reducing the food need of the growing world's population, decreases poverty, promotes food security and improves livelihoods [2]. Sweet potato tubers are great hunger-combating food materials which significantly saved millions of people during some daring moments in the history of the world such as the period critical food insecurity at china (in the 1960s), Uganda (during the 1990s) and Japan, that is the time during which there was a development of phytosanitary problems in fields of cassava in Uganda and the destructions of rice farms by typhoons in Japan [61,2].

In addition to food, Sweet potato has got multiindustrial uses because of its wonderful attributes such as cheap and simple cultivation, ability to tolerate infections and pests, high output and rich in carbohydrate, minerals and vitamins in both its roots and leaves [62]. The vitamin A precursor (Beta- carotene) rich orange fleshed Sweet potato can importantly contributes to the global fight against the deficiency of vitamin A among people, which has started yielding positive results in improving status quo of vitamin A among lactating and expectant mothers and children in Africa [9,63].




Fig. 5. Food and industrial products from sweet potato [9]

Kojo and Fosu; Asian J. Food Res. Nutri., vol. 1, no. 4, pp. 37-49, 2022; Article no.AJFRN.94710

Property	Mechanism of reaction	References
Anti-oxidant action	Flavonoids and n-hexane, ethanol and ethyl- acetate extracts from leaves help to remove oxygen-based reactive radicals in the cell. Reduces the chances of getting degenerative disorders like senile, dementia, cancer, diabetes, asthma, which are highly caused by the actions of free harmful radicals.	[31,32] [33]
Boost of immune system	The leaves of purple Sweet potato regulate the activities of lytic, T-lymphocyte deleterious cell.	[34]
	Generation of antibody production in healthy humans The derivatives of ethyl-acetate from Sweet potato have immune-modulatory actions	[35]
Anti-cancer property	The polyphenol constituent of Sweet potato extract hinders the proliferation and also leads to apoptosis in prostate cancer cells The anthocyanin in the Sweet potato with purple flesh is potent against the development of cancers of the breast and stomach.	[36,37]
Anti-diabetic property	The white-fleshed Sweet potato increases the adiponectin level in the blood, a hormone that	[38]
	modifies the metabolism of insulin. Extracts of white-fleshed Sweet potato potentially reduces blood sugar level	[39]
Antimicrobial property	Extracts of ethanol and acetone obtained from the leaves of Sweet potato respectively have antimicrobial inhibitory actions on <i>Pseudomonas</i> <i>aeruginosa</i> and <i>Salmonella typhimurium</i> Liquid extracted from the leaf of Sweet potato highly inhibits the growth of <i>E. coli, Bacillus cereus</i> and <i>Staphylococcus aureus</i>	[40] [41]
Cardiovascular enhancing property	Extract rich in phytochemicals, obtained from the leaves of Sweet potato inhibits the oxidation of less dense lipoprotein in humans.	[42]
	Anthocyanin in purple skinned Sweet potato decreases the chances of getting coronary infections.	[43]
Anti-ulcer property	Extract rich in methanol, obtained from the roots of Sweet potato protects the stomach of rat against	[44]
	ulcer, induced by aspirin. The flour obtained from the roots of Sweet potato inhibits gastric ulcer induced by ethanol.	[45]
Anti-inflammatory property	Inhibition of inflammatory infection of the brain by preventing the inflammatory reactions caused by lipopolysacharide.	[46]
Haematological property	The leaf of Sweet potato has haematinic action, and it is used in indigenous treatment of anaemia.	[47]
	Powder from the leaf of Sweet potato increases the levels of red blood cells and haemoglobin of mice.	[48]

Table 2. Pharmacological properties of sweet potato

Poverty reduction

- •Source of sustainable income for small holder farmerss
- •Generation of biofuel due to richness in starch
- •Rapid multiplication of vines for sale
- •High yields of tubers for food and sales

•[72-74]

Fight against hunger and malnutrition

- Highly nutritious foods from the leaves and tuberss
- •Supply tubers for food at lean season for households in some countries
- does well on poor soils and has low cost of production and short maturity
- Rich in vitamins, macro and micronutients
 [75-76]

Promote resiliency against climate change on peasant farming system

- •It requires less water and has short maturity
- •It is tolerant to drought and does well in environment of higher level of carbon dioxides
- •Use in making bioethanol for stoves, serves alternative way of reducing greenhouse gases emission by petroleum products •[77-82]

Fig. 6. The roles of sweet potato in promoting food security

In the populous semi-arid plains of Eastern Africa, Sweet potato is a cherishable food material used to combat food insecurity by thousands of rural communities [64]. The development and demands of snack-foods developed from Sweet potato have seen a surge in recent years, covering a significant portion of the food commodity market, meeting the taste and preference of consumers [65]. Sweet potato cultivation is very profitable, with high sale-value and has the potentials of earning its growers more incomes [66].

Okoye [67] has noted that Sweet potato has a good comparative advantage in Nigeria, in terms of its production and small holder farmers earn significant amount of income from the vines, tubers and the leaves. In terms of nutrients and biomass per hectare, Sweet potato is very excellent and tops all other crops worldwide, and as such it is considered as one of the best crops with the capacity to fight hunger and poverty [63]. Various primary and secondary processed food products such as puree, flour, chips, noodles, pasta and biscuits can be made from the tubers of Sweet potato [68], and therefore, it can be investigated in advance level as a key crop for the formulation of sustainable nutritionally resourced food items (Fig. 6).

The tubers of the purple fleshed type of Sweet potato contribute to the keeping of the body stable; they can best substitute rice in our food menu due to their richness in carbohydrate, and also contain more nutrients than the white rice [69]. Ice cream is made from Sweet potato [70] and the purple type Sweet potato can best be used in place of corn in terms of starch for bread making [71].

4. FUTURE RESEARCH FOCUS AND PROSPECTS

Sweet potato (Ipomoea batatas L.) is noted globally as one of the great cultivated crops among the roots and tubers, being a cheap and sustainable source of nutrition. It has several underutilized health advantages, market and export potentials and above all, the crop and its products possess greater prospects in the areas of sustainable food, nutrition and income security for every player along its value chain in Africa, most particularly in the Sub-Saharan African sub region where majority of the rural peasant farming populations are bedeviled with acute hunger, income instability and poverty. The starch of Sweet potato is of promising industrial importance, and can be explored vastly by research to develop more starch production efficient varieties for expansion of the processing industry. The molecular, chemical and mineral richness as well as the medicinal potentials of the edible portions of Sweet potato could be exploited in to detail and well incorporated in to infant foods, other dietary packages for children to improve health and reduce the incidence of infantile minerals and nutritional deficiencies related diseases. The crop is naturally endowed with anthocyanin, anticancer, anti-inflammatory, antibacterial and numerous other disease fighting properties, which can be further explore in medical and pharmaceutical products such as capsules, mixtures, ointment for enhance administering and use. Currently, less priority is given to research focus on the development of more industrially efficient varieties, especially for starch, medicinal and pharmaceutical purposes. Some national and regional agricultural institutions in Africa have demonstrated little weak commitment towards interests and investment in the Sweet potato sub-sector by helping rural and urban growers to expand production and helping to create the enabling local and international market spaces to fully exploit the market and export opportunities of the crop and its related products. Sweet potato is a kind of crop which is climate smart, cheap to cultivate and serves as a good source of sustainable income and food security, and as such. developing nations that have the comparable advantages in Sweet potato cultivation can invest adequately in the sector to help improve livelihoods, and income for rural development and growth of national economy.

5. CONCLUSIONS

This comprehensive review has revealed some of the promising potentials of Sweet potato. These characteristics show how multipurpose and versatile the crop is. It has the capacity to fight hunger, employment and income generations, reduction in poverty and raising the standard of living in the. Sweet potato has the capacitv to improve lives and promote development if the potentials along its value chains from production to processing and utilizations are well harnessed. It is nutritionally rich and can be processed industrially and locally in to several food products and feed forms. Major attention from the research community should be given to the development of Sweet potato varieties that are viable for industry and value addition. The numerous potentials of the crop should be adequately harnessed for development.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Mohanraj R. Sweet potato: Bioactive compounds and health benefits. In the

Bioactive Molecules in Food. Reference Series in Phytochemistry; Mérillon JM, Ramawat KG, Eds., Springer International Publishing, Cham, Switzerland. 2018:1-16.

- 2. El-Sheikha AF, Ray RC. Potential impacts of bioprocessing of Sweet potato: Review. Critical Review of Food Science and Nutrition. 2017;57:455-471.
- Cartabiano-Leite CE, Porcu OM, De Casas AF, Sweet potato (*Ipomoea batatas* L.) nutritional potential and social relevance: A review. International Journal of Engineering, Research and Applications. 2020;10(6):23-40. ISSN: 2248-9622
- 4. FAO. STAT. Food and Agriculture Organization Statistics, Food and Agricultural Association of the United Nations Database; 2018.
- 5. FAO. STAT. Food and Agriculture Organization Statistics, Food and Agricultural Association of the United Nations – Database; 2019.
- 6. Mu TH, Li PG. Chapter 2 Sweet potato: Origin and production, sweet potato. Academic Press. 2019:5-25.
- Alam MK. A comprehensive review of sweet potato (*Ipomoea batatas* L.) Lam): Revisiting the associated health benefits. Trends in Food Science and Technology. 2021;115:512-529.
- Alam MK, Rana ZH, Islam SN. Comparison of the proximate composition, total carotenoids and total polyphenol content of nine orange-fleshed sweet potato varieties grown in Bangladesh. Foods. 2016;5(4):64.
- Truong VD, Avula RY, Pecota KV, Yencho GC. Sweet potato production, processing, and nutritional quality." Handbook of Vegetables and Vegetable Processing, Volume II, Second Edition. Edited by Muhammad Siddiq and Mark A. Uebersax. John Wiley and Sons Ltd., USA; 2018.
- Teow CC, Truong VD, McFeeters RF, Thompson RL, Pecota KV, Yencho GC. Antioxidant activities, phenolic and βcarotene contents of Sweet potato genotypes with varying flesh colours. Food Chemistry. 2007;103:829-838.
- 11. Padda MS, Picha DH. Quantification of phenolic acids and antioxidant activity in Sweet potato genotypes. Science of Horticulture. 2008;119:17-20.
- 12. Park SY, Lee SY, Yang JW, Lee JS, Oh SD, Oh S, Lee SM, Lim MH, Park SK, Jang JS, et al. Comparative analysis of

phytochemicals and polar metabolites from colored sweet potato (*Ipomoea batatas*, L.) tubers. Food Science and Biotechnology. 2016;25:283-291.

- Tumwegamire S, Kapinga R, Rubaihayo PR, LaBonte DR, Grüneberg WJ, Burgos G, Felde TZ, Carpio R, Pawelzik E, Mwanga ROM. Evaluation of dry matter, protein, starch, sucrose, β-carotene, iron, zinc, calcium, and magnesium in East African Sweet potato (*Ipomoea batatas* L.) germplasm. Horticultural Science. 2011;46: 348-357.
- 14. USDA. United States Department of Agriculture, Agricultural Research Service, Nutrient Data Laboratory, USDA National Nutrient Database for Standard Reference, Release. 2015;28.
- 15. Fanta SW, Neela S. Review on nutritional composition of orange-fleshed sweet potato and its role in management of vitamin a deficiency. Food Science and Nutrition. 2019;7:1920-1945.
- Musilová J, Bystrická J, Árvay J, Harangozo L. Polyphenols and phenolic acids in sweet potato (*Ipomoea batatas*, L.) roots. Potravin. Slovak Journal of Food Science. 2017;11:82-87.
- Olatunde GO, Henshaw FO, Idowu MA, Tomlins K. Quality attributes of sweet potato flour as influenced by variety, pretreatment and drying method. Food Sci. Nutr. 2016;4(4):623–635.
- Rumbaoa RGO, Cornago DF, Geronimo IM. Phenolic content and antioxidant capacity of Philippine sweet potato (*Ipomoea batatas* L.) varieties. Food Chemistry. 2009;113:1133-1138.
- 19. Mohanraj R, Sivasankar S. Sweet potato (*Ipomoea batatas* L.) a valuable medicinal food: A Review. Journal of Medicine and Food. 2014;17:733-741.
- López-Yerena A, Jaime-Rodríguez C, González-Coria J, Lamuela-Raventós RM, Vallverdú-Queralt A, Romanyà J, Pérez M. Sweet potato is not simply an abundant food crop: A comprehensive review of its phytochemical constituents, biological activities, and the effects of processing. Antioxidants. 2022;11:1648.
- Padmaja G. Uses and nutritional data of Sweet potato. In the Sweet potato (Loebenstein G, Thottapilly G, editors, Springer, Dordrecht, Netherlands. 2009;189-233.
- 22. Waramboi JG, Gidley MJ, Sopade PA. Influence of extrusion on expansion,

functional and digestibility properties of whole Sweet potato flour. Food Science and Technology. 2014;59:1136-1145.

- Karan YB, Şanli OG. The assessment of yield and quality traits of sweet potato (*Ipomoea batatas* L.) genotypes in Middle Black Sea region, Turkey. PLoS ONE. 2021;16(9):e0257703.
- 24. Wang S, Nie S, Zhu F. Chemical constituents and health effects of sweet potato. Food Research International. 2016;89:90-116.
- 25. Zhu F. Interactions between starch and phenolic compound. Trends in Food Science and Technology. 2015;43:129-143.
- 26. Truong V, Avula RY. Sweet potato purees and powders for functional food ingredients. Sweet potato: postharvest aspects in food. 2010:117-61.
- Kiribuchi T, Kubota K. Studies on cooking of Sweet potato (part 1) changes in sugar content and beta-amylase activity during cooking. Kaseigaku Zassi. 1976;27(6):418-22.
- Ishida H, Suzuno H, Sugiyama N, Innami S, Tadokoro T, Maekawa A. Nutritive evaluation on chemical components of leaves, stalks and stems of sweet potatoes (*Ipomoea batatas* poir). Food Chemistry. 2000;68(3):359-67.
- 29. Sanoussi AF, Dansi A, Ahissou H, Adebowale A, Sanni LO, Orobiyi A, Dansi M, Azokpota P, Sanni A. Possibilities of sweet potato (*Ipomoea batatas* L.) value chain upgrading as revealed by physicochemical composition of ten elites landraces of Benin. African Journal of Biotechnology. 2016;15(13)481-9.
- Oloo BO, Shitandi A, Mahungu S, Malinga JB, Ogata BR. Effects of lactic acid fermentation on the retention of beta carotene content in orange fleshed sweet potatoes. International Journal of Food Studies. 2014;3:13-33.
- 31. Tang Y, Cai W, Xu B. Profiles of phenolics, carotenoids and antioxidative capacities of thermal processed white, yellow, orange and purple sweet potatoes grown in Guilin, China. Food Science and Human Wellness. 2015;4:123-132.
- 32. Ji H, Zhang H, Li H, Li Y. Analysis on the nutrition composition and antioxidant activity of different types of sweet potato cultivars. Food and Nutrition Science. 2015;6:161.

- Florence TM. The role of free radicals in disease. Australian and New Zealand Journal of Agriculture, Food Chemistry and Ophthalmology. 1995;23:3-7.
- 34. Chen YC, Chang HS, Lai HM, Jeng ST. Characterization of the wound inducible protein IPOMOELIN from sweet potato. Plant Cell Environment. 2005;28:251-259.
- 35. Chen CM, Li SC, Liao YY, Liu JF, Shyur JF, Chen PR, Chen CY. *In vitro* immunomodulatory effect of *Ipomoea batatas* L. tuber bioactives. The FASEB Journal. 2013:862-869.
- Sugata M, Lin CY, Shih YC. Antiinflammatory and anticancer activities of Taiwanese purple-fleshed sweet potatoes (*Ipomoea batatas* L. Lam) extracts. BioMed Research International. 2015;2015:10. Article ID 768093
- 37. Wang LS, Stoner GD. Anthocyanins and their role in cancer prevention. Cancer Letters. 2008;269:281-290.
- Ludvik B, Hanefeld M, Pacini G. Improved metabolic control by *Ipomoea batatas* (Caiapo) is associated with increased adiponectin and decreased fibrinogen levels in type 2 diabetic subjects. Diabetes Obesity and Metabolism. 2008;10:586-592.
- Ayeleso TB, Ramachela K, Mukwevho E. A review of therapeutic potentials of sweet potato: Pharmacological activities and influence of the cultivar. Tropical Journal of Pharmaceutical Research. 2016;15(12):2751-2761.
- 40. Adsul VB, Khatiwora E, Torane R, Deshpande NR. Antimicrobial activities of *Ipomoea carnea* leaves. Journal of Natural Product and Plant Resources. 2012;2(5):597-600.
- 41. Promprom W, Chatan W. GC-MS Analysis and antioxidant activity of Bauhinia nakhonphanomensis leaf ethanolic extract. Pharmacognosy Journal. 2017;9(5).
- 42. Nagai M, Tani M, Kishimoto Y, Iizuka M, Saita E, Toyozaki M, Kondo K. Sweet potato (*Ipomoea batatas* L.) leaves suppressed oxidation of low density lipoprotein (LDL) *in vitro* and in human subjects. Journal of Clinical Biochemistry and Nutrition. 2011;48:203.
- 43. Mazza G. Anthocyanins and heart health. Ann I Super Sanita. 2007;43:369.
- 44. Panda V, Sonkamble M. Anti-ulcer activity of *Ipomoea batatas* L. tubers (sweet potato). Functional Foods in Health and Disease. 2012;2:48-61.

- Sathish R, Rani M, Natarajan K. Study of ulcer protective effect of *Ipomoea batatas* L. dietary tuberous roots (Sweet Potato). Iranian Journal of Pharmacology and Therapeutics Collection. 2012;11:36-39.
- 46. Osime EO, Ediale GE, Omoti CE, Famodu AA. Effect of sweet potato leaf (*Ipomoea batatas* L.) extract on some haematological parameters using rabbits. Journal of Medicine and Biomedical Research. 2008;7(68):12-15.
- Montejo JF, Mondonedo JAB, Lee MGA, Ples MB, Vitor RJS. Hematological effects of *Ipomoea batatas* (camote) and *Phyllanthus niruri* (sampa-sampalukan) from philippines in the ICR mice (*Mus musculus*). Asian Pacific Journal of Tropical Biomedicine. 2015;5:29-33.
- 48. Islam S. Nutritional and medicinal qualities of Sweet potato tops and leaves. Plant Science, In the Cooperative Extension Program, University of Arkansas at Pine Bluff. Arkansas, FSA6135; 2014.
- 49. Etela I, Kalio GA. Yields components and 48-h rumen dry matter degradation of three sweet potato varieties in Ndama steers as influenced by date of harvesting. Journal of Agriculture and Social Research, 2011;11(2):15-21.
- 50. Ali AIM, Wassie SE, Joergensen RG, Korir D, Goopy JP, Butterbach-Bahl K, Merbold L, Dickhoefer U, Schlecht E. Feed quality and feeding level effects on faecal composition in East African cattle farming systems. Animals, 2021;11:564.
- Mibach M, C.F. Demarco, A. A. Barbosa, L. De Oliveira, M. N. Corrêa, F.A.B.D. Pino, V.R. Rabassa E, Schmitt GV, Kozloski CC, Brauner sweet potato flour as a replacement for ground corn as an energetic concentrate. Animal Production. Ciência Rural, Santa Maria. 2021; 51(10):e20200838. ISSNe: 1678-4596.
- 52. Murugan S, Paramasivam SK, Nedunchezhiyan M. Sweet potato as animal feed and fodder. Global Science Books, Fruit, Vegetable and Cereal Science and Biotechnology. 2012;6(1): 106-114.
- Beckford RC, Bartlett JR. Inclusion levels of sweet potato root meal in the diet of broilers I. Effect on performance, organ weights, and carcass quality. Poultry Science. 2015;94(6):1316-1322.
- 54. Mozafari O, Ghazi S, Moeini MM. The effects of different levels of edible potato

(*Solanum tubresum*) replacing maize on performance, serum metabolite and immune system of broiler chicks. Iran Journal of Animal Science. 2013;3(3):583-588.

- Okereke CO, Oti E, Nwauzor EC. Effect of dietary inclusion of orange-fleshed sweet potato meal on egg quality of laying hens. In the Proceedings of 14th Annual Conference of Animal Science Association of Nigeria, Nigeria. 2009:382-384.
- 56. Edache JA, Musa U, Ehizokale MUM, Esilonu JO, Okpala EJ, Karsin PD, Yisa A, Zwandor NJ. Replacement value of sweet potato (*Ipomea Batatas* L.) Meal for maize in practical diets fed to quail (*Cotumix cotumix* japonica) chicks. Nigeria Journal of Animal Production, 2009;36(11):34-40.
- 57. Tsega W, Tamir B. The effect of increasing levels of dried leaves of sweet potato (*Ipomoea batatas*) on dry matter intake and body weight gain performance of broiler finisher chickens. Livestock Research for Rural Development. 2009;21. Article #208.
- 58. Heuzé V, Tran G, Hassoun P. Sweet potato (*Ipomoea batatas* L.) forage. Feedipedia.org; 2013.
- Mosebework K, Negesse T, Nurfeta A. Effect of replacing maize (*Zea Mays* L.) with orange fleshed sweet potato tuber (*Ipomoea Batatas* L.) on performance of cobb 500 broiler chickens. Agricultural Research & Technology: Open Access Journal. 2018;13(5):555-895.
- 60. Tagesse T. Review on feed resources with more emphasis on use of cassava and sweet potato meal in chicken diet: Ethiopia. Animal and Veterinary Sciences. 2021;9(2):32-38.
- Scott GJ, Maldonado L. Sweet potato for the New Millennium: Trends in production and utilization in developing countries. CIP Program Repository. 1998;98:329-335.
- Amagloh FC, Yada B, Tumuhimbise GA, Amagloh FK, Kaaya AN. The potential of Sweet potato as a functional food in Sub-Saharan Africa and its implications for health: A review. Molecules, 2021;26:2971.
- 63. Bach D, Bedin AC, Lacerda LG, Nogueira A, Demiate IM. Sweet potato (*Ipomoea batatas* L.): A versatile raw material for the food industry. Review-Food/Feed Science and Technology, Brazilian Archives of Biology and Technology. 2021;64:1678-4324.

- 64. Rozi F, Prasetiaswati N, Elisabeth DAA. Study on sweet potato market behavior in supporting food security. In the IOP Conference Series, Earth and Environmental Science, 2021;756:012082.
- 65. Low JW, Arimond M, Osman N, Cunguara B, Zano F, Tschirley D. A food-based approach introducing orange-fleshed Sweet potato increased vitamin A intake and serum retinol concentrations in young children in rural Mozambique. Journal of Nutrition, 2007;137:1320-1327.
- Low J, Lynam J, Lemaga B, Crissman C, Barker I, Thiele G, Namanda S, Wheatley C, Andrade M. Chapter 16, Sweet potato in Sub-Saharan Africa. Loebenstein G, Thollappilly G. (eds.), the Sweet potato. © Springer Science-Business Media B.Y. 2009:359.
- 67. Mu TH, Sun HN, Ma MM. Sweet potato snack foods. Chemistry, Processing and Nutrition. 2019:303-324.
- 68. Dereje B, Girma A, Mamo D, Chalchisa T. Functional properties of sweet potato flour and its role in product development: A review. International Journal of Food Properties. 2020;23(1):1639-1662.
- 69. Okoye AC. Gender deferential in market competitiveness among Sweet potato smallholder farmers in South Eastern Nigeria. A Ph.D. dissertation presented in the Department of Agribusiness and Management, Michael Okpara University of Agriculture, Umudike, Nigeria. 2018:10-65.
- Giri NA, Ambedkar B, Sakhale BK. Sweet potato (*Ipomoea batatas* L.): A valuable tropical tuber crop: A review. The Pharmaceutical Innovation Journal. 2019;8(6):182-191.
- 71. Wijaya H, Slay A. Abdullah N. Ice cream products made from processed purple sweet potatoes: A product organoleptic study. In the IOP Conference Series, Earth and Environmental Science. 2021;807:042074.
- 72. Affandi DR, Handajani S. Es krim ubi jalar ungu (*Ipomoea batatas* L.): Tinjauan sifat sensoris, fisik, kimia. Dan Aktivitas Antioksidannya J. Teknol. Has. Pertan. 2011:4.
- 73. Anugrah RM, Suryani E. Kandungan Gizi Donat dengan Penambahan Ubi Ungu (*Ipomoea batatas* L.). Sebagai Makanan Jajanan Berbasis Pangan Lokal Bagi Anak Sekolah J. Gizi. 2020;9:150-158.

- 74. Lirag TBM. Determinants of profitability of sweet potato production in Camarines Sur, Philippines, International Journal of Advanced Science, Engineering, Information and Technology. 2019;9:467-472.
- 75. Virgínioe Silva JO, Almeida MF, Da Conceição Alvim-Ferraz M, Dias JM. Integrated production of biodiesel and bioethanol from sweet potato. Renew, Energy. 2018;124:114-120.
- 76. Mudege NN, Mayanja S, Muzhingi T. Women and men farmer perceptions of economic and health benefits of orange fleshed sweet potato (OFSP) in Phalombe and Chikwawa districts in Malawi. Food Security. 2017;9:387-400.
- 77. Neela S, Fanta SW. Review on nutritional composition of orange-fleshed sweet potato and its role in management of vitamin A deficiency. Food Science and Nutrition, 2019;7:1920-1945.

- Laurie S, Faber M, Adebola P, Belete A. Biofortification of sweet potato for food and nutrition security in South Africa. Food Research International. 2015;76:962-970.
- 79. FAO. Food and Agriculture Organization, Land and Water; 2020.
- UNIDO. United Nations Industrial Development Organization, Baseline Report of Clean Cooking Fuels in the East African Community (EAC) Region. 2020; 2-34.
- Sakai P, Afionis S, Favretto N, Stringer LC, Ward C, Sakai M, Weirich Neto PH, Rocha CH, Alberti Gomes J, De Souza NM, et al. Understanding the implications of alternative bioenergy crops to support smallholder farmers in Brazil. Sustainability. 2020;12:2146.
- 82. Su MH, Huang CH, Li WT, Tso CT, Lur HS. Water footprint analysis of bioethanol energy crops in Taiwan. Journal of Cleaner Production. 2015;88:132-138.

© 2022 Kojo and Fosu; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history: The peer review history for this paper can be accessed here: https://www.sdiarticle5.com/review-history/94710