



## **Effect of Zinc Fortification on Growth and Yield of Finger Millet (*Eleusine coracana* L.)**

**P. Rathnakar<sup>a\*</sup>, N. Mohan<sup>a#</sup>, Boyina Aparna<sup>a#</sup> and Pasala Ramya<sup>a</sup>**

<sup>a</sup> Department of Agronomy, NAI, SHUATS, Prayagraj, U. P, India.

### **Authors' contributions**

*This work was carried out in collaboration among all authors. 'I' authors read and approved the final manuscript.*

### **Article Information**

DOI: 10.9734/IJPSS/2022/v34i2031129

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/87751>

**Original Research Article**

**Received 20 March 2022**

**Accepted 30 May 2022**

**Published 07 June 2022**

### **ABSTRACT**

A field experiment was conducted during *kharif* 2021 at CRF (Crop Research Farm), Department of Agronomy, SHUATS, Prayagraj, (U.P). The soil of the experimental field is sandy loam in texture with a neutral soil pH. The experiment was laid out in a Randomized Block Design with nine treatments and three replications. The treatments consisted of three levels of Zinc *viz.*, 10 kg/ha, 15 kg/ha, 25 kg/ha and three levels of Zinc sulphate at 0.1%, 0.3% and 0.5%, whose effect was observed in finger millet. The results revealed that the treatment with application of 15 kg/ha zinc + 0.5% Znso<sub>4</sub> recorded higher plant height (81.22 cm), number of tillers/plant (7.85/plant), plant dry weight (26.32g/plant), and the yield attributes *viz.*, seed yield (2.79 t/ha), straw yield (4.72 t/ha) and harvest index (37.19%) were also recorded highest in the treatment with application of 15 kg/ha zinc and 0.5% Znso<sub>4</sub>.

**Keywords:** Zinc; zinc sulphate; growth parameters; yield attributes and finger millet.

### **1. INTRODUCTION**

Finger millet (*Eleusine coracana* L.) is an important small millet crop grown in India, and it

is distinguished by having the highest productivity among millets. "In India finger millet is cultivated over an area of 1.19 million hectares with a production of 1.98 million tones giving an

<sup>o</sup> M.Sc. Scholar,

<sup>#</sup> Assistant Professor,

\*Corresponding author: E-mail: putlurathnakar756@gmail.com;

average productivity of 1661 kg per hectare” (<http://www.indiastst.com>). “Small-seeded grains belonging to different variety of annual grasses that are cultivated primarily as grain crops on marginal lands in dry areas in temperate, subtropical and tropical regions are collectively referred as millets. They are the most important cereals of semi-arid zones of the world and are staple food for millions of people in Africa and Asia” [1]. It is called finger millet, because the inflorescence resembles the fingers of a human hand. It is an important crop in drought-prone regions because of its outstanding ability to withstand adverse weather conditions and grow in marginal and poor soils.

Finger millet is locally known as ragi and mandua (India). “Ragi is commonly known as “Nutritious millet” as the grain is nutritionally superior to many cereals (rice, corn and sorghum) providing proteins, minerals, iron, calcium and vitamins in abundance. When consumed as food, it provides a sustaining diet, especially for people doing hard work. Straw makes valuable fodder for both draught and milch animals. Finger millet is considered as a wholesome food for diabetic patients. Grain may also be malted and flour of the malted grain is used as cakes or porridge and a nourishing food for infants and invalids” [2].

“Zinc - Micronutrients are essential for plant growth and play an important role in balanced crop nutrition. Micronutrients are as important to plant nutrition as primary and secondary nutrients, though plants do not require as much of them. They play a major role in plant growth like protein synthesis, improving seed quality, cell division and pollen tube growth” [3]. “Deficiency of micronutrients during the last three decades as grown in both magnitude and extent because of increased use of high analysis fertilizers, use of high yielding crop varieties and increasing in cropping intensity. The deficiency of Zn, B and Fe are 49%, 33% and 12% respectively in Indian soils. Among the seven micronutrient elements (Fe, Mn, Cu, Zn, B, Mo, Ca) essential for plant growth, Zn has assumed extensively important place in Indian agriculture” [3].

“Zinc plays a key role in plants with enzymes and proteins involved in carbohydrate metabolism, protein synthesis, gene expression, auxin metabolism, pollen formation, maintenance of biological membranes, protection against photo-oxidative damage and

heat stress, and resistance to infection by pathogens” (Alloway, 2008).

Zinc deficiency in plants retards photosynthesis and nitrogen metabolism and reduces flowering. This results in delayed maturity, a decreased yield, and poor quality of finger millet.

“Zinc hampers the productivity of cereals, millets and oil seed crops. In addition to the nutritional value of zinc, it is a component of various enzyme systems. It also plays a vital role in biosynthesis of Indole Acetic Acid. It helps in formation of nucleic acids and the synthesis of proteins” [4].

“Finger millet requires considerable amount of zinc as well as calcium for its growth and development. Deficiency of secondary and micronutrients leads to reduction in the number of effective tillers and improper grain filling” [5]. Finger millet is being increasingly incorporated in breakfast cereals, beverages and infant foods, which makes it an important crop that deserves attention for fortification of its grains with zinc [6-8]. Therefore, the present study was undertaken to study the effect of zinc fortification on growth and yield of finger millet.

## 2. MATERIALS AND METHOD

The experiment was conducted during Kharif season 2021. The experiment was laid out in Randomized Block Design consisting of nine treatments each replicated thrice. The treatment combinations are T1: Zinc 10kg/ha+ZnSo<sub>4</sub> 0.1% at 30DAS, T2: Zinc 10kg/ha+ Znso<sub>4</sub> 0.3% at 30DAS, T3: Zinc 10kg/ha+ Znso<sub>4</sub> 0.5% at 30DAS, T4: Zinc 15kg/ha + Znso<sub>4</sub> 0.1% at 30DAS, T5: Zinc 15kg/ha+ Znso<sub>4</sub> 0.3% at 30DAS, T6: Zinc 15kg/ha+ Znso<sub>4</sub> 0.5% at 30DAS, T7: Zinc 25kg/ha+ Znso<sub>4</sub> 0.1% at 30DAS, T8: Zinc 25kg/ha+ Znso<sub>4</sub> 0.3% at 30DAS, T9: Zinc 25kg/ha+ Znso<sub>4</sub> 0.5% at 30DAS. In each plot five plants were chosen and tagged randomly for recording observations at 20, 40, 60, 80, 100 days after sowing of the crop. Later on statistical analysis was performed using the method of analysis of variance with 5% level of significance for the F-Test.

## 3. RESULTS AND DISCUSSION

**Growth attributes:** The data pertaining to growth attributes given in the Table 1.

**Plant Height:** At harvest, the highest plant

height (81.22 cm) was recorded with application of zinc 15 kg/ha + zinc sulphate 0.5% which was significantly superior over rest of the treatments except the treatment with application zinc 10 kg/ha + zinc sulphate 0.5%. The increased plant height might be due to profound influence of Zinc fertilizers on height of plant as increased metabolic process in plant which has prompted meristematic activities causing higher apical growth and photosynthetic area.

**Number of tillers:** Maximum number of tillers (7.85/plant) was recorded at harvest with application of zinc 15 kg/ha + zinc sulphate 0.5% which was superior over the other treatments and the treatment with zinc 10 kg/ha + zinc sulphate 0.3%, zinc 15 kg/ha + zinc sulphate at 0.3% was statistically at par with treatment 6.

**Dry weight (g/plant):** At harvest, highest plant dry weight (26.32 g/plant) was recorded with application of zinc 15 kg/ha + zinc sulphate 0.5% which was significantly superior over rest of all the treatments except with application of zinc 15 kg/ha + zinc sulphate 0.3% and zinc 10 kg/ha + zinc sulphate 0.5%. "It might be due to the increase in plant dry weight might be attributed to the optimum and uniform availability of micronutrients in the entire growth period by means of application of nutrients through the soil. There was increased dry weight at successive stages of growth, with the Zinc and iron application" Pradhan et al. [9]. "The improved dry matter production and nutrient uptake in finger millet with foliar application of Zinc sulphate at 0.5%" was reported by Ajay Kumar et al. [10].

**Crop growth rate and relative growth rate:** Maximum crop growth rate (18.36 g/m<sup>2</sup>/day)

was recorded at 80 DAS to harvest with the application of zinc at 10 kg/ha + zinc sulphate 0.5%. was superior over the rest of the treatments and statistically on par with the zinc 15 kg/ha + zinc sulphate 0.1%, zinc 15 kg/ha + zinc sulphate 0.3% and zinc 15 kg/ha + zinc sulphate 0.5%.

At 80 DAS to harvest, maximum relative growth rate (0.0195 g/m<sup>2</sup>/day) was recorded with the application of zinc at 15 kg/ha + zinc sulphate at 0.3%, which was superior over the rest of the treatments and there is no statistically significant difference between them.

**Yield Attributes:** The data pertaining to yield attributes given in the Table 2.

At harvest the Superior test weight was recorded with application of Zinc 15 kg/ha + Zinc sulphate 0.5% (3.79 g) and there was no significant difference between the treatment combinations.

**Seed Yield:** Maximum seed yield was obtained with application of zinc 15 kg/ha + zinc sulphate 0.5% (2.79 t/ha), which was significantly superior over rest of all the treatments except with application of zinc 10 kg/ha + zinc sulphate 0.5% and the treatment with zinc 15 kg/ha + zinc sulphate 0.3%.

"The increase in seed yield and straw yield when zinc were sprayed on foliage at tillering stage in finger millet. Foliar application of zinc plays a critical role in crop growth, involving in photosynthesis processes, respiration and other biochemical and physiological activities and thus their importance in achieving higher yields" reported by Zeidan et al. [11].

**Table 1. Effect of Zinc fortification on growth attributes of finger millet at harvest**

Treatments Combinations	Plant height	No. of Tillers/ plant	Dry Weight	CGR	RGR
1. Zinc 10 kg/ha + Zinc sulphate 0.1%	77.34	6.37	24.48	16.34	0.0178
2: Zinc 10 kg/ha + Zinc sulphate 0.3%	78.52	6.92	23.59	15.12	0.0170
3: Zinc 10 kg/ha + Zinc sulphate 0.5%	80.34	7.64	25.86	18.36	0.0191
4: Zinc 15 kg/ha + Zinc sulphate 0.1%	78.51	6.47	24.78	17.68	0.0193
5: Zinc 15 kg/ha + Zinc sulphate 0.3%	79.20	7.44	25.43	18.24	0.0195
6: Zinc 15 kg/ha + Zinc sulphate 0.5%	81.22	7.85	26.32	18.25	0.0186
7: Zinc 25 kg/ha + Zinc sulphate 0.1%	78.70	6.45	24.93	17.30	0.0187
8: Zinc 25 kg/ha + Zinc sulphate 0.3%	78.38	6.53	23.61	15.04	0.0168
9: Zinc 25 kg/ha + Zinc sulphate 0.5%	76.92	6.06	22.93	13.70	0.0156
<b>SEm(±)</b>	0.62	0.18	0.43	0.91	0.00
<b>CD (P=0.05)</b>	1.87	0.53	1.30	2.72	-

**Table 2. Effect of zinc fortification on yield and yield attributes of finger millet**

Treatments	Test weight (gm)	Seed yield (t/ha)	Straw yield (t/ha)	Harvest Index (%)
1. Zinc 10 kg/ha + Zinc sulphate 0.1%	3.51	2.48	4.48	35.64
2: Zinc 10 kg/ha + Zinc sulphate 0.3%	3.57	2.59	4.58	36.09
3: Zinc 10 kg/ha + Zinc sulphate 0.5%	3.67	2.67	4.67	36.37
4: Zinc 15 kg/ha + Zinc sulphate 0.1%	3.41	2.51	4.52	35.67
5: Zinc 15 kg/ha + Zinc sulphate 0.3%	3.57	2.63	4.57	36.57
6: Zinc 15 kg/ha + Zinc sulphate 0.5%	3.79	2.79	4.72	37.19
7: Zinc 25 kg/ha + Zinc sulphate 0.1%	3.48	2.43	4.41	35.56
8: Zinc 25 kg/ha + Zinc sulphate 0.3%	3.53	2.36	4.39	34.99
9: Zinc 25 kg/ha + Zinc sulphate 0.5%	3.42	2.29	4.31	34.65
<b>SEm (±)</b>	0.08	0.04	0.06	0.47
<b>CD (P=0.05)</b>	-	0.13	0.19	1.42

**Straw Yield:** At harvest significantly higher straw yield was obtained with application of zinc 15 kg/ha + zinc sulphate 0.5% (4.72 t/ha), However the treatments with zinc 10 kg/ha + zinc sulphate 0.5%, zinc 10 kg/ha + zinc sulphate 0.3% and zinc 15 kg/ha + zinc sulphate 0.3% are statistically at par with treatment 6. "This might be due to profound influence of zinc fertilizers on growth attributes as increased metabolic process in plants which has promoted meristematic activities and photosynthetic process, ultimately better growth resulted in higher yield and yield attributes" [12].

**Harvest Index (%):** Maximum harvest index was obtained with application of zinc 15 kg/ha + zinc sulphate 0.5% (37.19 %), which was significantly superior over rest of all the treatments except with application of zinc 15 kg/ha + zinc sulphate 0.3%, zinc 10 kg/ha + zinc sulphate 0.5% and zinc 10 kg/ha + zinc sulphate 0.3%.

The increase in yield and yield attributes might be due to foliar application of zinc sulphate might be due to increase seed weight. Sai Divya et al. [3].

The application of zinc in a soil deficient in zinc improved the overall growth and development of plants and ultimately the grain and straw yields was increased. These findings are also supported by Shrivatsava et al. and Sammauria 2007. The improvement in yield attributes is the manifestation of better growth, higher photosynthetic activity and transport of photosynthates from source to sink. The higher values of yield parameters might be due to increased availability of Zinc application.

#### 4. CONCLUSION

On the basis of the present experiment on finger millet crop in kharif season, it is concluded that the treatment T6 with application of zinc at 15 kg/ha and 0.5% zinc sulphate recorded higher plant height, dry weight, number of tillers, seed yield and straw yield respectively. Therefore, it is recommended to the farmers to receive higher yield of finger millet.

#### ACKNOWLEDGEMENT

The authors are thankful to Dr. Shikha Singh Assistant Professor, Department of Agronomy, SHUATS, Prayagraj, and U.P. for constant support and guidance to carry out the experiment. I must also express my gratitude to my friends for their constant support throughout my research work.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Thippeswamy TG, Junna L, Shinde M. Proximate composition, resistant starch and other phytochemical constituents of native finger millet cultivar. International journal of food and nutritional science. 2016;3(5):2320-7876.

2. Radhika Chaturvedi, Sarita Srivatsava. Genotype variations in physical, nutritional and sensory quality of popped grains of amber and dark genotypes of finger millet. *Journal of Food Science and Technology Mysore*. 2008;45(5):443-446.
3. Sai Divya B, Rajesh Singh and Wasim Khan. Effect of foliar application of iron and zinc on yield and economics of finger miller (*Eleusine coracana L.*). *The Pharma Innovation Journal*. 2021;10(4): 897-899.
4. Dholariya HP, Zinzala VJ, Patel JV, Patel VM. Zinc nutrition in finger millet (*Elusine coracana L. Gaertn.*) for better nutritional security, *International Journal of Current Microbiology and Applied Sciences*. 2020;11:1082-1086.
5. Shetty YV, Sheshadri T, Vasuki N, Gajanan GN. Effect of organic and inorganic nutrient sources on yield of ragi and groundnut under rainfed conditions. *Annual Report*. 1993;40:45-46.
6. Bhumika T, Kalpana P. Finger millet flour as vehicle for fortification with Zinc. *Journal of Trace Elements in Medicine and Biology*. 2010;24(1): 46-51.
7. Cakmak I. Enrichment of cereal grains with Zinc agronomic or genetic bio fortification. *Plant and Soil*. 2008;302(1-2): 1-17.
8. Sammauria R. Response of fenugreek to phosphorous and zinc application and their residual effect on succeeding pearl millet. (*Pennisetum glaucum*) under irrigated conditions of North West Rajasthan. Ph.D. Thesis, Rajasthan Agricultural University. Bikaner; 2007.
9. Pradhan A, Abhinav S, Nag SK, Chandrakar TP. Effect of zinc fertilization on growth and yield of finger millet (*Eleusine coracana L. Gaertn.*). *International Journal of Science Environment and Technolog*. 2016;5(3): 1477-1487.
10. Ajay Kumar, E., Surekha K., Bhanu Rekha K. and Harish Kumar Sharma S. Effect of Various sources of Zinc and Iron on Grain yield, Nutrient uptake and Quality parameters of Finger millet (*Elusine corocona L.*). *International Research Journal of Pure and Applied Chemistry*. 2020;21(2):46-55.
11. Zeidan E, Satyanarayanswamy Y, Triveni C. Effect of micronutrient application with different sources of application on growth and yield of finger millet crop in laterite zone. *Journal of Agricultural Science and Technology*. 2010;4:403-416.
12. Saraswathi, Shetty YV, Diesh Kumar M, Gurumurthy KT. Effect of Nano Zno on growth and yield of finger millet. *International journal of current microbiology and applied sciences*. 2019; 8(2):1365-1371.

© 2022 Rathnakar et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/87751>