

CrossMark
click for updates

Effect of Early Physical Activity Programs on Motor Performance and Neuromuscular Development in Infants Born Preterm: A Randomized Clinical Trial

Leila Valizadeh¹, Mahnaz Sanaeefar^{1*}, Mohammad Bager Hosseini², Mohammad Asgari Jafarabadi³, Aryan Shamili⁴

¹Department of Pediatric Nursing, Nursing and Midwifery Faculty, Tabriz University of Medical Sciences, Tabriz, Iran

²Department of Pediatrics, Medicine Faculty, Tabriz University of Medical Sciences, Tabriz, Iran

³Department of Statistics and Epidemiology, Faculty of Health, Tabriz University of Medical Science, Tabriz, Iran

⁴Department of Occupational Therapy, Faculty of Rehabilitation, Iran University of Medical Sciences, Tehran, Iran

ARTICLE INFO

Article Type:

Original Article

Article History:

Received: 21 Jan. 2016

Accepted: 19 Dec. 2016

ePublished: 1 Mar. 2017

Keywords:

Preterm

Infant

Low birth weight

Physical activity

Motor performance

Neuromuscular development

ABSTRACT

Introduction: Although the survival rate of infants born preterm has increased, the prevalence of developmental problems and motor disorders among this population of infants remains the same. This study investigated the effect of physical activity programs in and out of water on motor performance and neuromuscular development of infants born preterm and had induced immobility by mechanical ventilation.

Methods: This study was carried out in Al-Zahra hospital, Tabriz. 76 premature infants were randomly assigned into four groups. One group received daily passive range of motion to all extremities based on the Moyer-Mileur protocol. Hydrotherapy group received exercises for shoulders and pelvic area in water every other day. A combination group received physical activity programs in and out of water on alternating days. Infants in a containment group were held in a fetal position. Duration of study was two weeks 'from 32 through 33 weeks post menstrual age (PMA). Motor outcomes were measured by the Test of Infant Motor Performance. Neuromuscular developmental was assessed by New Ballard scale and leg recoil and Ankle dorsiflexion items from Dubowitz scale. Data were analyzed using SPSS version 13.

Results: TIMP and neuromuscular scores improved in all groups. Motor performance did not differ between groups at 34 weeks PMA. Postural tone of leg recoil was significantly higher in physical activity groups post intervention.

Conclusion: Physical activities and containment didn't have different effects on motor performance in infants born preterm. Leg recoil of neuromuscular development items was affected by physical activity programs.

Please cite this paper as: Valizadeh L, Sanaeefar M, Hosseini MB, Jafarabadi MA, Shamili A. Early physical activity programs on motor performance and neuromuscular development in infants born preterm: a randomized clinical trial. J Caring Sci 2016; 6 (1): 69-81. doi:10.15171/jcs.2016.008.

Introduction

The improvements in prenatal and postpartum care coupled with advances made in medical technology have increased the survival rate of extremely preterm and low birth weight infants.^{1,2} Despite the innovative interventions and improved survival rates for immature infants,³ the incidence of developmental and motor disorders among these infants has remained almost unchanged.^{4,5} Secondary concerns are about increased rates of

developmental disability and the declining quality of life in preterm infants.⁶ Motor developmental disorders as well as hearing and sight disorders are associated with preterm birth and very low birth weight. It has been observed in one out of every three infants born premature.⁴ Complex biological, medical, and environmental elements contribute to developmental problems in infants born preterm.⁷⁻¹² Small for gestational age premature infants are at a higher risk for developmental and cognitive delays. Those

*Corresponding Author: Mahnaz Sanaeefar, (MSc), email: Mahnaz.Sanaeefar@gmail.com. This study was approved and funded by the deputy of research of Tabriz University of Medical Sciences (Project number: 466).



© 2017 The Author(s). This work is published by Journal of Caring Sciences as an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by-nc/4.0/>). Non-commercial uses of the work are permitted, provided the original work is properly cited.

born at extremely low birth weights are at a double risk and require special clinical attention and care.¹³

There has been some research indicating that early motor intervention can facilitate motor development and minimize the harmful effects of the NICU environment.⁷ The interventions in this area might have a significant impact on differentiation of muscle fibers and consequently on the development of postural tone in infants born preterm.^{14,15}

As for the effectiveness of early developmental interventions on preterm infants' motor development at infant, pre-school, and school ages as well as in adulthood, some studies revealed no significant differences among the infants of 0-3 years of age with respect to the motor outcomes. Nor were there any differences in motor outcomes between infants in the intervention and control groups who weighed more or less than 2000 grams. This review demonstrated that the effect size of intervention on motor development is relatively small; however, there was a statistically significant difference at infancy.

Programs that began in hospital were found to have a slightly greater impact on motor outcomes at infancy than did the ones that were initiated following hospital discharge. The programs that focused on infant development alone had a slightly greater impact on motor outcomes at infancy than those that simultaneously focused on the parent-infant relationship as well as infant development; however, the effect in neither group was significant.^{16,17}

A meta-analysis study found that the children born LBW or preterm who participated in early intervention programs exhibited significant improvements in their mental or neuromusculoskeletal and movement functional developments.¹⁸ There is some evidence suggesting that physical activity can be useful in decreasing the risk of osteopenia in infants born preterm by promoting bone mass and weight gain.¹⁹⁻²¹

Some benefits of hydro-kinesiotherapy for accelerating the growth and development of

biological systems are found in premature infants hospitalized in NICUs. Sweeney found hydrotherapy to be useful for promoting weight gain and feeding tolerance.²² Vignochi indicated that aquatic physical therapy was effective in reducing pain and improving sleep quality. These authors suggested that hydrotherapy can be used as a non-pharmacological method for relieving pain and improving deep sleep quality and duration and can contribute to the reduction of the harmful effects of the NICU in infants, without depriving them of the tactile-kinesthetic stimulation necessary for their neurodevelopment.²³ In 1983, Sweeney compared exercises in water with immersion only and found normal vital signs during both procedures.²² Recently Tobinaga has shown that preterm newborns have a significant reduction in both heart rate and respiratory rate (both within normal limits), and that their salivary cortisol levels were significantly reduced after hydrotherapy, suggesting short-term relief from stress.²⁴ Based on this evidence, aquatic physical therapy might be introduced as a safe intervention for infants born preterm.

In view of the alarming prevalence of preterm birth in Iran almost 9.2%²⁵ it seems necessary that adequate planning should be made to improve the quality of life of the preterm infants. We are endeavoring to enhance our knowledge and understanding of the physical therapy programs for different purposes in infants born preterm. Even though there is no shortage of such programs, further studies are required to help us identify the effectiveness of interventions. As such, this study was designed and implemented as a step towards identifying appropriate interventions for premature infants.

Studies on effectiveness of physical activity and hydrotherapy programs in promoting motor development in this population of infants are limited.²⁶ Knowledge of what influences neurodevelopmental outcomes is the key to developing better strategies.¹²

The purpose of the present study was to assess the immediate (short-term) effects of

physical activity in and out of water versus a control group on motor performance and neuromuscular development of infants born preterm hospitalized in a NICU. This study hypothesized that physical activity in or out of water may have different effects on functional motor performance and neuromuscular development, i.e., postural tone, based on the type of intervention and the environment (e.g., diminished force of gravity in the water). This study aimed to compare the effects of different types of physical activities on motor performance and neuromuscular development in infants born preterm. Researchers hypothesized that physical activity in bed would have a larger effect on neuromuscular development and consequently improve functional motor performance than hydrotherapy would. Furthermore, we expected that a combination of both programs would be the most effective in improving both functional performance and postural tone.

Materials and methods

This study was a Randomized Clinical Trial. The setting for the study was the Neonatal Intensive Care Unit (NICU) in Al-Zahra hospital affiliated to Tabriz University of Medical sciences, Tabriz, Iran. The Study was conducted from August 2014 to September 2015, following the acquisition of approval from the ethics committee of Tabriz University of Medical Sciences (code: 9360) and registration in the Iran Registry of Clinical Trials (IRTC) (code: N7201405208315).

The inclusion criteria were gestational age (GA) at birth between 25 and 30 weeks and corrected age of 32 weeks PMA at study time, minimum birth weight of 800 grams and weight appropriate for GA, Apgar score of more than 7 at 5th minute after birth, the weight range of 1000 to 2000 grams at entry point to the study, no congenital anomalies, being subjected to mechanical ventilation and sedative and muscle relaxants drugs for at least 3 days, receiving no aggressive treatment at the time of the study, no maternal substance abuse, no septicemia, no evidence of

metabolic problems or grade 2 and above intraventricular hemorrhage (IVH), and no neurological problems or chromosomal disorders. The infants who had late diagnosis of congenital neurological anomalies, chromosomal disorders, cardiopulmonary or neuro-musculoskeletal diseases and other conditions that needed surgery or phototherapy at the time of study and who had been transferred to another center were excluded from the study.

For study groups and interventions were as follow:

I- The physical activity was based on the Moyer-Mileur protocol.¹⁹ Both extension and flexion were performed five times at the wrist, elbow, shoulder, ankle, knee, and hip joints. The sequence was performed slowly and in the cephalo-caudal direction with gentle pressure within the range of motion. To cause a minimum of stress for the patient, the movements began with one side (the right arm and leg) and then finished on the other side (the left arm and leg). A total of 14 sessions was provided within 14 days beginning at 32 weeks postmenstrual age (PMA).

II- The hydrotherapy program was implemented based on the methods used in Sweeney²² and Vignochi's²³ studies. The infant was placed in water at 99-101 °F (37.2-38.3°C) with the head, neck and pelvis supported. In the first few moments, we gave the baby an opportunity to adapt to the water environment. Then for 10 minutes rotational movements were applied to the pelvis while the head and the neck were held still and then the pelvis was held still to perform movements on the upper limbs. The program was performed every other day for over 14 days (7 hydrotherapy sessions overall) beginning at 32 weeks PMA.

III- Infants in combination group received the previously described hydrotherapy and physical activity programs on alternating days over 14 days. Overall, 7 sessions of hydrotherapy and 7 sessions of physical activity were provided.

IV- In containment group, infants were placed on one side in a fetal position with a hand (not

too soft, not too hard) on the neonate at the top of the head and over the trunk and hip area to flex the hips for 10 minutes daily for 14 days.

All treatments were performed 30 minutes before or an hour after being fed for 10 minutes daily. Infants put in the nest in side-lying position 10 minutes before and after interventions without any manipulation.

The use of standardized tests is necessary for adequate assessment of the outcomes of early intervention.²⁷ Among the specific tests used to assess the motor development of babies, the Test of Infant Motor Performance (TIMP) was developed for early identification of delayed functional motor development and to assess the efficacy of physical and occupational therapy in clinical practice. It considers the influences of the infant's neurological maturation, the environment, the force of gravity, and posture on motor development.²⁸⁻³⁰

The TIMP consists of 42 test items in 2 distinct sections: 13 observed items and 29 elicited items, which test the infant's functional postural and motor control. Each item has its own scale; the number of points varies from 0 to 6. A total raw score is summed from item scores and then z scores were calculated. Results of raw scores are classified as "average" (within -0.5 to $+1.0$ standard deviations (SD) of age mean), "low average" (-0.5 to -1.0 SD below age mean), "below average" (-1.0 to -2.0 SD below age mean), and "far below average" (more than -2.0 SD below age mean).³¹ The TIMP has been shown to be valid as an outcome measure in three clinical trials involving intervention in the NICU.³²⁻³⁴ Infants were assessed with the TIMP at 32 and 34 weeks PMA by a pediatric occupational therapist blind to group assignment. The evaluator studied the TIMP user's manual, watched the self-instructional program on the TIMP which has been shown to produce reliable scoring, and practiced application of the test with infants born full term and infants born preterm at a variety of postmenstrual ages (PMA) over a 2-month period.

Neuromuscular examination consisted of 6 limb tone patterns based on New Ballard

score³⁵ which was adapted for infants born extremely preterm and two items from the Dubowitz GA assessment included leg recoil and ankle dorsiflexion.³⁶ Each item was measured by using a standardized form. The infants were examined undressed on an open bed or in their cot in a warm, quiet room. The neuromuscular assessment was completed by a neonatologist blind to group assignment at 31/6 weeks PMA, a day before the interventions began, and at 35/1 weeks PMA, a day after the interventions ended.

Evaluation of TIMP scoring reliability was performed by an expert pediatric occupational therapist by making video recordings of the assessment of 20 infants, rescoring based on the videos and comparing the results of the observations, the reliability of the evaluator was acceptable to achieve an intra-rater correlation coefficient (ICC=0.8). The neuromuscular test was performed by neonatologists. Scoring of 10 infants was done simultaneously by two persons. The intra-rater correlation coefficient of the observations was acceptable (ICC = 0.8).

Based on the information from twenty infants born preterm, sample size was estimated according to the mean (standard deviation) of the neuromuscular development items and motor performance scores on a pilot study. G power software as well as assuming the confidence interval of 95%, the test's power to be 0.8 and the effect size to be 0.4, the total sample size required was 76 infants and at least 19 infants for each group. The infants were allocated to physical activity, hydrotherapy, combination, and containment groups through random blocks of sizes 4 and 8. The statistical analysis and randomization were done by an associate professor of statistics who was blinded and independent from the study. The same effect size was observed in another recent study.³³ Motor function and postural tone outcomes were considered primary, because either or both could have been affected by the interventions. 132 infants were screened for inclusion in the present study and 56 infants did not meet the inclusion criteria (Fig. 1). Thus 76 eligible




Fig1: Flowchart of the study

infants at 32 weeks PMA were recruited for the study. Informed parental consent was obtained for each infant. The infants were randomly assigned into one of the four groups and received their programs. Before the interventions and assessments began, the physician and the nurse in charge were asked for permission to proceed. The infants' heart rate and oxygen saturation level were monitored during the intervention, using a

portable oximeter. In order to ensure the security and safety of children, the interventions were done by a nurse who practiced the protocol of interventions under the supervision of a pediatric occupational therapist, with infants born preterm over a 2-month period and she was familiar with stress and stability behavioral cues according to the assessment of infants born preterm behavior (APIB).³⁷ The battery-operated heart rate and

blood oxygen saturation was monitored through using a pulse oximetry sets, model Nova Metrix 512 made in USA. When physiological instability occurred once during the program (i.e., heart rate below 100 or over 180 bpm and drop of SaO₂ to less than 88% twice during the intervention)³⁸ the intervention was modified (particular exercise paused for 15 s and resumed if parameters recovered) based on the infant's physiological and behavioral stress responses in order to prevent overloading of the sensory inputs. Moreover, when physiological instability lasted more than 12 s and occurred twice within 24 hours after intervention, the baby was deemed to be intolerant of intervention and he/she was excluded from the study.

During the assessment, both the examiners and the statistical analysis were blind about the type of intervention received by the infants. Two infants were excluded because of physiological instability.

Normality were assessed using Kolmogorov-Smirnov test. Mean and standard deviation, frequency and percentage, maximum- minimum were calculated for description of variables. The comparison of the groups in terms of qualitative variables was performed by chi-square test. Post-test group means for TIMP scores were analyzed, using One-way analysis of variance (ANOVA). The comparison of the groups in terms of delay percentage was performed through variance analysis. Analysis of covariance (ANCOVA) was used to adjust confounding variables.

Pearson test was used to determine the correlation between motor function and duration of hospital stay, days submitted to mechanical ventilation, the duration of treatment with sedative drugs and the number of surfactant received. Sign test was performed to compare the results for the variable 'neuromuscular development' obtained before and after the intervention. Chi-square test was used for post-comparisons neuromuscular difference between groups. The data were analyzed using SPSS version 13.

Results

Overall, the results for 76 infants were available for analysis. 59.2% of the participants were female. Average birth weight was 1036 (SD=196) grams. The most common cause of hospitalization in the intensive care unit was prematurity with respiratory distress syndrome (100%). There were no statistically significant differences in the demographic and clinical characteristics between the groups ($P>0.05$). All the groups were similar, regarding gender, days with mechanical ventilation, and Apgar score at 5 minutes after birth. All of the infants were at 32 weeks PMA and the mean chronological age was 28.98 (SD=12.64) days at the beginning of the study. (Table 1 describes clinical and demographic variables.)

The analysis revealed that there were no statistically significant differences on the TIMP between the groups at beginning of 32 weeks PMA (pretreatment) ($P = .18$). All infants improved their raw scores on the TIMP at the end of 34 weeks PMA (the post intervention), but there was no significant difference in mean motor performance for raw ($P = .11$) or Z ($P = .11$) scores between the groups at the end of 34 weeks PMA. The post-intervention ANOVA showed no differences between the groups (Table 2). The proportions of infants in the categories of average motor performance between the study groups was not significantly different ($P = 0.26$) (Table 3). About neuromuscular development, the analysis indicated that there were not significant differences in post intervention on Posture, Arm Recoil, Popliteal Angle, Scarf Sign, Heel to Ear, Square window and Ankle dorsiflexion items. Just, leg recoil was a significant difference on physical activity and Hydrotherapy groups ($P<0.05$). Comparison of the scores for the 4 groups at 34 weeks PMA indicated that passive physical activities affected leg recoil of postural tone and that the other neuromuscular development changes over time were most likely due to maturation of all groups (Table 4).

Table 1. Comparison between study groups on baseline Characteristics (n=19)

Variable	Physical activity N (%)	Hydrotherapy N (%)	Combination N (%)	Containment N (%)	P
Gender[‡]					
Male	11(57)	8 (43)	7 (36.8)	5 (26.3)	0.25
Female	8 (43)	11 (57)	12 (63.2)	14 (73.7)	
Type of delivery[‡]					
Vaginal	2 (10.5)	5 (26.3)	6 (31.6)	8 (42.1)	0.17
Cesarean section	17 (89.5)	14 (73.7)	13 (68.4)	11(57.9)	
Pregnancy[‡]					
Single	15 (78.9)	12 (63.2)	15 (78.9)	13 (68.4)	0.69
Multiple	4 (21.1)	7 (21.1)	4 (21.1)	6 (31.6)	
Apgar score[§]					
first minute	2 - 9	4 - 9	3 - 9	3 - 8	0.53
5 th minute	6 - 10	7 - 10	5 - 10	6 -10	0.93
Birth weight[§] (g)	1038 (235.8)	1089 (176.1)	1033 (196.5)	1093 (179.1)	0.68
GA at birth[§] (wk)	25-30	25-30	25-30	26-30	0.64
Age at beginning study[§] (day)	8-55	10-53	12-64	9-56	0.52
Diagnosis[‡]					
Prematurity	19(100)	19(100)	19(100)	19(100)	1
Respiratory distress syndrome	19(100)	19(100)	19(100)	19(100)	1
Premature rupture of membrane	11(57.9)	9(47.4)	14(73.7)	12(63.2)	0.44
Preeclampsia	7(36.8)	7(36.8)	8(42.1)	4(21.1)	0.66
Vaginal bleeding	1(5.3)	2(10.5)	2(10.5)	0(0.0)	0.74
Number of surfactant administrate[§]	0-4	0-2	0-2	0-3	0.81
mechanical ventilation long stay[§] (day)	3-23	3-15	3-28	3-39	0.58
Length of sedative drug administration[§]	0-12	0-5	0-7	0-10	0.87

[‡] Chi-Square test was used. [§]The data are presented as mean (SD) and Maximum- Minimum using ANOVA one way test. [¶]The data are presented as Maximum- Minimum (Rang) using Kruskal-Wallis test.

Table 2. Test of infant motor performance scores by groups (n=19)

Variable	Physical activity Mean(SD)	Hydrotherapy Mean(SD)	Combination [§] Mean(SD)	Containment Mean(SD)	P [¶]
TIMP Score					
32 weeks	40.42 (6.43)	37.73(4.78)	38.89 (4.61)	41.42 (5.99)	0.18
34 weeks	50. 21(6.67)	48. 05(5.60)	52.00 (5.14)	51.57 (3.87)	0.11
Z score [‡]	0.08 (.44)	-0.06(0.37)	0.19(0.34)	0.17 (0.25)	0.11

[¶]ANOVA one way test was used. [§]Combination group received each other day physical activity and hydrotherapy interventions. [‡] Z score calculated as their raw score minus the mean for the 34 weeks group divided by the SD for the normative group.

Table 3. TIMP performance categories by group at 34 weeks PMA (n=19)

Variable	Physical activity N (%)	Hydrotherapy N (%)	Combination N (%)	Containment N (%)	P [¶]
TIMP category (34weeks)					
Average range [§]	17 (89.5)	18 (94.7)	18 (94.7)	19 (100)	0.26
low average [‡]	2 (10.5)	1 (5.3)	1 (5.3)	0 (0)	

[¶] Chi-Square Test was used. [§] Average Range =scores within -0.5 to+ 1 SD of the mean for the age group. [‡] Low average=Range of scores between -0.5 to -1 SD below the mean, a subset of the average range and the threshold for delay

Table 4. Post comparison of 8 items neuromuscular difference scores between groups

Variable		Physical activity N (%)	Hydrotherapy N (%)	Combination N (%)	Containment N (%)	P ^ε
Diff posture score	-1	1(5.3)	2(10.5)	0(0.0)	0(0.0)	0.62
	0	8(42.1)	5(26.3)	4(21.1)	12(63.2)	
	1	7(36.8)	10(52.6)	13(68.4)	6(31.6)	
	2	3(15.8)	2(10.5)	2(10.5)	1(5.3)	
	-2	0(0.0)	0(0.0)	0(0.0)	1(5.3)	
Diff Square window score	-1	3(15.8)	6(31.6)	3(15.8)	2(10.5)	0.41
	0	12(63.2)	10(52.6)	9(47.4)	10(52.6)	
	1	4(21.1)	1(5.3)	7(36.8)	4(21.1)	
	2	0(0.0)	2(10.5)	0(0.0)	2(10.5)	
	-2	1(5.3)	0(0.0)	0(0.0)	0(0.0)	
Diff Arm recoil score	-1	2(10.5)	3(15.8)	1(5.3)	2(10.5)	0.91
	0	5(26.3)	8(42.1)	9(47.4)	9(47.4)	
	1	9(47.4)	6(31.6)	7(36.8)	7(36.8)	
	2	2(10.5)	1(5.3)	1(5.3)	1(5.3)	
	3	0(0.0)	1(5.3)	1(5.3)	0(0.0)	
Diff popliteal angle score	0	6(31.6)	12(63.2)	5(26.3)	11(57.9)	0.89
	1	12(63.2)	5(26.3)	9(47.4)	7(36.8)	
	2	1(5.3)	2(10.5)	3(15.8)	1(5.3)	
	3	0(0.0)	0(0.0)	2(10.5)	0(0.0)	
Diff Heel to ear score	-1	1(5.3)	5(26.3)	2(10.5)	2(10.5)	0.96
	0	10(52.6)	3(15.8)	6(31.6)	7(36.8)	
	1	3(15.8)	7(36.8)	7(36.8)	5(26.3)	
	2	4(21.1)	3(15.8)	4(21.1)	5(26.3)	
	3	1(5.3)	1(5.3)	0(0.0)	0(0.0)	
Diff scarf sign score	-2	1(5.3)	0(0.0)	0(0.0)	0(0.0)	0.95
	-1	0(0.0)	2(10.5)	2(10.5)	1(5.3)	
	0	10(52.6)	8(42.1)	3(15.8)	11(57.9)	
	1	7(36.8)	6(31.6)	12(63.2)	7(36.8)	
	2	1(5.3)	4(21.1)	1(5.3)	0(0.0)	
Diff Leg recoil score	3	0(0.0)	0(0.0)	1(5.3)	0(0.0)	0.04*
	-1	1(5.3)	0(0.0)	0(0.0)	2(10.5)	
	0	9(47.4)	11(57.9)	15(78.9)	13(68.4)	
	1	9(47.4)	8(42.1)	4(21.1)	4(21.1)	
	-2	7(36.8)	0(0.0)	3(15.8)	2(10.5)	
Diff Ankle dorsiflexion score	-1	4(21.1)	4(21.1)	4(21.1)	7(36.8)	0.49
	0	5(26.3)	10(52.6)	8(42.1)	7(36.8)	
	1	3(15.8)	3(15.8)	1(5.3)	3(15.8)	
	2	0(0.0)	2(10.5)	2(10.5)	0(0.0)	
	3	0(0.0)	0(0.0)	1(5.3)	0(0.0)	

The different of after and before scores calculated and the data are presented as a frequency and percent. ^ε Chi-square test was used. *statistically significant

Discussion

The present study compares the effect of various types of physical activity programs on functional motor performance and neuromuscular development of postural tone in infants born preterm. The findings of this study showed that different types of physical activity programs have the same containment effects on functional motor performance as the development of postural tone except leg recoil. The groups did not differ statistically from each other, indicating that change was most likely due to maturation. The results of the study also indicated that there was no relationship between motor function and duration of hospital stay, days with mechanical ventilation, the duration of treatment with sedative drugs, and having received surfactant. The infants in this study, although born prior to 32 weeks GA, were relatively low risk without major medical complications, and 97% performed within the normal range on the TIMP at the end of the study (34 weeks PMA). The results might differ in older infants or those with more medical complications.

Several studies have investigated the effect of active therapy programs on motor performance in infants born preterm at older ages with varying results. For example, Piper et al., found that performance at 12 months of age in an experimental group was not superior to the control group in any of the assessed areas of development. They indicated that infants born at earlier gestational ages ($P < 32$ weeks), when compared with later ages (32-36 weeks), scored lower on tests of gross motor development. Short gestational age may relate to the quality of motor performance at a later age in infants born preterm.³⁹

Other studies indicated that experiences such as prolonged hospital stay, disorders related to medical complications or limitations resulting from medical technologies, septicemia and need for

mechanical ventilation that cause inactivity might be associated with infant's motor performance and postural stability.⁴⁰⁻⁴² On the other hand, Girolami and Campbell reported that a neuro-developmental treatment program was an effective method of improving motor performance of infants born preterm at high risk for developmental disabilities during their stay in the intensive care unit.³⁴ Resnick et al., reported that developmental intervention programs for high-risk infants born preterm resulted in a significant improvement in physical abilities, and the quality of infants' interaction with their parents in the treatment group was better than that in the control group.⁴³

The studies by Girolami and Campbell and by Resnick et al., showed that active physical activity has a positive effect on functional performance of premature infants. These results were not similar to the results in the current study due to different interventions (active versus passive), duration of treatment and follow up timing (immediately after intervention versus longer follow up periods). Piper et al., and Saylor et al., indicated there was no difference between the outcome of assessments for those infants who received early interventions at the intensive care unit and infants who received the treatment at older ages.^{39,44} These results were similar to the current study.

We used the parameters derived from New Ballard's assessment³⁵, an accepted measure of the functional state of the nervous system, to clinically measure the effect of different type physical activity programs on the level of maturation of the neuromuscular system. This study appears to provide the first evidence that physical activities in or out of the water during their hospital stay does not produce significant change in neuromuscular development, except leg recoil, of infants born preterm. Leg recoil provides information on the reaction of the lower limbs to physical activity. The post comparison of 8 items

indicated that change in leg recoil was significantly better in the physical activity, hydrotherapy and combination groups. The literature search did not reveal any previous studies which investigated the short term effects of passive physical activities on neuromuscular development in preterm infants.

The duration of the intervention was short, and we believe that had we applied the movements for a longer period, a greater advancement in neuromuscular maturity could have been achieved in the trunk and limbs, which mature at a later age. Additional studies are needed to determine if, with optimal timing and duration, or an integrated different kind of passive and active intervention can yield a better neuromuscular outcome in infants born preterm.

Although this study did not show a positive effect for the physical activity on motor functional performance, we cannot yet ignore the effectiveness of this program in other ways. Some evidence demonstrated that physical activity programs might promote short-term weight gain and bone mineralization in premature infants^{20,45} and hydrotherapy had a positive effect on pain reduction, improvement of sleep quality, weight gain and feeding tolerance among infants born preterm.^{22,23} The procedures used in this study were tolerated well by the infants and thus passive exercise, either in or out of the water, appears to be a safe intervention.

The strength and exclusive aspect of the present study was its comparison of different types of passive exercise with development in a containment group. In order to minimize potential biases, the principles of clinical trial studies, including complete randomization of the participants and blinding the examiners and the data analyzers, was done.

The limitation of this study was the length of the intervention and the shortness of the follow-up time. Another limitation of the study was that the hydrotherapy group

received less intervention overall than did the other groups, so its effectiveness might have been compromised by being an insufficient dose. Thus, it is suggested that another study perform the interventions for a longer period, with similar dosages or other protocols with different design, and assess motor performance and neuromuscular development. When the infants reach 34 weeks PMA, active physical therapy might be added to investigate the combined effect on bone density, functional movement performance, and neuromuscular development of a longer period of passive and active intervention. Finally, the parents were not involved in this study and a new study could investigate the benefits of having interventions done by the family.

Conclusion

This study demonstrates that passive physical activities have as the same containment effects in infants born preterm on motor performance and most items of postural tone except leg recoil. Understanding the effect of passive versus active physical therapy by using reliable and valid tests such as the TIMP may be useful to clinicians to make decisions regarding what kind of physical activity or containment can help high risk preterm infants and what dosage is useful to compensate for the complications of preterm birth.

Acknowledgments

We would like to thank Dr. Suzann K. Campbell for advice on use of the TIMP and help with scientific writing and Dr. Jeanne Ballard who helped us to correctly conduct this study with her professional instruction. We also appreciate the personnel of the neonatal intensive care units of Al-Zahra teaching hospital in Tabriz as well as the parents who helped us in conducting this study by giving permission to include their infants in the study.

Ethical issues

None to be declared.

Conflict of interest

The authors declare no conflict of interest in this study.

References

1. Saigal S, Doyle LW. An overview of mortality and sequelae of preterm birth from infancy to adulthood. *Lancet* 2008 ; 371 (9608) : 261-9. doi: [10.1016/S0140-6736\(08\)60136-1](https://doi.org/10.1016/S0140-6736(08)60136-1)
2. Blencowe H, Lee ACC, Cousens S, Bahalim A, Narwal R, Zhong N, et al. Preterm birth-associated neurodevelopmental impairment estimates at regional and global levels for 2010. *Pediatr Res* 2013; 74 (S1) : 17-34. doi: [10.1038/pr.2013.204](https://doi.org/10.1038/pr.2013.204).
3. Kinney MV, Lawn JE, Howson CP, Belizan J. 15 million preterm births annually: what has changed this year? *Reproductive Health* 2012; 9 (1) : 28. doi: [10.1186/1742-4755-9-28](https://doi.org/10.1186/1742-4755-9-28)
4. Wilson-Costello D, Friedman H, Minich N, Siner B, Taylor G, Schluchter M, et al. Improved neurodevelopmental outcomes for extremely low birth weight infants in 2000–2002. *Pediatrics* 2007 ; 119 (1) : 37-45.
5. Robertson CM, Watt MJ, Dinu IA. Outcomes for the extremely premature infant: what is new? And where are we going? *Pediatr Neurol* 2009 ; 40 (3): 189-96. doi: [10.1016/j.pediatrneurol.2008.09.017](https://doi.org/10.1016/j.pediatrneurol.2008.09.017)
6. Arpino C, Compagnone E, Montanaro ML, Cacciatore D, De Luca A, Cerulli A, et al. Preterm birth and neurodevelopmental outcome: a review. *Childs Nerv Syst* 2010 ; 26 (9): 1139-49. doi: [10.1007/s00381-010-1125-y](https://doi.org/10.1007/s00381-010-1125-y).
7. Vanderveen JA, Bassler D, Robertson CM, Kirpalani H. Early interventions involving parents to improve neurodevelopmental outcomes of premature infants: a meta-analysis. *J Perinatol* 2009 ; 29 (5) : 343-51. doi: [10.1038/jp.2008.229](https://doi.org/10.1038/jp.2008.229).
8. Grunau RE, Whitfield MF, Petrie-Thomas J, Synnes AR, Cepeda IL, Keidar A, et al. Neonatal pain, parenting stress and interaction, in relation to cognitive and motor development at 8 and 18 months in preterm infants. *Pain* 2009 ; 143 (1-2): 138-46. doi: [10.1016/j.pain.2009.02.014](https://doi.org/10.1016/j.pain.2009.02.014).
9. Volpe JJ. Brain injury in premature infants: a complex amalgam of destructive and developmental disturbances. *Lancet Neurol* 2009; 8 (1) : 110-24.
10. de Kieviet JF, Piek J, Aarnoudse C, Oosterlaan J. Motor development in very preterm and very low-birth-weight children from birth to adolescence: a meta-analysis. *JAMA* 2011; 31 (1): 41-2. doi: [10.1001/jama.2009.1708](https://doi.org/10.1001/jama.2009.1708)
11. Limperopoulos C, Bassan H, Gauvreau K, Robertson RL, Sullivan NR, Benson CB, et al. Does cerebellar injury in premature infants contribute to the high prevalence of long-term cognitive, learning, and behavioral disability in survivors? *Pediatrics* 2007 ; 120 (3) : 584-93.
12. Allen MC. Neurodevelopmental outcomes of preterm infants. *Curr Opin Neurol* 2008; 21 (1): 123-8. doi: [10.1097/WCO.0b013e3282f88bb4](https://doi.org/10.1097/WCO.0b013e3282f88bb4).
13. Feldman R, Eidelman AI. Neonatal state organization, neuromaturation, mother-infant interaction, and cognitive development in small-for-gestational-age premature infants. *Pediatrics* 2006 ; 118 (3) : e869-e78. doi: [10.1542/peds.2005-2040](https://doi.org/10.1542/peds.2005-2040).
14. Downs JA, Edwards AD, McCormick DC, Roth SC, Stewart AL. Effect of intervention on development of hip posture in very preterm babies. *Arch Dis Child* 1991 ; 66 (7 Spec No): 797-801. doi: [10.1136/adc.66.7.Spec.No.797](https://doi.org/10.1136/adc.66.7.Spec.No.797).
15. Korner AF. Infant stimulation. Issues of theory and research. *Clinics in perinatology* 1990 ; 17 (1) : 173-84.
16. Spittle A, Orton J, Anderson PJ, Boyd R, Doyle LW. Early developmental

- intervention programmes provided post hospital discharge to prevent motor and cognitive impairment in preterm infants. *Cochrane Database Syst Rev* 2015 ; (11). doi: [10.1002/14651858.CD005495.pub3](https://doi.org/10.1002/14651858.CD005495.pub3)
17. Spittle A, Orton J, Anderson P, Boyd R, Doyle LW. Early developmental intervention programmes post-hospital discharge to prevent motor and cognitive impairments in preterm infants. *Cochrane Database Syst Rev* 2012 ; 12 : Cd005495. doi: [10.1002/14651858.CD005495](https://doi.org/10.1002/14651858.CD005495).
 18. Park HY, Maitra K, Achon J, Loyola E, Rincón M. Effects of early intervention on mental or neuromusculoskeletal and movement-related functions in children born low birthweight or preterm: a meta-analysis. *Am J Occup Ther* 2014 ; 68 (3) : 268-76. doi: [10.5014/ajot.2014.010371](https://doi.org/10.5014/ajot.2014.010371).
 19. Moyer-Mileur LJ, Ball SD, Brunstetter VL, Chan GM. Maternal-administered physical activity enhances bone mineral acquisition in premature very low birth weight infants. *J Perinatol* 2008 ; 28 (6) : 432-7. doi: [10.1038/jp.2008.17](https://doi.org/10.1038/jp.2008.17).
 20. Vignochi CM, Miura E, Canani LH. Effects of motor physical therapy on bone mineralization in premature infants: a randomized controlled study. *J Perinatol* 2008 ; 28 (9): 624-31. doi: [10.1038/jp.2008.60](https://doi.org/10.1038/jp.2008.60).
 21. Tosun Ö, Bayat M, Güneş T, Erdem E. Daily physical activity in low-risk pre-term infants: positive impact on bone strength and mid-upper arm circumference. *Ann Hum Biol* 2011; 38 (5) : 635-9. doi: [10.3109/03014460.2011.598187](https://doi.org/10.3109/03014460.2011.598187).
 22. Sweeney JK. Neonatal hydrotherapy: an adjunct to developmental intervention in an intensive care nursery setting. *Phys Occup Ther Pediatr* 1983 ; 3 (1) : 39-52.
 23. Vignochi C, Teixeira PP, Nader SS. Efeitos da fisioterapia aquática na dor e no estado de sono e vigília de recém-nascidos pré-termo estáveis internados em unidade de terapia intensiva neonatal. *Rev Bras Fisioter* 2010 ; 14 (3) : 214-20.
 24. de Oliveira Tobinaga WC, de Lima Marinho C, Abelenda VLB, de Sá PM, Lopes AJ. Short-Term effects of hydrokinesiotherapy in hospitalized preterm newborns. *Rehabil Res Pract* 2016 ; 2016. doi: [10.1155/2016/9285056](https://doi.org/10.1155/2016/9285056)
 25. Vakilian K, Ranjbaran M, Khorsandi M, Sharafkhani N, Khodadost M. Prevalence of preterm labor in Iran: a systematic review and meta-analysis. *Int J Reprod Biomed* 2015 ; 13 (12): 743-8.
 26. Blauw-Hospers CH, Hadders-Algra M. A systematic review of the effects of early intervention on motor development. *Dev Med Child Neurol* 2005 ; 47 (6) : 421-32. doi: [10.1111/j.1469-8749.2005.tb01165.x](https://doi.org/10.1111/j.1469-8749.2005.tb01165.x).
 27. Spittle AJ, Doyle LW, Boyd RN. A systematic review of the clinimetric properties of neuromotor assessments for preterm infants during the first year of life. *Dev Med Child Neurol* 2008 ; 50 (4) : 254-66. doi: [10.1111/j.1469-8749.2008.02025.x](https://doi.org/10.1111/j.1469-8749.2008.02025.x).
 28. Rose RU, Westcott SL. Responsiveness of the test of infant motor performance (TIMP) in infants born preterm. *Pediatr Phys Ther* 2005 ; 17 (3) : 219-24.
 29. Campbell SK, Kolobe TH, Osten ET, Lenke M, Girolami GL. Construct validity of the test of infant motor performance. *Phys Ther* 1995 ; 75 (7) : 585-96.
 30. Byrne E, Campbell SK. Physical therapy observation and assessment in the neonatal intensive care unit. *Phys Occup Ther Pediatr* 2013; 33 (1) : 39-74. doi: [10.3109/01942638.2012.754827](https://doi.org/10.3109/01942638.2012.754827).
 31. Campbell SK. The test of infant motor performance, test user's manual version 3.0 for the timp version 5. 1st ed. Chicago I: Infant motor performance scales,LLC; 2012.
 32. Fucile S, Gisel EG. Sensorimotor interventions improve growth and motor function in preterm infants. *Neonatal Netw* 2010 ; 29 (6) : 359-66. doi: [10.1891/0730-0832.29.6.359](https://doi.org/10.1891/0730-0832.29.6.359).
 33. Ustad T, Evensen KA, Campbell SK, Girolami GL, Helbostad J, Jorgensen L, et

- al. Early parent-administered physical therapy for preterm infants: a randomized controlled trial. *Pediatrics* 2016 ; 138 (2). doi: [10.1542/peds.2016-0271](https://doi.org/10.1542/peds.2016-0271).
34. Girolami GL, Campbell SK. Efficacy of a neuro-developmental treatment program to improve motor control in infants born prematurely. *Pediatric Physical Therapy* 1994 ; 6 (4) : 175-84.
35. Ballard JL, Khoury JC, Wedig K, Wang L, Eilers-Walsman BL, Lipp R. New ballard score, expanded to include extremely premature infants. *J Pediatr* 1991 ; 119 (3) : 417-23.
36. Dubowitz LM, Dubowitz V, Goldberg C. Clinical assessment of gestational age in the newborn infant. *J Pediatr* 1970 ; 77 (1) : 1-10.
37. Als H, Butler S, Kosta S, McAnulty G. The Assessment of Preterm Infants' Behavior (APIB): Furthering the understanding and measurement of neurodevelopmental competence in preterm and full-term infants. *Ment Retard Dev Disabil Res Rev* 2005 ; 11 (1) : 94-102. doi: [10.1002/mrdd.20053](https://doi.org/10.1002/mrdd.20053).
38. Burns K, Cunningham N, White-Traut R, Silvestri J, Nelson MN. Infant stimulation: Modification of an intervention based on physiologic and behavioral cues. *J Obstet Gynecol Neonatal Nurs* 1994 ; 23 (7) : 581-9.
39. Piper MC, Kunos VI, Willis DM, Mazer BL, Ramsay M, Silver KM. Early physical therapy effects on the high-risk infant: a randomized controlled trial. *Pediatrics* 1986 ; 78 (2) : 216-24.
40. Case-Smith J. Postural and fine motor control in preterm infants in the first six months. *Phys Occup Ther Pediatr* 1993 ; 13 (1): 1-17. doi: [10.1080/J006v13n0101](https://doi.org/10.1080/J006v13n0101).
41. Georgieff MK, Bernbaum JC. The High incidence of abnormal shoulder girdle muscle tone in premature Infants during the first year of life. *Pediatric Research* 1984; 18 (S4) : 104A. doi: [10.1203/00006450-198404001-00065](https://doi.org/10.1203/00006450-198404001-00065) .
42. De Vries NKS, Erwich JJHM, Bos A. General movements in the first fourteen days of life in extremely low birth weight (ELBW) infants. *Early Hum Dev* 2008 ; 84 (11) : 763-8. doi: [10.1016/j.earlhumdev.2008.05.003](https://doi.org/10.1016/j.earlhumdev.2008.05.003).
43. Resnick MB, Armstrong S, Carter RL. Developmental intervention program for high-risk premature infants: effects on development and parent-infant interactions. *J Dev Behav Pediatr* 1988 ; 9 (2): 73-8.
44. Saylor CF, Casto G, Huntington L. Predictors of developmental outcomes for medically fragile early intervention participants. *J Pediatr Psychol* 1996;21 (6): 869-87.
45. Schulzke SM, Kaempfen S, Trachsel D, Patole SK. Physical activity programs for promoting bone mineralization and growth in preterm infants. *Cochrane Database of Systematic Reviews* 2014(4). doi: [10.1002/14651858.CD005387](https://doi.org/10.1002/14651858.CD005387)