



Study the Interactive Effects of Different Nitrogen Doses and Spacing on Quality Characteristics of Coriander (*Coriandrum sativum* L.)

Chaman Vats ^a, T. P. Malik ^a, Mukesh Bishnoi ^{b*},
Uday Singh ^a, Randeep Singh ^c and Abhishek Kumar ^d

^a Department of Vegetable Science, Chaudhary Charan Singh Haryana Agricultural University, Hisar, Haryana 125004, India.

^b Department of Horticulture, Chaudhary Charan Singh Haryana Agricultural University, Hisar, Haryana 125004, India.

^c Department of Horticulture, Govt. of Haryana, India.

^d Department of Plant Pathology, Chaudhary Charan Singh Haryana Agricultural University, Hisar, Haryana 125004, India.

Authors' contributions

This work was carried out in collaboration among all authors. Authors CV and TPM performed the conceptualization, methodology, investigation, writing-original draft preparation. Author MB did the reviewing and editing. Authors RS, US and AK did the reviewing the final draft and editing. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i213942

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/107765>

Original Research Article

Received: 01/08/2023

Accepted: 07/10/2023

Published: 13/10/2023

ABSTRACT

Coriander (*Coriandrum sativum* L.) is one of the most important spice crops worldwide, possessing varied medicinal impacts. It is majorly grown for its leaves; however, its seeds occupy a major place in various cuisines around the world. Several studies have reported the interactive effects of

*Corresponding author: E-mail: mukeshbishnoi1996@gmail.com;

different nitrogen doses and spacing in coriander and provided valuable insights into the optimal conditions for its growth and development. In this sequence, the present study explores the effect of various nitrogen doses and spacing on coriander under field conditions at Research Farm of Vegetable Science, Chaudhary Charan Singh Haryana Agricultural University, Hisar (India) during *Rabi*, 2019-20. Amongst all the treatments, treatment S_3N_3 (40×15; N@75 kg/ha) was found best for harvest index, Test weight, Standard Germination, Seedling Length, Seedling Dry weight, Vigour Indices I & II, and S_2N_1 (30×15; N@50 kg/ha) for minimum Electrical Conductivity. Based on this study, the following treatments can be exploited for sustainable coriander production.

Keywords: *Rabi*; spacing; sustainable.

1. INTRODUCTION

Coriander (*Coriandrum sativum* Linn.), a member of *Apiaceae* family which is also known as *Dhania*, cilantro, and Chinese parsley; is grown as an annual herb around the world. It is originated from the Mediterranean region, however, extensively cultivated in Central Europe North Africa, and Asia as a medicinal as well as culinary herb [1]. Its matured fruits and fresh leaves are used as spices and traditional medicine. The essential oil content in matured coriander fruits ranges from 0.03% to 2.7%. Linalool, the primary component of the oil, constitutes approximately two-thirds of its composition [2]. India is the largest producer of coriander in the world. Within Indian courtesy, it is mainly cultivated in Rajasthan, Madhya Pradesh, Gujarat, Andhra Pradesh, Tamil Nadu, Karnataka, Haryana and Punjab, West Bengal, and Himachal Pradesh. The crop was grown over an area of 583 thousand hectares with the production of 784 thousand MT (NHB, 2018-19) and in Haryana, the crop occupied an area of 2.40 thousand hectares with the production of 4.4 thousand metric tonnes during 2018-19 [3].

In India, it is primarily grown in *Rabi* season specifically for its leaves and seeds [4] Besides various culinary applications, it possess a range of medicinal properties [5] owing to its broad phytochemical profile and a variety of chemical compounds present in every plant part (root, fruits, leaves, and seeds) [6]. There are several documented health benefits of coriander in literature i.e., swellings, diarrhoea, mouth ulcers, anaemia, digestion, menstrual disorders, smallpox, eye care, conjunctivitis, skin disorders, blood sugar disorders, and protection and soothing liver, etc. [7-9]

The application of nitrogen fertilizer has been found to enhance plant growth, as well as increase the production of essential oil, fixed oil, total carbohydrates, and soluble sugars. Nitrogen

plays a crucial role in key plant processes such as photosynthesis, respiration, and protein synthesis. It contributes to the development of dark green leaves, stimulates vigorous vegetative growth, improves the utilization of available resources, and ultimately results in higher productivity [2]. In the coriander cultivation system, nitrogen fertilization is considered a major production factor that imparts a direct effect on the quality, composition, and the yield of volatile compounds. As per previous documentations, the application of nitrogen fertilizer has improved seed yield and biomass of coriander by 43-68% and 25-42% [10]. In contrast, a nitrogen deficiency in plants leads to the yellowing of lower leaves, stunted growth, and the premature shedding of both leaves and fruits, which directly impacts overall yield negatively.

On the other hand, excessive nitrogen application results in excessive shoot growth, rendering plants more vulnerable to pest and disease attacks. It also hampers root development, leading to lodging, and causes a delay in crop maturity. Consequently, the excessive application of nitrogen diminishes crop yield and degrades the quality of the produce [11]. Therefore, the incorporation of essential plant nutrients in the right proportion and adequate quantity concerning soil and climatic conditions is very essential [10]. Besides, maintaining the optimum plant population is also a beneficial, non-monetary agronomical practice of crop production. Maintaining optimum density of plants results in reduced nutrient competition within the plant population. Various studies have reported that optimum plant density increased the growth and yield of coriander significantly [12]. However, Wider or too close spacing beyond the optimum can negatively affect the yield of a crop [13]. Therefore, the present investigation was undertaken to demonstrate the effect of various nitrogen and spacing effects on coriander under field conditions.

2. MATERIALS AND METHODS

2.1 Experimental Site

The field trial was conducted at Research Farm of Vegetable Science Chaudhary Charan Singh [14] Haryana Agricultural University, Hisar (India) (29° 10' N, 75° 46' E, Elevation 215.2 m) in the *Rabi* season of 2019-20. The temperature varies from 40 °C to 48 °C during summer to as low as to freezing point accompanied with chilling frost in winters. The approximate average rainfall was 450 mm.

2.2 Properties of the Soil before the Experiment Conducted

The soil of the experimental field was analyzed for mechanical and chemical properties, and cropping history details are given below in Table 1.

2.3 Field Preparation

2.3.1 Experimental details

The coriander variety DH 220, developed by department of vegetable science, was chosen for experiment. The seeds were sown in a plot size of 3.0 m x 2.4 m with Randomized Block Design

(RBD) in three replications. A total of twelve treatments (Table 2) were formulated and raised with the standard agronomic practices. In the study, half of the nitrogen was applied at the sowing time and the rest of the dose was applied after 6 weeks of the first dose. The five were selected randomly and tagged data collection.

2.4 Evaluation of Physical Parameters

Harvest Index: The calculation of the harvest index, expressed as a percentage, is determined by dividing the economic yield by the biological yield using the following formula:

$$\text{Harvest index (\%)} = (\text{Economic yield} / \text{Biological yield}) \times 100$$

Test Weight: The count and weight of one thousand seeds were measured in triplicate for each treatment, and the average seed weight for each treatment was calculated.

Standard Germination: In each replication of the experiment, fifty seeds from each treatment were placed between germination papers and kept at a temperature of 20°C with 90-95% relative humidity in a germinator. The first count of germinated seeds was recorded on the 7th day, and the final count was taken on the

Table 1. Detailed description of mechanical, and chemical analysis of soil and cropping history

Mechanical analysis of the soil			
Sr. No.	Soil parameters	Proportion (%)	Methods and reference
1	Sand	56	International pipette method
2	Silt	32	[15]
3	Clay	12	
4	Soil texture	Sandy - loam	
Chemical analysis of the soil at the start of the experiment			
S. No.	Soil Parameters	Value	Methods and reference
1	pH (1:2 soil: water suspension)	8.1	Potentiometric method [16]
2	EC (ds/m) at 25°C (1:2 soil: water suspension)	0.36	Conductometric method [16]
3	Organic Carbon (%)	0.35	Wet oxidation method [17]
4	Available nitrogen (kg/ha)	138	Kjeldhal- distillation method [18]
5	Available phosphorus (kg/ha)	22.5	NaHCO ₃ extraction and colorimetry method [19]
6	Available potassium (kg/ha)	227	N NH ₄ OAC extraction and Flame photometry method [16]
Cropping history of the experimental field			
Crop seasons			
Year	<i>Kharif</i>	<i>Rabi</i>	
2017-18	<i>Okra</i>	<i>Potato</i>	
2018-19	<i>Okra</i>	<i>Fennel</i>	
2019-20	Bottle gourd	Coriander	

Table 2. Detailed treatment dose of N with combinatorial application with various spacing

Treatment (number)	A detailed description of treatments
1	N ₁ : 0 kg/ha (control)+ S ₁ : 20x15 cm
2	N ₁ : 0 kg/ha (control)+ S ₂ : 30x15 cm
3	N ₁ : 0 kg/ha (control)+ S ₃ : 40x15 cm
4	N ₂ : 25 kg/ha+ S ₁ : 20x15 cm
5	N ₂ : 25 kg/ha+ S ₂ : 30x15 cm
6	N ₂ : 25 kg/ha+ S ₃ : 40x15 cm
7	N ₃ : 50 kg/ha+ S ₁ : 20x15 cm
8	N ₃ : 50 kg/ha+ S ₂ : 30x15 cm
9	N ₃ : 50 kg/ha+ S ₃ : 40x15 cm
10	N ₄ : 75 kg/ha+ S ₁ : 20x15 cm
11	N ₄ : 75 kg/ha+ S ₂ : 30x15 cm
12	N ₄ : 75 kg/ha+ S ₃ : 40x15 cm

21st day. The recorded observations for seed germination were then converted into a percentage to determine the per cent seed germination.

Seedling Length: At the end of the germination test, the root and shoot length of five randomly chosen seedlings from each treatment in each replication were measured in centimetres. These measurements were then averaged to determine the total root and shoot length.

Seedling Dry Weight: After the final count in the standard germination test (21 days), the dry weight of seedlings was evaluated. Five seedlings from each treatment, replicated three times, were selected for assessment. The seedlings were dried in a hot air oven for 48 hours at a temperature of 80±10 °C. Following the drying process, the seedlings from each replication were weighed, and the average seedling dry weight for each treatment was calculated.

Vigour Indices: Seedling vigour indices were calculated according to the method suggested by Abdul- Baki and Anderson (1973):

- I. Vigour index-I (on seedling length basis):
Vigour index-I = Standard germination (%) × Average seedling length (cm)
- II. Vigour Index-II (on seedling dry weight basis):
Vigour index-II = Standard germination (%) × Average seedling dry weight (mg)

2.5 Electrical Conductivity Test

To assess the electrical conductivity, three replications of 50 intact and undamaged seeds were immersed in 75 ml of distilled water within

100 ml beakers. The seeds were fully submerged in the water, and the beakers were covered with foil. Subsequently, these samples were left at a temperature of 25°C for 24 hours. The electrical conductivity of the leachates from the seeds was measured using a direct reading conductivity meter, with the conductivity value expressed in micro Siemens per centimetre per gram ($\mu\text{S cm}^{-1}\text{g}^{-1}$).

2.6 Statistical Analysis

The data was statistically analyzed in Randomized Block Design using SPSS software (IBM, SPSS Inc., USA). The p-values were calculated and the results were expressed as CD at 5% level of significance. Data was subjected to two-way ANOVA (Gomez and Gomez 2010).

3. RESULTS AND DISCUSSION

3.1 Harvest Index

Harvest index was recorded maximum in spacing 40 x 15 cm (37.50) which was followed by 30 x 15 cm (37.12). In their study, Sharma et al. (2016) found that spacing played a significant role in promoting vigorous plant growth, as evidenced by abundant branching and increased biomass accumulation per plant. The extensive branching not only facilitated enhanced flower initiation but also ensured an optimal supply of metabolites. The higher biomass per plant likely contributed to better retention of flowers, resulting in greater seed formation and seed growth (Table 3).

Harvest index was recorded maximum at nitrogen dose 75 kg ha⁻¹ (39.22) which was followed by 50 kg ha⁻¹ (38.67) Choudhary et al. [14] and they stated that the enhancement in

yield of coriander is directly associated with the parallel increase in growth and yield attributing characters. As nitrogen is a major nutrient of biological significance, it is mainly required in the synthesis of protein, chlorophyll, and other organic compounds. Therefore, with an increased availability of nitrogen in the soil medium and thereafter, effective absorption and translocation in various parts of plants imparts active cell division and elongation which results in greater plant height.

3.2 Test Weight

The maximum test weight was observed in spacing 40 x 15 cm (9.05 g) which was followed by 30 x 15 cm (8.87 g), Katar and Katar [20] and Diwan et al. [21] they stated that it might be due to broader spacing as it provides adequate water, nutrients, air and light which endorses

better growth of plants and help in generating the maximum weight of seeds. The test weight (g) was recorded maximum at nitrogen dose 75 kg ha⁻¹ (9.58) which was followed by 50 kg ha⁻¹ (9.06). According to Diwan et al. [21], the impact of nitrogen on seed production occurs indirectly through an increase in the availability of nutrients to the floral components. This implies that maintaining a large and photo synthetically efficient leaf area during the flowering stage is crucial for maximizing seed weight (Table 4).

3.3 Standard Germination

The maximum standard germination (%) was recorded at nitrogen dose 75 kg ha⁻¹ (88.89) followed by 50 kg ha⁻¹ (86.56) and maximum standard germination (%) was observed in spacing 40 x 15 cm (85.58) followed by 30 x 15 cm (84.00) (Table 5).

Table 3. Interaction effect of nitrogen levels under different spacing on harvest index on coriander (*Coriandrum sativum* L.)

Treatments	Harvest Index (%)			Mean N
	S ₁ (20 x 15 cm)	S ₂ (30 x 15 cm)	S ₃ (40 x 15 cm)	
N ₀ = Control (0 kg/ha)	31.96	33.49	34.33	33.26
N ₁ = 25 kg/ha	34.81	35.67	35.67	35.38
N ₂ = 50 kg/ha	37.00	39.33	39.67	38.67
N ₃ = 75 kg/ha	37.33	40.00	40.33	39.22
Mean S	35.28	37.12	37.50	
CD (5%)	N= 0.75	S= 0.65	N x S= NS	

Table 4. Interaction effect of nitrogen levels under different spacing on test weight on coriander (*Coriandrum sativum* L.)

Treatments	Test Weight (g)			Mean N
	S ₁ (20 x 15 cm)	S ₂ (30 x 15 cm)	S ₃ (40 x 15 cm)	
N ₀ = Control (0 kg/ha)	7.97	8.13	8.23	8.11
N ₁ = 25 kg/ha	8.33	8.33	8.40	8.36
N ₂ = 50 kg/ha	8.50	9.17	9.50	9.06
N ₃ = 75 kg/ha	8.83	9.83	10.07	9.58
Mean S	8.41	8.87	9.05	
CD (5%)	N= 0.49	S= 0.39	N x S= NS	

Table 5. Interaction effect of nitrogen levels under different spacing on Standard Germination on coriander (*Coriandrum sativum* L.)

Treatments	Standard Germination (%)			Mean N
	S ₁ (20 x 15 cm)	S ₂ (30 x 15 cm)	S ₃ (40 x 15 cm)	
N ₀ = Control (0 kg/ha)	78.00	78.33	79.67	78.67
N ₁ = 25 kg/ha	81.00	81.67	83.33	82.00
N ₂ = 50 kg/ha	85.00	86.67	88.00	86.56
N ₃ = 75 kg/ha	86.00	89.33	91.33	88.89
Mean S	82.50	84.00	85.58	
CD (5%)	N= 2.30	S= 1.99	N x S= NS	

Amare et al. [22] observed similar findings in onions, where the combined effect of narrower plant spacing and increased fertilization led to a higher proportion of seeds originating from primary umbels. This improvement in seed quality can be attributed to the larger size of embryos obtained from primary umbel seeds compared to those obtained from secondary and tertiary umbels. This is likely due to the active involvement of minerals in embryo development and the activation of hormones during seed germination.

3.4 Seedling Length

The maximum seedling length (cm) was obtained with nitrogen dose 75 kg ha⁻¹ (15.4 cm) followed by 50 kg ha⁻¹ (15.1 cm) and maximum seedling length (cm) was found with spacing 40 x 15 cm (14.5 cm) followed by 30 x 15 cm (14.3 cm) (Table 6).

Harshita et al. [23] found similar outcomes in coriander, and Kumar et al. [24] observed comparable results in fennel. In both studies, it was observed that an increased synthesis of photosynthates resulted in the formation of larger, bolder seeds. This can be attributed to a higher translocation of photosynthates into the seeds. Bolder seeds were found to contain more reserve food in their cotyledons, leading to greater seedling length in fennel plants.

3.5 Seedling Dry Weight

The maximum seedling dry weight (mg) was observed with nitrogen applied at 75 kg ha⁻¹ (5.11 mg) followed by 50 kg ha⁻¹ (4.83 mg) and maximum seedling dry weight (mg) was found with spacing 40 x 15 cm (4.58 mg) followed by 30 x 15 cm (4.45 mg). Similar results were observed by Harshita et al. (2017) in coriander and Kumar et al. (2019) in fennel that accumulation of more photosynthates which aided in formation of bolder seeds due to higher

translocation of photosynthates in the seed. Bolder seeds have more reserve food present in the cotyledons, which resulted into higher seedling dry weight (mg) of the fennel plant (Table 7).

3.6 Vigour Indices

Maximum vigour index-I was recorded with nitrogen applied at 75 kg ha⁻¹ (1348.03) followed by nitrogen supplied at 50 kg ha⁻¹ (1302.98) and maximum vigour index-I was found with spacing 40 x 15 cm (1236.19) followed by 30 x 15 cm (1202.77). Similar results were observed by Harshita et al. [23] in coriander and Amare et al. [22] in fennel that in larger seed size there is more reserved food in the cotyledon of the seeds to sustain the seedlings growth and make seedlings to be vigorous than the smaller seed sizes whose reserved food could be exhausted very soon. These result into higher vigour index-I (Tables 8 and 9).

Maximum vigour index-II (449.62) was recorded with nitrogen applied at 75 kg ha⁻¹ followed by nitrogen supplied at 50 kg ha⁻¹ (410.84) and maximum index-II was found with spacing 40 x 15 cm (393.27) followed by 30 x 15 cm (375.27). Similar results were observed by Harshita et al. [23] in coriander and Amare et al. [22] in fennel stated that broader spacing with adequate application of fertilizer produced larger size seeds and gave high vigour index-II. These results showed the highest vigour index-II.

3.7 Electrical Conductivity

Minimum electrical conductivity ($\mu\text{S cm}^{-1}\text{g}^{-1}$) (129 $\mu\text{S cm}^{-1}\text{g}^{-1}$) was recorded with nitrogen applied at 50 kg ha⁻¹ followed by nitrogen supplied at 75 kg ha⁻¹ (137 $\mu\text{S cm}^{-1}\text{g}^{-1}$) and minimum electrical conductivity ($\mu\text{S cm}^{-1}\text{g}^{-1}$) was found with spacing 30 x 15 cm (138 $\mu\text{S cm}^{-1}\text{g}^{-1}$) followed by 40 x 15 cm (140 $\mu\text{S cm}^{-1}\text{g}^{-1}$). In coriander, Harshita et al. [23] and in fennel, Kumar et al. [24] obtained

Table 6. Interaction effect of nitrogen levels under different spacing on Seedling Length on coriander (*Coriandrum sativum* L.)

Treatments	Seedling Length (cm)			Mean N
	S ₁ (20 x 15 cm)	S ₂ (30 x 15 cm)	S ₃ (40 x 15 cm)	
N ₀ = Control (0 kg/ha)	12.4	13.0	13.3	12.9
N ₁ = 25 kg/ha	13.0	13.4	13.6	13.3
N ₂ = 50 kg/ha	14.4	15.3	15.4	15.1
N ₃ = 75 kg/ha	14.8	15.5	15.7	15.4
Mean S	13.6	14.3	14.5	
CD (5%)	N= 0.29	S= 0.25	N x S= NS	

Table 7. Interaction effect of nitrogen levels under different spacing on Seedling Dry Weight on coriander (*Coriandrum sativum* L.)

Treatments	Seedling Dry Weight (mg)			Mean N
	S ₁ (20 × 15 cm)	S ₂ (30 × 15 cm)	S ₃ (40 × 15 cm)	
N ₀ = Control (0 kg/ha)	3.23	3.63	3.73	3.53
N ₁ = 25 kg/ha	3.90	3.97	4.10	3.99
N ₂ = 50 kg/ha	4.47	4.93	5.10	4.83
N ₃ = 75 kg/ha	4.70	5.27	5.37	5.11
Mean S	4.08	4.45	4.58	
CD (5%)	N= 0.10	S= 0.09	N × S= 0.17	

Table 8. Interaction effect of nitrogen levels under different spacing on Vigour Index - I on coriander (*Coriandrum sativum* L.)

Treatments	Vigour Index - I			Mean N
	S ₁ (20 × 15 cm)	S ₂ (30 × 15 cm)	S ₃ (40 × 15 cm)	
N ₀ = Control (0 kg/ha)	966.22	1016.43	1052.45	1011.70
N ₁ = 25 kg/ha	1059.14	1097.63	1131.05	1095.94
N ₂ = 50 kg/ha	1223.66	1327.18	1358.08	1302.98
N ₃ = 75 kg/ha	1271.10	1369.84	1403.15	1348.03
Mean S	1130.03	1202.77	1236.19	
CD (5%)	N= 46.44	S= 40.22	N × S= NS	

Table 9. Interaction effect of nitrogen levels under different spacing on Vigour Index - II on coriander (*Coriandrum sativum* L.)

Treatments	Vigour Index - II			Mean S
	S ₁ (20 × 15 cm ²)	S ₂ (30 × 15 cm ²)	S ₃ (40 × 15 cm ²)	
N ₀ = Control (0 kg/ha)	252.92	292.17	304.47	283.19
N ₁ = 25 kg/ha	324.88	328.63	339.66	331.06
N ₂ = 50 kg/ha	377.89	410.02	444.60	410.84
N ₃ = 75 kg/ha	393.14	470.27	485.44	449.62
Mean N	337.21	375.27	393.54	
CD (5%)	N= 16.69	S= 14.45	N × S= 28.91	

Table 10. Interaction effect of nitrogen levels under different spacing on Electrical Conductivity on coriander (*Coriandrum sativum* L.)

Treatments	Electrical Conductivity (μS cm ⁻¹ g ⁻¹)			Mean N
	S ₁ (20 × 15 cm)	S ₂ (30 × 15 cm)	S ₃ (40 × 15 cm)	
N ₀ = Control (0 kg/ha)	152	150	149	150
N ₁ = 25 kg/ha	146	142	144	144
N ₂ = 50 kg/ha	132	127	129	129
N ₃ = 75 kg/ha	141	132	138	137
Mean S	143	138	140	
CD (5%)	N= 3.66	S= 3.17	N × S= NS	

similar findings regarding the increase in electrical conductivity of seeds. This increase was attributed to the leaching of electrolytes, nitrogen, and amino acids, which subsequently led to a loss of membrane integrity in aged seeds. Consequently, this reduction in membrane integrity resulted in decreased seed vigor and viability (Table 10).

4. CONCLUSION

The study on the interactive effects of different nitrogen doses and spacing in coriander has provided valuable insights into the optimal conditions for growing this important herb. The results of the study indicate that both nitrogen dose and spacing have a significant impact on

the growth, yield, and quality of coriander. The findings can be useful for farmers and researchers in developing effective strategies for improving the production of coriander, which can have significant economic and nutritional benefits. Further research can be conducted to explore the long-term effects of different nitrogen doses and spacing on coriander and to identify additional factors that may impact its growth and development.

ACKNOWLEDGEMENTS

The authors gratefully acknowledge the Department of Vegetable Science, Chaudhary Charan Singh Haryana Agricultural University, for providing the opportunity to carry out this research work.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Seidemann J. World Spice Plants: Economic, Usage, Botany, Taxonomy. Publisher: Springer-Verlag Berlin Heidelberg. 2005;591.
2. Izgi MN. Effects of nitrogen fertilization on coriander (*Coriandrum sativum* L.): yield and quality characteristics. Applied Ecology and Environmental Research. 2020;18(5):7323-7336.
3. Anonymous. Area production and productivity of coriander in Haryana. Area and production of horticulture crops: All India (NHB, 2018-19); 2019.
4. Gil A, De La Fuente EB, Lenardis AE, López Pereira M, Suárez SA, Bandoni A, et al. Coriander essential oil composition from two genotypes grown in different environmental conditions. Journal of Agriculture and Food Chemistry. 2002; 50:2870-2877.
5. Prachayasittikul V, Prachayasittikul S, Ruchirawat S, Prachayasittikul V. Coriander (*Coriandrum sativum*): A promising functional food for well-being. Food Research International. 2018; 105:305-323.
6. Mandal S, Mandal M. Coriander (*Coriandrum sativum* L.) essential oil: Chemistry and biological activity. Asian Pacific Journal of Tropical Biomedicine. 2015;5(6):421-428.
7. Diwan G, Bisen BP, Maida P. Effect of nitrogen doses and row spacing on growth and seed yield of coriander (*Coriandrum sativum* L.). International Journal of Chemical Studies. 2018;6(4):2768-2772
8. Wangensteen H, Samuelson AB, Malterud KE. Antioxidant activity in extracts from coriander. Food Chemistry. 2004;88:293-297.
9. Maroufi K, Farahani HA, Darvishi HH. Importance of Coriander (*Coriandrum Sativum* L.) between the medicinal and aromatic plants. Advances in Environmental Biology. 2010;4(3):433-436.
10. Ali H, Ayub G, Elahi E, Shahab M, Ahmed S, Ahmed N. Response of coriander (*Coriandrum sativum* L.) to different nitrogen levels and sowing dates. Asian Journal Agriculture Biological. 2015;3(4): 155-158.
11. Javiya PP, Solanki JN, Kaneria SC, Rupareliya VV. Response of coriander (*Coriandrum sativum* L.) to nitrogen and phosphorus in south Saurashtra condition. International Journal of Pure and Applied Bioscience. 2017;5(4):860-866.
12. Katiyar RS, Nainwal RC, Singh D, Chaturvedi V, Tewari SK. Effect of spacing and varieties on growth and yield of coriander (*Coriandrum sativum* L.) on reclaimed sodic waste soil. Progressive Research. 2014;9:811-814.
13. Kaium Abdul, Islam M, Sultana S, Hossain E, Shovon SC, Mahjuba A. Yield and yield contributes of coriander (*Coriandrum sativum* L.) as influenced by spacing and variety. International Journal of Scientific and Research Publications. 2015;5(3): 2250-3153.
14. Choudhary I, Yadav SS, Yadav LR, Sharma OP, Yadav BL. Effect of weed and nitrogen management on coriander (*Coriandrum sativum* L.) yield and economics. Journal of Spices and Aromatic Crops. 2014;23(1):38-44
15. Piper CS. Soil and water analysis. Published by University Adelaide, Australia; 1950.
16. Jackson ML. Soil chemical analysis. Prentice Hall of India Pvt. Ltd., New Delhi. 1973;214-221.
17. Walkley A, Black CA. An examination of the digtjareff method for determination of soil organic matter and a proposed modification of chronic acid titration method. Soil Science. 1934;37:29-39.

18. Subbaiah BV, Asija GL. A rapid procedure for the determination of available nitrogen in the soil. *Current Science*. 1956;25:259-260.
19. Olsen SR, Cole CV, Watanabe FS, Dean LA. Estimation of available phosphorous in soils by extraction with sodium bicarbonate. *Circulation from United States department of agriculture*, 939. USDA, Washington, D.C; 1954.
20. Katar D, Katar N. Determination of the effect of plant density on yield and yield components for two different coriander cultivars (*Coriandrum sativum* L.). *International Journal of Agriculture and Wildlife Science*. 2016;2(1):33-42.
21. Diwan G, Bisen BP, Maida P. Effect of nitrogen doses and row spacing on growth and seed yield of coriander (*Coriandrum sativum* L.). *International Journal of Chemical Studies*. 2018;6(4):2768-2772.
22. Amare G, Mohammed W, Tana T. Effect of plant spacing and np fertilizer levels on growth, seed yield and quality of onion (*Allium cepa* L.) at Shewa Robit, Northern Ethiopia. *The Open Biotechnology Journal*. 2020;14:12-22.
23. Singh Harshita, Panghal VPS, Mor VS, Rana MK. Seed quality of coriander as affected by irrigation and fertilizer levels. *International Journal of Current Microbiology and Applied Sciences*. 2017; 6(12):81-84
24. Kumar KV, Malik TP, Kumar V, Mor VS, Vikram. Effect of integrated nutrient management on seed quality of fennel (*Foeniculum vulgare* Mill.). *Journal of Pharmacognosy and Phytochemistry*. 2019;8(1):511-515.

© 2023 Vats et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/107765>