



# Irrigation Scheduling and Nitrogen Management Practices Affected the Nutrient Concentration of Moth Bean under Arid Ecosystem

R. L. Jat <sup>a\*</sup>, L. K. Rolaniya <sup>a</sup>, Monika Punia <sup>a</sup>  
and Raja Ram Choudhary <sup>b</sup>

<sup>a</sup> ICAR-Indian Institute of Pulses Research, Regional Research Centre, Bikaner-334006, India.

<sup>b</sup> ICAR-Directorate of Groundnut Research, Regional Research Station, Bikaner-334006, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJECC/2023/v13i123778

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/111372>

**Original Research Article**

**Received: 18/10/2023**

**Accepted: 24/12/2023**

**Published: 26/12/2023**

## ABSTRACT

Efficient utilization of water and fertilizers is important for maximizing yield for moth bean production through improvement the nutrient use efficiency under arid ecosystem. The aim of this study was to evaluate the effects of irrigation and nitrogen fertilizers on growth, yield and nutrient content of moth bean in the western zone of Rajasthan during 2020 & 2021. Treatments included three levels of nitrogen (N<sub>1</sub>:15 kg N as basal + foliar spray of urea (2%) at flowering stage, N<sub>2</sub>: 15 kg N + foliar spray of urea (2%) at pod development stage and N<sub>3</sub>: 15 kg N + foliar spray of urea (2%) at flowering and pod development stages) and levels three levels of irrigation (I<sub>1</sub>: single irrigation at branching stage, I<sub>2</sub>: two irrigation at branching + pod development stages and I<sub>3</sub>: two irrigation at flowering + pod development stages) were laid down in factorial combinations of randomized block design with three replications. The results revealed that 15 kg N + foliar spray of urea (2%) at

\*Corresponding author: E-mail: khokharagro1986@gmail.com;

flowering and pod development stage recorded significantly higher nitrogen, phosphorus and potassium content in seed and haulm over 15 kg N + foliar spray of urea (2%) at flowering stage and 15 kg N + foliar spray of urea (2%) at pod development stage. However, the phosphorus content in seed and haulm was not significantly influenced by N<sub>3</sub> and statistically equivalent with N<sub>2</sub> and N<sub>1</sub> during 2020 and 2021. Potassium content in seed showed significant enhancement under N<sub>3</sub>, while K content in haulm was not affected significantly during 2020 and pooled basis. Further, irrigation management practices did not show any significant improvement on N, P, K content in seed and haulm during years of 2020 and 2021. While, N content in seed and haulm was significantly improved under I<sub>2</sub> over I<sub>3</sub> and I<sub>1</sub> on pooled basis. Thus, 15 kg N as basal application + foliar spray of urea (2%) at flowering and pod development stages with two irrigations at branching + pod development stages improved the nutrient content in moth bean in arid condition.

**Keywords:** Nitrogen; irrigation; foliar spray; urea; nutrient content; moth bean.

## 1. INTRODUCTION

Moth bean (*Vigna aconitifolia*) also known as kidney bean, dew bean, dew gram, moth mat, mat bean, is an important leguminous crop cultivated in arid and semi-arid regions, known for its resilience and ability to thrive in low-input agricultural systems. It is native crop of Indian subcontinent and widely cultivated in India, Sri Lanka, China, Pakistan, and also grown in the South-Western states of the USA and in Canada [1,2]. It was grown on acreage of 9.93 lakh hectares of land with annual production of 3.43 lakh tonnes, and average productivity was 346 kg per hectare during the 2020–21 in India. [3]. It is a drought-tolerant pulse crop cultivated for its high protein content and adaptability to adverse growing conditions. *Vigna aconitifolia* is grown from sea-level up to an altitude of 1300 m and it does outstandingly well in arid and hot regions [4,5]. It performs better under average temperature of 24–32°C with a well-distributed annual rainfall of 500–750 mm. The plant provide yield even with as little as 50–60 mm in 3–4 showers during the growing period [5]. Moth bean can be grown on a variety of soils however, it is particularly suitable for dry, light sandy soils and does not tolerate water logging. It is somewhat salt tolerant and can grow on a wide pH range (3.5–10). Due to multipurpose use, it provides hot-season pasture and hay for livestock [6] and immature green pods can be boiled and consumed as a vegetable. At maturity, the seeds can be cooked or fried, and they can be ground and mixed with flour to make dhal and unleavened bread [5]. It is capable of sustaining under long dry spells and high temperatures beyond 40°C. Thus, the erratic and low rains hardly 130 mm with long dry spells of 30-35 days interval between two rains and high temperatures beyond 40°C during crop season affect its growth and productivity drastically often

restricting the productivity much below its potential (>200 Kg/ha). Plants cannot take up all the nitrogen applied to the soil as a fertilizer leading to runoff and groundwater contamination. Thus, nitrogen management in light soils needs careful attention due to high potentials for leaching losses of nitrate with water flow. The sandy soils have low water holding capacity which favors the negative charged nitrate (NO<sub>3</sub><sup>-</sup>) ions to leach in deeper layer below the root zone in light sandy soils. The soil organic matter content in light textured soil is generally lower than fine textured soil. The soil organic matter content is the mains source of nitrogen which supplies nitrogen slowly to the crops as per their demand. The SOM content in soils is space for holding the water and nutrient in the soil and supplying the nutrient to soil solution as per need of the crops. Nitrogen availability is often a limiting factor for high productivity in legume crops. Hence, nitrogen fertilizers are commonly applied to legume crops in countries where they are readily available. The crop yield drastically decreases either under excess water as well as water deficit conditions. Water stress is more common and important in the rain fed arid conditions that reduce the productivity of crops. Water stress during the active crop growth phase results into cessation of growth as it influences the photosynthesis and other physiochemical processes and or death, by desiccation. Irrigation management strategies have become more vital aspect of research in arid region to improve the productivity. The ideal scheduling of irrigation enhances the crop yield though efficient utilization of irrigation water on most critical stages under water stress situation which is depends upon the soil, climate and plant characteristics. Water stress reduces the total chlorophyll and relative water content significantly in different genotypes of moth bean and it is more pronounced in late flowering

genotypes [7]. The overall objective of this work is to improve nutrient content through enhancement of nutrient use efficiency with better agronomic management practices in moth bean. This would be a major advance in profitability for the moth bean industry in India and would significantly improve the ecological footprint of the crop.

## 2. MATERIALS AND METHODS

A field experiment was conducted to evaluate the effect of nitrogen levels and irrigation scheduling on nutrient content (N, P, & K) of moth bean under hyper arid condition of western Rajasthan i.e. Bikaner during kharif 2020 and 2021. The region comes under agro climatic zone “hyper arid partially irrigated western plain” of Rajasthan. It is situated at 28°10' N latitude, 73°18' E longitude and 223.88 meter above mean sea level. The experimental location is characterized by typically tropical arid climate with very hot and dry summer with an average temperature of 40° c and relative humidity of 10-20 % (March-June), cool winter average minimum temperature 4°C and average maximum relative humidity 70% (October to February) and wet monsoon season (July to Sept). The average annual rainfall is 300 mm of which 90% is received in monsoon season (July to Sept). Before the commencement of the experiment, soil sampling was done in a zigzag pattern from a depth of 15 cm. The chemical properties of the soil are as low in organic carbon & nitrogen, medium in phosphorous and high in potassium. The moth bean was sown in lines with spacing 30x10 (30 cm row to row and 10 cm plant to plant) using seed rate of 15 kg ha<sup>-1</sup>. Prior

to sowing, recommended dose of minerals N and P fertilizers were applied as basal application using Urea and Di-ammonium phosphate at rates of 15 kg N/ha and 40 kg P<sub>2</sub>O<sub>5</sub>/ha. Nitrogen applied to all the plot at uniform dose of 15 kg N ha<sup>-1</sup> through DAP as basal application at the time of seeding followed by spray of urea as per the proposed treatment at different growth stages. The size of the plots was 32.4 m<sup>2</sup> (6.0 x 5.4 m) and net plot size for observation was 16.8 m<sup>2</sup> (4.2 x 4 m) maintained with double bunds (1 m wide) boundaries which were compacted to reduce the lateral movement of water. Irrigations was applied in each experimental unit through surface flood irrigation, which involved delivering water to the experimental units by sub water channels from main water channels. Pendimethalin was applied as pre-emergence herbicide followed by hand weeding at 30-35 DAS to control weeds, While, all other agricultural practices were uniform across the experimental treatments except for irrigation. For plant chemical analysis the seed and haulm samples collected at harvest were oven dried at 70°C till constant weight was achieved. The dried samples were finely ground and passed through 40 mesh sieve and used for determination of nutrient content (N, P, K) as per method furnished in Table 1.

A two-way ANOVA analysis was performed using JMP®, Version 16 [11] on observed traits, including nitrogen phosphorus and potassium content in seed and haulm to compare the effect of nitrogen application and irrigation scheduling in moth bean. All effects were statistically significant at the 0.05 level of significance and the results are given below.

**Table 1. Chemical methods for plant nutrient analysis**

S. No.	Nutrient	Method	Reference
1.	Nitrogen	Nessler’s reagent colorimetric method	Lindner [8]
2.	Phosphorus	Ammonium Vanadomolybdo-phosphoric acid yellow colour method	Richards [9]
3	Potassium	Triple acid digestion method	[10]

## 2. RESULTS AND DISCUSSION

### 2.1 Nitrogen Content

**N content in grain :-** Nitrogen treatments significantly affected N content in grain of moth bean (Table 2). The treatment N<sub>3</sub> (15 kg N ha<sup>-1</sup> + 40 kg P<sub>2</sub>O<sub>5</sub> with foliar spray urea @ 2% at flowering and pod development stages) recorded significantly maximum value of N content in seed during both the years of experimentation and pooled basis, which was significantly higher by 7.23, 5.38 & 6.35 and 8.23, 6.32 & 5.26 per cent over N<sub>2</sub> and N<sub>1</sub> during 2020, 2021 and pooled basis, respectively. The

increment in N content might be due to more and continuous availability of nitrogen through soil and foliar application, which is associated with an increase in protoplasm, cell division, and cell enlargement, resulting in higher absorption of nutrients [12,13]. Moreover, significant variation in N content in seed was observed under irrigation scheduling practices. The N content was not enhanced significantly during both the years of experimentation. However, the N content was significantly improved under treatment I<sub>2</sub> (irrigation at branching and pod development stages) on pooled data basis, which was increased by 5.69 percent over I<sub>1</sub> (irrigation at branching stage), however, it was statistically on par with I<sub>3</sub> (irrigation at flowering and pod development stages) on pooled basis. Adequate water management, particularly through irrigation scheduling, plays a pivotal role in optimizing crop yield and nutritional quality. The higher N, P and K content in grain with two irrigations may be due to higher soil moisture content which facilitate the nutrients to bring into soil solution. These results are in agreement with those obtained by Singh et al. [14] and Maske et al. [15].

**N content haulm:** - The N content in haulm was significantly influenced by different nitrogen management levels. The higher value of N content in haulm recorded with N<sub>3</sub> (15 kg N ha<sup>-1</sup> + 40 kg P<sub>2</sub>O<sub>5</sub> with foliar spray urea @ 2% at flowering and pod development stages). The percent increase in the N content was 6.69, 8.52 & 7.74 and 5.41, 7.50 & 6.42 % over N<sub>1</sub> and N<sub>2</sub> during 2020, 2021 and pooled basis, respectively. The N content in haulm increased with higher nitrogen rate might be primarily due to an increase in the chlorophyll content of leaves, which leads to a higher photosynthetic rate, and ultimately plenty of photosynthates available during grain development [16]. Further, irrigation scheduling approaches did not show any significant variations on N content in haulm during both the years of experimentation. While irrigation practices were significantly improved the N content on pooled data basis. The maximum value of N content recorded under treatment I<sub>2</sub> (Irrigation at branching and pod development stages) which was significantly higher by 8.12 percent over I<sub>1</sub> (Irrigation at branching stage). However, it was statistically equivalent with I<sub>3</sub> (irrigation at flowering and pod development stages) on pooled basis. The increase in N content in haulm may be due to more availability of N in soil solution with two irrigations. The observed results are in close conformity with Maske et al. [14].

### 3.2 Phosphorus Content

**P content in grain:** - Nitrogen management levels recorded significant improvement in P content in grain of moth bean (Table 3). The P content in grain was not significantly influenced under different nitrogen management options during both the years of experimentation, however, pooled data shows significant improvement in P content (0.50%) of grain with N<sub>3</sub> (15 kg N ha<sup>-1</sup> + 40 kg P<sub>2</sub>O<sub>5</sub> with foliar spray urea @ 2% at flowering and pod development

stages) over N<sub>2</sub> (15 kg N ha<sup>-1</sup> + 40 kg P<sub>2</sub>O<sub>5</sub> with foliar spray urea @ 2% at pod development stage) and N<sub>1</sub> (15 kg N ha<sup>-1</sup> + 40 kg P<sub>2</sub>O<sub>5</sub> with foliar spray urea @ 2% at flowering stage). Nitrogen, obtained through nitrogen fixation and soil uptake, serves as a catalyst for various metabolic processes that contribute to enhanced phosphorus uptake, transport, and incorporation into seeds. These results are in line with Britz et al. [17]. Further, irrigation management option did not show any significant impact on P content in grain of moth bean during both the years as well as pooled basis.

**P content in haulm:** - Different nitrogen management options were significantly influenced the P content in grain of moth bean (Table 3). On the pooled basis, the treatment N<sub>3</sub> (15 kg N ha<sup>-1</sup> + 40 kg P<sub>2</sub>O<sub>5</sub> with foliar spray urea @ 2% at flowering and pod development stages) was significantly enhanced the P content in haulm by 28.57 and 8.00 percent over N<sub>1</sub> and N<sub>2</sub> respectively. While, P content was not significantly affected in individual year of 2020 and 2021. Adequate nitrogen supply can enhance the overall nutrient status of the plant, potentially improving the uptake and translocation of potassium. This result is consistent with the findings of Coblenz et al. [18] and Xie et al. [19]. Moreover, Irrigation scheduling practices did not express any significant improvement in P content in haulm during both the year as well as pooled basis.

### 3.3 Potassium Content

**K content in grain:** - Significant improvement in K content in grain was recorded under different nitrogen levels applied on moth bean (Table 4). The K content in grain was significantly enhanced under the treatment of N<sub>3</sub> (15 kg N ha<sup>-1</sup> + 40 kg P<sub>2</sub>O<sub>5</sub> with foliar spray urea @ 2% at flowering and pod development stages), which was significantly higher by 23.33, 23.08 &

**Table 2. Effect of nitrogen and irrigation management on nitrogen content in grain and haulm of moth bean**

Treatments	Grain			Haulm		
	2020	2021	Pooled	2020	2021	Pooled
<b>Nitrogen Management</b>						
N <sub>1</sub>	3.29b	3.17b	3.23b	1.90b	1.83b	1.87b
N <sub>2</sub>	3.33b	3.20b	3.27b	1.93ab	1.85b	1.89b
N <sub>3</sub>	3.51a	3.44a	3.48a	2.03a	1.96a	1.99a
<b>Irrigation Management</b>						
I <sub>1</sub>	3.27a	3.14a	3.20b	1.90a	1.83a	1.87b
I <sub>2</sub>	3.49a	3.43a	3.46a	2.01a	1.92a	1.97a
I <sub>3</sub>	3.38a	3.24a	3.31a	1.95a	1.88a	1.92a

**Table 3. Effect of nitrogen and irrigation management on phosphorus content in grain and haulm of moth bean**

Treatments	Grain			Haulm		
	2020	2021	Pooled	2020	2021	Pooled
<b>Nitrogen Management</b>						
N <sub>1</sub>	0.43a	0.39a	0.41b	0.22a	0.20a	0.21b
N <sub>2</sub>	0.47a	0.43a	0.45ab	0.26a	0.24a	0.25a
N <sub>3</sub>	0.51a	0.49a	0.50a	0.28a	0.26a	0.27a
<b>Irrigation Management</b>						
I <sub>1</sub>	0.46a	0.43a	0.45a	0.23a	0.21a	0.22a
I <sub>2</sub>	0.49a	0.47a	0.48a	0.27a	0.25a	0.26a
I <sub>3</sub>	0.45a	0.41a	0.43a	0.25a	0.24a	0.25a

**Table 4. Effect of nitrogen and irrigation management on potassium content in grain and haulm of moth bean**

Treatments	Grain			Haulm		
	2020	2021	Pooled	2020	2021	Pooled
<b>Nitrogen Management</b>						
N <sub>1</sub>	0.30b	0.26b	0.28b	0.71a	0.64b	0.68a
N <sub>2</sub>	0.32b	0.28b	0.30b	0.74a	0.66ab	0.70a
N <sub>3</sub>	0.37a	0.32a	0.35a	0.77a	0.69a	0.73a
<b>Irrigation Management</b>						
I <sub>1</sub>	0.32a	0.28a	0.30a	0.72a	0.65a	0.68a
I <sub>2</sub>	0.35a	0.31a	0.33a	0.77a	0.69a	0.73a
I <sub>3</sub>	0.32a	0.28a	0.30a	0.74a	0.66a	0.70a

28.00% and 15.62, 14.29 & 16.67 % over N<sub>1</sub> an N<sub>2</sub> during 2020, 2021 and pooled basis, respectively. However, N<sub>2</sub> statistically equivalent with N<sub>3</sub> during the 2020 and pooled basis. Adequate nitrogen supply can enhance the overall nutrient status of the plant, potentially improving the uptake and translocation of potassium [8]. Additionally, irrigation management practices were not influenced the K content of moth bean during both the years of experimentations as well as pooled data basis.

**K content in haulm:** - Nitrogen management levels observed the significant increment in the K content in the haulm of moth bean. The treatment N<sub>3</sub> (15 kg N ha<sup>-1</sup> + 40 kg P<sub>2</sub>O<sub>5</sub> with foliar spray urea @ 2% at flowering and pod development stages) registered significantly higher K content in haulm over N<sub>2</sub> an N<sub>1</sub> during 2021. The increment in the K content was 7.81 & 4.54 % higher over N<sub>1</sub> and N<sub>2</sub>, respectively. While, nitrogen levels did not show any significant improvement in the K content during

2020 and pooled basis. In addition, irrigation scheduling approaches were not proved any significant improvement in K content in haulm of moth bean.

#### 4. CONCLUSION

The results of a two-year study validated that irrigation scheduling and nitrogen fertilizer management extensively affected the nutrient concentration of moth bean. Based on the experimental results, it may be concluded that 15 kg N ha<sup>-1</sup> with foliar spray of urea of 2% at flowering and pod development stages in combination with irrigations twice at branching + pod development stages could be more beneficial to improvement in nutrient content of moth bean under arid conditions.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Kochhar SL. Mat or Moth bean [*Phaseolus aconitifolius*]. In: Kochhar (Ed.), Economic botany: A comprehensive study, 5th edition, Cambridge Univ. Press. 2016;430.
2. Munro DB. CNRC; Small, E. Les légumes du Canada. NRC Research Press; 1998. Available:<https://books.google.fr/books?id=BpaUNli-4r4C&pg=PA389>
3. Kanishka RC, Gayacharan Basavaraja T, Chandora R, Rana JC. Moth bean (*Vigna aconitifolia*). A Minor Legume with Major Potential to Address Global Agricultural Challenges. *Front. Plant. Sci.* 2023;14.
4. NARO. NARO Genebank Databases. Genebank Project, NARO, Japan; 2020. Available:[https://www.gene.affrc.go.jp/data/bases-plant\\_images\\_detail\\_en.php](https://www.gene.affrc.go.jp/data/bases-plant_images_detail_en.php)
5. Brink M, Jansen PCM. *Vigna aconitifolia* (Jacq.) Maréchal. In: Brink, M.; Belay, G. (Eds.). PROTA (Plant Resources of Tropical Africa / Ressources végétales de l'Afrique tropicale), Wageningen; 2006.
6. Saravanan M, Ignacimuthu S. Hypocholesterolemic effect of indian medicinal plants - a review. *Medicin. Chem.* 2015;5:1.
7. Garg BK, Burman U, Kathju S. Effect of water stress on moth bean [*Vigna aconitifolia* (JACQ) Marechal] genotypes. *Indian J. Plant Physiol.* 2004;9(1):29-35.
8. Lindner RC. Rapid analysis methods of some or more common inorganic substances of plant and soil. *Plant Physiology.* 1944;19:76-84.
9. Richards LA. Diagnosis and improvement of saline and alkali soils. USDA Handbook No. 60, Oxford and IBH Publishing Co., New Delhi; 1968.
10. Jackson ML. Soil chemical analysis. Prentice Hall of India Pub. Ltd., New Delhi; 1973.
11. JMP®, Version 16. SAS Institute Inc., Cary, NC, 1989–2023.
12. Khan AK, Hussain A, Ganai MA, Sofi NR, Hussain ST. Yield, nutrient uptake and quality of sweet corn as influenced by transplanting dates and nitrogen levels. *Journal of Pharmacognosy and Phytochemistry.* 2018;7(2):3567-3571.
13. Pal V, Singh G, Dhaliwal SS. Agronomic biofortification of chickpea with zinc and iron through the application of zinc and urea. *Commun. Soil Sci. Plant Anal.* 2019;50(15):1864–1877.
14. Singh MV, Kumar N, Srivastava RK. Effect of nutrient management on maize (*Zea mays*) hybrid in eastern zone of Uttar Pradesh. *Annals of Plant and Soil Research.* 2018;20(1):99-102.
15. Maske NM, Pramanik K, Asewar B. Effect of irrigation scheduling and nutrient management on productivity and nutrient uptake by sweet corn (*Zea mays* var. *saccharata*) under vertisols of Marathwada. *Annals of Plant and Soil Research.* 2020;22(3):310-314.
16. El-Sanatawy AM, Ash-Shormillesy SM, El-Yazied AA, El-Gawad HGA, Azab E, Gobouri AA, Sitohy M, Osman A. Enhancing grain yield and nitrogen accumulation in wheat plants grown under a Mediterranean arid environment by foliar spray with papain-released whey peptides. *Agronomy.* 2021;11(10):1913.
17. Britz E, Cyster L, Samuels I, Cupido C, Masemola L, Ngcobo N, Manganyi F, Müller F. Nitrogen fertilization increases the growth and nutritional quality of the forage legume, *Calobota sericea* – A preliminary investigation. *Heliyon.* 2023;9(2):e13535.
18. Coblenz WK, Daniels MB, Gunsaulis JL, Truter JE, Scarbrough DA, Humphry JB, Coffey KP, Pas PA, Moore JR, Teague KA, Spiegth JD. Effects of nitrogen fertilization on phosphorus uptake in Burmudagrass

- forage growth on high soil-test phosphorus sites. Prof. Anim. Sci. 2004;20: 146–154.
19. Xie K, Li X, Zhang Y, Wan L, Hennaway DB, Wang D, Qin Y, Fadul GMA. Effect of nitrogen fertilization on yield, N content, and nitrogen fixation of alfalfa and smooth brome grass grown alone or in mixture in greenhouse pots. J. Integr. Agric. 2015;14:1864–1876.

---

© 2023 Jat et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/111372>