



# Field Efficacy of Insecticides against *Conopomorpha sinensis* Bradley (Lepidoptera: Gracillariidae): A Major Pest of Litchi and its Management in Bihar, India

Swati Suman <sup>a+++\*</sup>, Manendra Kumar <sup>a++</sup>  
and Braj Kishore Prasad Singh <sup>a++</sup>

<sup>a</sup> Department of Zoology, B.R.A Bihar University, Bihar, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJPSS/2024/v36i44470

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/113538>

Original Research Article

Received: 24/12/2023  
Accepted: 27/02/2024  
Published: 04/03/2024

## ABSTRACT

Litchi (*Litchi chinensis* Sonn.) is a subtropical fruit; family Sapindaceae. Litchi offers a good potentiality for its cultivation in Bihar both nutritionally and economically. India is the second largest global producer of litchi, next only to China. Litchi fruit borer (*Conopomorpha sinensis* Bradley, 1986 *Lepidoptera: Gracillariidae*), litchi mite, leaf cutting weevils, litchi looper, and leaf roller are major pests which attack litchi plant, during different seasons. They cause damage to the litchi plant throughout the year but their seasonal abundance not only damage the plant, but also the fruits. Among these pests, *C. sinensis* is the prominent one remaining active throughout the year but it's the peak period was from July to November (as shoot borer) and from March to May (as fruit

<sup>++</sup> Ph.D Scholar;

<sup>\*</sup>Corresponding author: E-mail: swatisuman.12j@gmail.com;

borer). 30-70% of the dropped. The results revealed that the lowest number of fruits infestation was recorded with Lambda Cyhalothrin 5 EC @ 0.003% followed by the treatment with Spinosad which gave better result 25 % infestation was. Therefore, an attempt has been made to study management options for litchi fruit and shoot borer *C. sinensis* by using new control measures.

**Keywords:** Litchi; fruit borer; insecticides; new agents; management; pests; tropical fruits.

## 1. INTRODUCTION

“Litchi (*Litchi chinensis* Sonn.) is well known as queen of fruits, it is considered one of the most important subtropical and tropical evergreen fruits. This fruit is well-liked because of its mouth-watering flavor, eye-catching color, and excellent nutritional content” [1]. “Despite having its origins in China, the litchi fruit is now grown in over 20 nations, with China, India, Thailand and Vietnam serving as its top four producers” Zhao et al., [1] Jana, [2]. “It has gained Geographical Indication (GI) for Muzaffarpur, Bihar. Keeping in view its nutritional and economic importance for the area, factors affecting its production are being looked into by farmers as well as scientists. There are as many as 42 insects’ species and mite pests that have been reported to attack litchi trees and fruits at different stages of growth and fruiting” (Smith and Wagenknecht, [3] Siraj et al., [4]. “However, litchi fruit borer also known as stem end borer, *Conopomorpha sinensis* Bradley (Lepidoptera: Gracillariidae) has gained the status of major pest and it seen now as a significant concern for Indian litchi growers” (Fang et al., [5], Dubey et al., [6], Srivastava et al., 2019). According to Srivastava et al. (2019), “this insect causes significant economic losses to young shoots and fruits that range from 24–48% and 7–70%, respectively”. During the months of April and May, it infiltrates close to the peduncle of fruits (both developing and mature), after hatching, the larvae immediately enter into the fruit, pierce its peduncle and feed on the soft tissue within and damage the aril causing early fruit drop and such fruits lose their nutritional value. If preventative measures are not taken, this insect can cause up to 100% fruit loss in the field, lead to significant losses for Indian litchi growers. Currently, there is lack of information that comprehensively outlines the research on *C. sinensis*, which is very unfavorable for the development of research efforts on *C. sinensis*. Thus, the infestation remains hidden due to its systemic nature, the fact is that *C. sinensis* prevalence is far and wide both from geographical perspective as well as seasonal availability. But the hidden nature of infestation has led to comparatively poor study progress on

the species which is why it is thought to be a research progress limitation. The largest obstacle to developing effective pest control strategies has been our limited understanding of the ecology and behavior of this insect. The seasonality and hidden nature of the infection make it much more challenging to eradicate this insect. The present study was conducted to keep in view these urgencies for control this pest and aim to explore the possibility of *C. sinensis* control by various treatments scheduled on litchi plant.

## 2. MATERIALS AND METHODS

The infestation of litchi fruits by *C. sinensis* was recorded during 2022 and 2023 in a litchi orchard farm located at Mushahari area, Muzaffarpur district, Indian Council of Agriculture Research-National Research Centre. Twenty four trees of litchi orchard of same cultivar (Var. Shahi) were selected randomly. The experiment was laid out in a randomized complete block design with eight treatments, each treatment was replicated thrice. The efficacies of tested compounds; T<sub>1</sub> Spinetoram 11.7 SC @ 1 mL, T<sub>2</sub>, Flubendiamide 39.35 SC @ 1.5 ML, T<sub>3</sub>, Lambda cyhalothrin 5 EC @ 0.003%, T<sub>4</sub>, Flubendiamide 19.92% + Thiacloprid 19.92% @ 0.48%; T<sub>5</sub>, Chlorantraniprole 4.3% SC + Abamectin 1.7 @ 1.5 ml/L SC, T<sub>6</sub> Spirotetramat 11.01% EC + Imidacloprid 11.01 @ 0.36% EC, T<sub>7</sub> Spinosad 45 SC @ 1.75 mL/5L against *C. sinensis* borer were evaluated. Untreated control was also designed by spraying with water only and represented as T<sub>8</sub>. Each treatment and control was replicated for three times. The recommended doses of each insecticide was prepared by dilution with water and then sprayed separately by using a foot pump sprayer. Spraying was done by the insecticides on outer and inner canopy in all directions of trees through the season. Each season, the first spray was given at cardamom size fruit, while second spray after 15 days of first spray (about 15 days before harvest at color break of fruit).




Fig. 1. Depicting the infestation of fruits (%)

### 2.1 Data Collection and Analysis

Hundred ripened fruits were chosen from each replicate randomly and the numbers of healthy and infested fruits were recorded. Collected samples were opened, and damages caused by the *C. sinensis* were ascertained (Fig 1), by the presence of a *C. sinensis* larva inside the fruit or remains of its excreta in the interior. The collected data were statistically analyzed for which SPSS software was used.

### 3. RESULTS AND DISCUSSION

Litchi fruit borer larva is an internal feeder of fruits. During color break stage the egg laying was done on top of fruits near pedicel. The immature larvae bores in to the fruits and reach towards seed passing through pulp for feeding, the black spot near pedicel with small fruit size is an indicator of infestation by litchi fruit borer. It can be concluded that the larvae of this borer cause direct damage to litchi fruit. They can

escape contact with insecticides when they remain inside the fruit but systemic insecticides will be effective. But during their surface existence litchi fruit borer moth can be controlled by contact insecticides. The data in Table 1. pertaining the field efficacy of different pesticides against *C. sinensis* has been plotted. The data showed that treatment of Flubendiamide SC 19.92% and Thiacloprid SC 19.92% @ 0.48% was least effective and recorded highest fruits infestation 55 % followed by 52%, spinetoram11.7 SC @ 1mL/l, spirotetramat SC 11.01% + Imidacloprid SC 11.01% @ 0.36% showed 50 % infestation, while 48% infestation was recorded by Chlorantraniprole SC 4.3% + Abamectin SC 1.7% @ 1.5ml/L ,47% infestation was revealed by Flubendiamide 39.35 SC @ 1.5 ml/5L, 42% fruit infestation was shown by Lambda Cyhalothrin 5 EC @ 0.003%, treatment by Spinosad gave better result 25 % infestation was recorded. The data revealed that the Spinosad 45 SC @ 1.75 ml per liter of water was the most effective in controlling litchi fruit borer. It

Table 1. Depicting treatment and mean number of infested fruits by litchi fruit borer

Treatments	Mean No. of infested fruits by fruit borer	Standard error
T <sub>1</sub>	15.00	6.028
T <sub>2</sub>	12.00	4.726
T <sub>3</sub>	9.67	2.848
T <sub>4</sub>	19.00	5.568
T <sub>5</sub>	16.00	1.155
T <sub>6</sub>	15.67	8.172
T <sub>7</sub>	5.33	1.764
T <sub>8</sub>	19.00	3.464
<b>Total</b>	111.67	-
<b>C.D.</b>	N/A	-
<b>SE (m)</b>	4.34	-
<b>SE (d)</b>	6.14	-
<b>C.V.</b>	53.88	-

was more effective against fruit borer even on long rainy days after application than other insecticides and economically beneficial proved in the present study. So, it may be observed on a large scale in litchi orchards of farmers, and effect analysis needed.

It was moderately effective in reducing the percentage of fruit infestation and increased yield among systemic insecticides. However, its higher application costs reduced the profit margin and showed a lower benefit-cost ratio. These results were in accordance with Purbey [7] reported that spraying systemic insecticides like thiacloprid and imidacloprid is the most effective against LFB. In the current study, Chlorantraniliprole was found effective against *C. sinensis* borer which was also confirmed by various researchers [8]. Upadhyay et al. [9] showed “chlorantraniliprole effectiveness against brinjal shoot and fruit borer” and Larrain et al. (2014) proved “its effectiveness for tomato pinworm *Tuta absoluta* (Meyrick) in tomato”. Moreover, Kiss et al. [10] indicated that “at the injection of chlorantraniliprole in the apple trunk it caused moderate to high mortality of banded leaf roller *Choristoneura rosaceana* (Harrish)”. “Similar result was achieved against the rice stem borer, *Chilo suppressalis* (Walker)” [8]. Chlorantraniliprole is a novel anthranilic diamide insecticide, effective for control of lepidopteran insect pests, as well as some species in the orders coleoptera, Diptera, and Hemiptera. The sub-lethal concentration of Chlorantraniliprole significantly affected on the development of beet armyworm, *Spodoptera exigua* (Hubner) and diamond backmoth, *Plutella xylostella* (Linnaeus). Based on feeding cessation and reduction in feeding damage, chlorantraniliprole and flubendiamide is among the fastest-acting insecticides available for control of Lepidopteran pests.

Our results showed that flubendiamide was also effective against *C. sinensis* with low damage rate in both year. Related to this aspect, Taher et al. [11] demonstrated that “flubendiamide was efficient against yellow stem borer *Sciophaga incertulas* Walker”. Similarly, Shahid et al. (2015) confirmed that “less damage was caused by pod borer, *Helicoverpa armigera* (Hubner) in vegetable pea, legume pod borer, *Maruca vitrata* (Geyer) in pigeon pea, spotted stem borer, *Chilo partellus* (Swinhoe) (Lepidoptera: Crambidae) in maize at the treatment with flubendiamide. Treated sapota with the flubendiamide found that the yield was significantly higher than control against sapota fruit borer, *Phycita erythroliphia*

Hampson. Both chlorantraniliprole and flubendiamide was found effective in managing shoot and fruit borer and suggested that these insecticide could be used in integrated pest management system. The per cent infestation in our result varied from 12 to 85.5% in 2015 whereas it ranged from 15.25 to 88.75%”. Srivastava et al. [12] study was “in line with our study where he found the infestation of the insect up to 59.35% on dropped fruit”.

However, neem based insecticide was not found effective against *C. sinensis*. Similar result was found in legume pod borer, *Maruca vitrata* Fabricius (Lepidoptera: Pyralidae) and fruit borers of guava, *Psidium guajava* Linn. Yule and Srinivasan, [13] reported that “it was found effective against poplar borer, *Saperda calcarate*. It may be effective against stem borer than fruit borer”. Our result suggests either of chlorantraniliprole or flubendiamide could be used to minimize damage and loss caused by *C. sinensis*. Selectivity of the chlorantraniliprole and flubendiamide have more towards the insect ryanodine receptor (Lahm et al., 2009) thus have low mammalian toxicity (Tohnishi 2005; EFSA, 2013). [14-15].

#### 4. CONCLUSION

It appears that treatment by best, but various permutation and combination of different pesticide further should be tried so as to ascertain the least destructive combination which is most effective in controlling the pest and at the same time least destructive to the plant. The pest management against litchi fruit borer was cost effective as the one or two year time is not enough for the treatment to have a better results. It must be treated on larger scale in wide managed area (orchard) to increase the quality and quantity of the fruit.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Zhao L, Wang K, Wang K, Zhu J, Hu Z. Nutrient components, health benefits, and safety of litchi (*Litchi chinensis* Sonn.): A review. *Comprehensive Reviews in Food Science and Food Safety*. 2020;19(4): 2139-2163.

2. Jana BR, Das B, Sharma R, Bhat BP. Performance of late season litchi cultivars grown in Jharkhand Province of India. *Journal of Applied and Natural Science*. 2019;11(1):12-16.
3. Smith EH, Wagenknecht AC. The occurrence of cholinesterase in eggs of the peach tree borer and large milkweed bug and its relationship to the ovicidal action of parathion. *Journal of Economic Entomology*. 1956;49(6):777-783.
4. Siraj M, Singh S, Jindal J. Biology and morphological description of *Conopomorpha sinensis* Bradley (Lepidoptera: Gracillariidae): A serious pest of litchi; 2023. Available: <https://doi.org/10.21203/rs.3.rs-3368329/v1>
5. Fang HH, Lee WL, Chiu KT, Ma HY, Yang SH, Hung CY, Tsai, Y. C. Irradiation with green light at night has great effects on the management of *Conopomorpha sinensis* and maintains favorable litchi fruit quality. *Scientia horticulturae*. 2023;312:111830.
6. Dubey VK, Sahoo SK, Giri GS, Das A. Efficacy of thiamethoxam against whitefly, *Bemisia tabaci* (Gennadius) under open field conditions in okra. *Pest Management in Horticultural Ecosystems*. 2023;29(1):109-115.
7. Purbey SK. Round table conference on lychee borers' jointly organized by the ICAR-National research centre for lychee (NRCL). *Current Science*. 2016;110:758-759.
8. Wang SS, Huang SS, Liang GW, Zeng L. The rearing and the laboratory population life table of litchi fruit borer (*Conopomorpha sinensis* Bradley). *Acta Ecologica Sinica*. 2008;28(2):836-840.
9. Upadhyay SK, Aryal S, Bhusal B, Chaudhary B. Evaluation of Insecticides for the Management of Litchi Fruit and Shoot Borer. *Journal of Nepal Agricultural Research Council*. 2020;6:85-91.
10. Kiss M, Sörös C, Gutermuth Á, Ittész A, Szabó Á. Avermectin Trunk Injections: A Promising Approach for Managing the Walnut Husk Fly (*Rhagoletis completa*). *Horticulturae*. 2023;9(6):655.
11. Taher MA, Rahman MM, Islam KS, Uddin MM. Efficacy of different bagging techniques for the management of litchi fruit borer (*Conopomorpha sinensis*). *Journal of the Bangladesh Agricultural University*. 2022;20(3):238-242.
12. Srivastava K, Purbey SK, Patak RK, Nath, V. Managing fruit-borer for having healthy litchi. *Indian Horticulture*. 2016;61(3):56-63.
13. Yule S, Srinivasan, R. Combining bio-pesticides with chemical pesticides to manage legume pod borer (*Maruca vitrata*) on yard-long bean in Thailand. *International Journal of Pest Management*. 2014;60(1):67-72.
14. Shahid M, Haq E, Mohamed A, Rizvi PQ, Kolanthasamy E. Entomopathogen-based biopesticides: Insights into unraveling their potential in insect pest management. *Frontiers in Microbiology*. 2023;14:1208237.
15. Srivastava K, Patel RK, Gupta AK, Sharma D. Scenario of Insect Pests on Litchi: Management Options. *Innovative Pest Management Approaches for the 21st Century: Harnessing Automated Unmanned Technologies*. 2020;461-480.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/113538>