



Biodegradation and Impact of Polycyclic Aromatic Hydrocarbons in the Environment: A Review

Emmanuel C. Onwujekwe^{a,b*}, James C. Ogbonna^a,
Chibuzor C. Eze^a and Chijioke N. Eze^c

^a Department of Microbiology, University of Nigeria Nsukka, Nigeria.

^b Department of Microbiology, Coal City University Enugu, Nigeria.

^c Department of Science Laboratory Technology, University of Nigeria Nsukka, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JABB/2022/v25i9597

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/92878>

Review Article

Received 30/09/2022
Accepted 02/11/2022
Published 12/12/2022

ABSTRACT

Polycyclic aromatic hydrocarbons are chemical substances that are produced from natural sources such as volcanic eruptions and from anthropogenic sources such as incomplete combustion of fuels such as coal, oil, gas and waste materials. They are mostly implicated among numerous toxic compounds in the ecological system. Exposure of such organic pollutants in an environment can endanger animals, plants, microbial, soil, and aquatic lives. Human beings are at a high risk of developing cancers due to pollutants exposure. The existence of such pollutants in the surroundings can endanger microbial life by affecting cell growth, shape and metabolic process, alters genetic composition and membranes of microbial cells. The major routes of microbial degradation of organic pollutants are via growth and co-metabolism. The processes enable microorganisms to metabolize toxic compounds in the cells milieu, so as to transform contaminants into harmless forms such as carbon dioxide and water. Bioremediation remains the safest and eco-friendly means of converting

*Corresponding author: E-mail: nuel.chijioke@yahoo.com;

toxic environment to a harmless state. This review seeks to provide an overview of biodegradation of polycyclic aromatic hydrocarbons and the adverse effect of polycyclic aromatic hydrocarbons compounds in the environment.

Keywords: *Biodegradation; bioremediation; environmental contaminants; hydrocarbons; microorganism; pollutants degradation.*

1. INTRODUCTION

The environment is witnessing significant pollution arising due to industrialization and other activities of man. Polycyclic aromatic hydrocarbons are chemical substances that are produced from natural sources such as volcanic eruptions and from anthropogenic sources such as partial combustion of fuels such as coal, oil, gas and waste materials. The natural sources of contamination and entry of polycyclic aromatic hydrocarbons in the environment are usually derived from natural sources such as volcanic eruptions and hydrothermal processes [1]. Generally, anthropogenic sources of PAHs contamination are petrogenic and pyrolytic. Petrogenic PAHs are usually derived from crude and refined petroleum, and are usually introduced on aquatic and soil environment through accidental oil spills, pipeline blowouts, improper drainage and discharges from vehicles. Pyrolytic sources are due to incomplete combustion of fuels such as oil, gas and biomass [2]. Polycyclic aromatic hydrocarbons (PAHs) are one of the most extensively disseminated toxic compounds in our surroundings due to its introduction from many anthropogenic sources. Significant proportions of PAHs that are emitted globally are due to human activities such as cooking with firewood, and farming activities such as burning of agricultural wastes. PAHs can as well emanate from other anthropogenic sources such as during manufacturing of products by industries, burning of diesel engines for power generations and combustion of fuels by automobiles. Johnsen et al., [3] noted that substance can form PAH compounds when subjected to heat at high temperature (500-800^oC) to low temperature (100-300^oC) at long duration. When released in surroundings, they portend great ecological and health concern. Choi et al., [4] reported that PAHs are one of the main elements of crude oil. Annually, large quantities of petroleum products are discharged into the soil and water bodies either accidentally or improperly due to mining, transportation, leakages from pipelines and blow outs.

Based on structure, PAHs are compounds with two or more fused benzene rings in its structural

configurations. They are made up of complex aromatic rings in linear, angular, or cluster arrangements [5]. This legacy compounds contaminates the environment and has been considered as emerging pollutants [6]. The contaminants at increased concentrations in environment become toxic to aquatic, plants, animals, microbial and other life forms. Human beings are vulnerable to carcinogenic, teratogenic, and mutagenic hazards when exposed to PAHs [7]. There are over 100 PAHs distributed in nature that usually occur as a complex mixture and not as a single compound [8]. Mixtures of PAHs can be found in incomplete burning of products such as black soot. PAHs can also be manufactured as a single compound for use in research. When found as unadulterated chemicals, they are usually colorless, white, or pale yellow-green solids. They also have a light and pleasing odor.

PAHs are ubiquitous contaminants in nature. Lim et al., [9] noted that they can be found as pollutants in air, sediments, soil and aquatic environments as well as surface and underground waters. The prototypic PAHs such as naphthalene, anthracene and phenanthrene usually serve as indicator compounds for the presence of PAHs in an environment. PAHs in an environment are generally extracted into an organic solvent prior to separation using analytical equipments [1]. Researchers have used numerous analytical instruments/methods in the extraction of PAHs such as gas chromatography (GC), high performance liquid chromatography (HPTLC), Fourier transform infrared spectroscopy (FTIR), supercritical carbon dioxide solvent extraction [9], Tenax extraction [11], and solid phase nonextraction-gas chromatography/mass spectrometry [12].

There are reports of some compounds of PAHs such as anthracene used for the production of the chemotherapeutic agent called Amsarcine [13]. The authors equally noted that Acenaphthene is used as a raw material (dye intermediate) in the production of plastics and as insecticides and fungicides. Fluoranthrene also has applications as a coating material in preserving the internal layer of steel and ductile-

iron drinking water pipes and storage tanks [14]. Organisms are exposed to compounds such as Benzol (a) pyrene (Bap) due to release from industries and automobiles, hazardous wastes dump, biomass burning, waste burning, municipal incinerators, volcanic eruptions and absorption from smoked food. The presence of the various forms of PAHs in the environment causes harmful effects on microorganisms and other life forms. The mutagenic and carcinogenic properties of PAHs are also worrisome. There are indications that polycyclic aromatic hydrocarbons in soil can hinder seed germination, reduced height and leaf area, decrease in total carbohydrates in roots and yield in plants [15]. The time it takes for the degradation of PAHs is determined by factors such as age of the co-existing contaminants in the environment [16]. Some factors that can affect the fate of PAHs in the environment include climate, biological diversity and properties such as molecular structure, polarity, solubility, hydrophobicity, temperature, pH and moisture content [16]. The intervention of degrading microbes unburdens toxicity rates in environment, decontaminates, and makes the environment beneficial for agricultural purposes. Microorganisms as silent operators in the environment have the responsibility to ensure that our surroundings are returned to normal despite the challenges. Microorganisms have developed machineries to surmount harmful nature contaminants. The metabolic expertise of microbes helps them in degrading toxic contaminants. Eze and Orjiakor [17] noted that the distributions of hydrocarbonclastic microorganisms in the environment are beneficial in hydrocarbon removal in the environment.

2. TOXICITY

There are over 100 different PAHs compounds in nature and 16 of these PAHs have been listed as environmental pollutants of main concern as a result of their toxicity, persistent, mutagenic and carcinogenic characteristics. PAHs are characterized by their electrochemical stability, persistency, structural angularity, hydrophobicity, volatility, carcinogenic, and recalcitrance toward biodegradation [16]. The extent of toxicity of PAHs depends on the molecular weight [18]. The low molecular weights (LMW) PAHs are described as acutely toxic and contain two or three aromatic rings. Meanwhile, high molecular weights (HMW) PAHs are regarded as genotoxic and contain four or more rings [19,20]. The high

molecular weight (HMW) PAHs do not dissolve easily in water, they are less explosive and more lipophilic than the low molecular weight (LMW) PAHs [21]. The observed relationship between PAHs and the rising quantity of benzene rings is consistent with the outcome of some findings linking biodegradation time and PAHs molecule size [22].

The main routes of PAHs exposure in humans is from breathing ambient and indoor air, smoking, and consumption of meals containing PAHs. Humans may be exposed to PAHs in their work places with observable symptoms such as irritation of the eyes, confusion, vomiting and nausea. Ingestion of high concentrations of Napthalene can cause haemolysis [23]. Additionally, there are numerous risks associated with exposure to polycyclic aromatic hydrocarbon concentrations in humans such as life time cancer risk, respiratory infections, coughing, impaired lung functions, cardiovascular diseases, eye irritations, stroke, and death [24]. Investigations by Gourdazi et al., [25] showed that air around industrial areas recorded higher levels of PAHs compared to the air surrounding residential homes. Therefore, individuals living close to industrial zones are likely to be exposed to higher concentrations of PAHs. Factors that could determine the effect of PAHs on humans include; concentrations of PAHs, length and route of exposures, toxicity of PAHs, pre-existing health conditions and age of the exposed individuals.

3. EFFECTS OF PAHs ON MICROORGANISMS

As noted earlier, microorganisms can be exposed to PAHs due to emissions from industries and automobiles, hazardous wastes dump, incinerations, biomass burning, waste burning, volcanic eruptions and absorption from smoked food. The pollutants can enter microbial system through absorption, respiration or by contact. There are reports of about fifty-four PAHs that were observed at various hazardous waste sites by the United States department of health and human services. Mumtaz and George [7] reported that concentrations of some indicator PAHs in the atmosphere are about 0.02-1.2 nanograms per cubic in rural areas and 0.15-19.3 ng/m³ in urban areas. The authors equally reported some concentrations of PAHs in potable water range from 4 to 24 nanograms per liter. The observed report shows that developed cities would have higher concentrations of PAHs compared to the rural communities.

Therefore, microorganisms can be exposed to greater concentrations of PAHs in regions with increased concentrations of PAHs. The occurrence of contaminants in high concentrations creates much danger to microbial cells, including the ability of bacteria to attack pollutants. They can upset the microbial populations, either by escalating or diminishing the mass of certain bacterial or fungal populations based on the character of the contaminants [26]. The existence of toxic metals in the ecological system endangers microbial life by affecting cell growth and shape, metabolic processes, alters genetic composition and membranes of microbial cells. Additionally, Eze et al., [27] reported that crude oil pollution inhibited bacterial and fungal counts on the soil and aquatic environment. The authors equally reported reductions in soil nutrients as the level of contamination by crude oil increased. Alimba and Faggio [28] noted that assimilation of contaminants by microorganisms can hinder metabolism and reproductive ability. The presence of contaminants in microorganisms interrupts physiological activities, energy storage, metabolism and survival of microorganisms. Environmental challenges such as the presence of contaminants can cause stress in microorganisms. Numerous environmental stress conditions that could be experienced by microbes include changes in concentration of toxins, carbon source, amino acid limitations, heat shock, acid shock as well as starvations from nutrients such as carbon, nitrogen, and phosphate. In order to stay alive and energetic in the presence of stress, microbes develop physiological adaptation mechanisms [29].

4. ENVIRONMENTAL FATE OF PAHs

The tendency of PAHs to resist biodegradation, with affinity to bioaccumulate and their deleterious effect on microorganisms is a significant concern. Bacteria have been found as principal microorganisms involved in detoxification of toxic compounds. The supply of vastly concentrated and specialized populations of microorganisms improves the ability and speed at which microbial populations degrade and convert toxic compounds [30]. Additionally, the addition of nutrients (biostimulation) that can stimulate autochthonous microorganisms in degrading pollutants is an effective means of detoxifying contaminants. Other environmental factors that could affect degradation of contaminants include oxygen, pH, temperature, moisture content, microbial population and

chemical structure of compounds. The wide range of temperature for PAHs degradation includes 30-40^oC in soil environment, 20-30^oC in fresh water and 15–20^oC in marine environments [16]. The authors equally reported successful degradation rates of PAHs at pH range 4-9. Kaiser [31] stated that microorganisms have the ability to acclimatize in the presence of contaminants found around their system through the process of horizontal gene transfer, so as to utilize such contaminants as nutritional source of carbon and nitrogen. The horizontal gene transfer is critical in the development of bacteria that can mineralize legacy contaminants such as polycyclic aromatic hydrocarbons [32]. Toxic compounds readily undergo bioconversion into less toxic compounds in natural ecosystem. However, processes involving incomplete biotransformation of complex molecules still persist. Some of these compounds are converted into intermediates that are exceedingly recalcitrant to microbial enzymes, thereby making detoxification difficult. Rohini et al., [33] enumerated reasons for the high resistant nature of some compounds in the ecosystem as follows:

1. The inability of biotransforming microorganisms to recognize the compounds as the substrate.
2. They are remarkably stable in ecosystem.
3. They possess large molecular weight which resists entry to microbial cells.
4. Less solubility in water.
5. They have high toxicity and affinity to release toxic products due to microbial actions.

However, the length of time each contaminant persists in the environment varies. It is a unique characteristic of contaminants to possess molecules of unusual elements, groupings and bond as compared biological standards [34]. The unique features make their recognition by naturally occurring microorganisms challenging. It is noteworthy that toxic compounds do not enter the common metabolic pathway of microorganisms easily [35]. PAHs are less soluble in water and can highly adsorb solid particles; this quality makes them to last long in the environment prior to biodegradation. Pannu et al., [36] reported that low molecular weights PAHs are unstable and highly soluble in water and as such highly disposed to microbial utilization.

Transformation of PAHs can take place through processes such as adsorption, volatilization,

photolysis, and chemical oxidation. However, the use of microorganisms for the conversion of contaminants remains the most ecologically sound means for the neutralization process of PAHs contaminated sites. There are two major pathways that could be effectively utilized by microorganisms in the degradation of PAHs via growth and co-metabolism. The growth pathway primarily involves the utilization of organic pollutants as a sole source of carbon and energy. Fritsche and Hofritcher [37] noted that this strategy could bring about mineralization of organic pollutants. Meanwhile, co-metabolism is a means of transformation of organic pollutants by microorganisms without nutritional benefit in the presence of a growth substrate. This mechanism involves microorganisms growing on a particular substrate gratuitously oxidizes a second substrate (co-substrate). Microorganisms do not assimilate the co-substrate, but the product may be available as substrate for other microorganisms of a mixed culture. When pure culture is involved, co-metabolism is a transformation without benefit to the organism. Meanwhile, this initial co-metabolic transformation may pave way for subsequent attack by another organism in the environment [35]. It is the basis of biotransformation used to convert a substance to a chemically modified form. Zhong et al., [38] stated that co-metabolism is an essential means of transforming non-substrate PAHs in the presence of growth substrate to enlarge the range and extent of PAHs degradation.

5. BACTERIAL DEGRADATION OF PAHs

The mineralization of hydrocarbons is essentially a development that relies heavily on the presence of oxygen. Most frequent aerobic degrading microbes are established in bacterial and fungal populations. The primary oxidation of benzene ring by the action of dioxygenase enzymes to form cis-dihydrodiols is the most important procedure for the aerobic bacterial metabolism [39]. The cleavage of the aromatic ring occurs at the rate limiting step as the first enzymatic process during aerobic PAHs degradation. Elliot et al; [40] reported that an essential system for the aerobic bacterial conversion of naphthalene and PAHs is through oxidative actions of the naphthalene dioxygenase enzyme complex which initiates molecular oxygen.

This enzyme complex explicitly integrates an oxygen atom to the two carbon atoms that were

formerly sharing a bond, forming a cis-dihydrodiol (Fig. 1). The intermediates further experience ortho or metaclevavage giving rise to Cis-muconic acid or 2-hydromuconic semialdehyde, respectively [41]. The ring cleavage give rise to the formation of succinic, fumaric, pyruvic, and acetic acids and aldehydes (Fig. 1), these products are precursors for the synthesis of cellular constituents and energy by the microorganisms [42]. Also, there are reports about mobile genetic elements (MGEs) like plasmids and transposon which are useful in the mineralization of organic contaminants such as PAHs [4].

Bacterial populations such as *Sphingomonas*, *Pseudomonas* and *Mycobacterium* were reported as the main microorganisms in PAHs mineralization in soil [43]. *Pseudomonas* strains such as *Pseudomonas sacharophila* P 15, isolated from creosote-polluted soil were observed to have efficiently enhanced the removal rates of fluoranthene, pyrene, benz[a]anthracene, chrysene and benzo[a]pyrene when salicylate was used as inducer of PAHs dioxygenase activity [21,41,44].

Gupta and Pathak [45] also identified the potentials of *Burkholderia* in degrading 2- or 3-ring PAHs such as naphthalene and anthracene, while noting *Mycobacterium* greater efficiency in degrading complex PAHs such as fluoranthene, benzo-alpha-pyrene. Kanaly and Harayama [22] reported on the ability of *Rhodococcus* sp strain UW1 isolated from hydrocarbon-polluted soil. The authors noted their potentials to metabolize Pyrene and Chrysene as sole sources of carbon and energy. Among PAHs degrading bacteria, genus bacteria *Rhodococci* are very distinctive due to their enormous catabolic versatility [4].

Studies by Xu et al., [46] on biodegradation of polycyclic aromatic hydrocarbons by bacterial mixture proved that bacterial mixture increased degradation rates of phenanthrene, anthracene, pyrene and benzo[a]pyrene to over 83%. The authors equally attributed the successful degradation of the compounds to the combined actions of dioxygenase and dehydrogenase enzymes. The findings of the research showed that bacterial consortium would better enhance degradation rates of PAHs than single bacterial.

6. FUNGAL DEGRADATION OF PAHs

Fungal utilization of PAHs is facilitated by the non-ligninolytic and ligninolytic fungi (the white

rot fungi). The non ligninolytic fungi uses cytochrome P450 monooxygenase enzyme which catalyze the oxidation of PAHs to arene oxides, the primary product of PAHs metabolic processes. The arene oxide intermediate is carcinogenic and can further be metabolized by epoxide hydrolase to form a trans-dihydrodiol Fig 1. However, the transdihydrodiol product cannot be used in organisms that use the cytochrome P-450 system as energy source [41,47]. The cytochrome is an essential enzyme in contaminants metabolism and are useful in the pharmaceutical industry due to their effectiveness in breakdown of medications.

The ligninolytic fungi popularly called white rot in wood. These organisms have the ability to absolutely convert PAHs to carbon dioxide and water, thus have made them efficient and extensively used in bioremediation [48]. The ligninolytic fungi have the ability to produce enzymes such as lignin peroxidase (LiP), manganese peroxide (MnP), versatile peroxidase (VP) and laccases (LAC) to degrade lignin in wood [49]. LiP was said to have played a significant role for the initial steps in benzo[a]pyrene metabolism by *P.chrysorium* [47]. The enzyme activity of LiP can be evaluated by the oxidation of veratryl alcohol (VA) which is preferred LiP substrate, to veratradehyde by the increase in absorbance at 310 nm [50].

The degradation of PAHs by MnP was originally studied in *P.chrysorium* and expressed as a lipid peroxidation dependent process [51]. There are also reports that PAHs degradation by some MnPs also takes place directly [47]. The efficiency of MnP in PAHs degradation was shown by Acevedo et al., [52] who reported that MnP produced by *Anthracoophyllum discolor* was able to degrade pyrene (>86%) and anthracene (>65%) alone or in mixture. Meanwhile, the authors noted lesser success in degrading flouranthene (<15.2 %) and phenanthrene (<8.6%).

Meanwhile, reports are sparse on versatile peroxidase production during PAHs degradation by white-rot fungi [53]. Pozdnyakova, [47], noted that PAHs oxidation by versatile peroxidase from *Bjerkandera adusta* in the presence and absence of Manganous ions.

PAHs oxidations by purified fungal laccases have been studied with laccase from *Trametes versicolor*, *Pleurotus ostreatus*, *Coriolus hirsutus*, and *Corioloopsis gallica* [54]. There was

simultaneous reduction of one dioxygen molecule to two molecules of water and oxidizing aromatic substrates by the Laccases [12,55]. The authors equally investigated the efficacy of laccase, produced by *Trametes versicolor* in the removal of aromatic compounds. The removal rate of acenaphthylene was found to have the highest removal rate of 37%, the anthracene and benzo(a)pyrene were 18 and 19%, respectively, and the removal rates for other PAHs investigated were found to be less than 10%. The addition of 1-hydroxybenzotriazole (HBT) to the reaction mixture induced the oxidation of PAHs resulting in virtually total removal of acenaphthylene, acenaphthene, fluorene, anthracene, benzo(a)pyrene, and perylene.

Pozdnyakova [47], reported that LAC from *Corioloopsis gallica* successfully oxidized flouranthene (75%) as well as its polycyclic heterocyclic analogs involving carbazole (100% loss), n-ethylcarbazole, (100%), and dibenothiophene (60%) in the presence of 1-hydroxybenzotriazole (HBT) and 2,2'-azinobis(3-ethylbenzthiazoline-6-sulfonic acid) as free radical mediators.

Ligninolytic fungi such as *Phanerochaete chrysosporium* and *Pleurotus ostreatus* can secrete both ligninolytic and non-ligninolytic type of enzymes, but uncertainties surround the extent each enzymes could take part in the degradation of PAHs [54]. Brown rot fungi are another type of ligninolytic fungi that essentially produces hydrogen peroxide for the degradation of hemicelluloses and cellulose. Brown rot fungi such as *Laetiporus sulphureus* and *Flammulina velutipes* can metabolize PAHs like phenanthrene, flouranthene, and fluorine [49,56].

7. MICROALGAL DEGRADATION OF PAHs

Microalgae are photosynthetic autotrophs which are microscopic and forms the major primary producers in the aquatic environment with applications in wastewater treatment, food and cosmetic industries, and bioenergy production. Its application for wastewater treatment has been subject of study for about a decade ago [57], and some strains have been identified with the potentials to biodegrade PAHs in aquatic environments [4, 58]. The potential of bio removal of fluorene by *Chlorella vulgaris* was studied in BG 11 medium supplemented with different concentrations of fluorine under phototrophic condition [58]. According to the

report, some biodegradable byproducts such as N-Hydroxymethylcarbazol, Dibutyl phthalate, Hexadecanoic acid, ethyl ester, and 1, 2-Benzenedicarboxylic acid, dioctyl ester were identified from the culture medium [58], which formed the proposed degradation pathway in Fig. 2. The fluorine degradation pathway by *C. vulgaris* started with the opening of the aromatic ring possibly by the activities of dioxygenase

enzymes as was obtained in the biotransformation pathway of naphthalene by microalgae *Oscillatoria* sp., strain JCM [59]. Dioxygenase enzyme system is well-known to be in charge for the degradation of PAHs in algal cells [60, 61, 62] just as in bacterial PAHs degradation systems but unlike those of fungi which utilizes monooxygenase systems [60].




Fig. 1. Microbial conversion pathway of PAHs, adapted from Cerniglia [19]

Table 1. Properties of the 16 USEPA PAHs, adapted from Mackay et al., [72]

	Ring number	Molecular weight	Aqueous solubility	Log (K_{ow})	Vapour pressure (Pa)
Naphthalene	1	128	31	3.37	1.1E+2
Acenaphthylene	2	152	16.1	4.00	0.9
Acenaphthene	2	154	3.8	3.92	0.3
Flourene	3	166	1.9	4.18	0.09
Phenanthrene	3	178	1.1	4.57	0.02
Anthracene	3	178	0.0045	4.54	0.001
Pyrene	4	202	0.13	5.18	0.0004
Flouranthene	4	202	0.26	5.22	0.00123
Benzo[a]anthracene	4	228	0.011	5.91	2.80E-5
Chrysene	4	228	0.006	5.91	5.70E-7
Benzo[b]flouranthene	5	252	0.0015	5.80	-
Benzo[k]flouranthene	5	252	0.0008	6.00	5.20E-08
Benzo[a]pyrene	5	252	0.0038	5.91	7.00E.07
Dibenzo[a,b]anthracene	6	278	0.0006	6.75	3.70E-10
Indenol[1,2,3-cd]pyrene	6	276	0.00019	6.50	-
Benzo[ghi]perylene	6	276	0.00026	6.50	1.40E-8




Fig. 2. A schematic overview of the possible biodegradation pathway of fluorene by *C. vulgaris* [58]




Fig. 3. Different strategies for biodegradation of PAHs by microalgae

A number of strategies that could be effectively employed in removal of PAHs include 1. the use of different microalga species; 2. use of mixture of PAHs, and 3. use of microalgae-bacteria consortium (Fig. 3). In a study on the mineralization competency for either fluoranthene or pyrene, or a combination of fluoranthene and pyrene by some microalgae species, it was observed that the existence of individual PAH acts synergistically in the elimination of the other PAH [63]. Hong et al. [64] observed that the degradation of fluoranthene by the two diatoms (*Skeletonema costatum* and *Nitzschia* sp.) were slower, compared to phenanthrene, and that the strains also showed comparable or greater competence in the elimination of the Phn-Fla combination than individual Phn or Fla. A report on the removal of benzo(a)pyrene (BaP) by two microalgal species *Selenastrum capricornutum* and *Scenedesmus acutus* revealed that *S. capricornutum* can remove 99% of BaP after 15h of exposure, whereas *S. acutus* can remove 95% after 72h of exposure under the same condition [65]. According to [66,67], the ability to neutralize PAHs from the ecological system and the efficacy of microbial cell population relies mainly on PAHs involved. In another study, *C. sorokiniana* and bacterial consortium was reported to have satisfactorily removed PAHs (> 85%) more than the algal system only which indicated the synergistic relationship between the algal-bacterial consortia [68]. Furthermore, algal-bacterial consortium was reported for the degradation of phenanthrene [69] and pyrene under photosynthetic conditions. It is cost-effective to use microbial consortium as they have proven to be competent and essential [70].

The emergence of biorefinery concept using microalgae feedstock highlights the economic advantage of PAHs bioremediation by microalgae. The microalgae biomass recovered from the process can be converted to biofuel and other value-added products of commercial value [71].

Additionally, there are numerous advantages in the use of various bioremediation techniques in degradation of hydrocarbon compounds such as; it is an environmentally friendly technique with a public acceptance. It converts recalcitrant compounds into harmless products such as biomass, carbon dioxide and water. Bioremediation technologies are relatively cost effective when compared to other remediation strategies. Microorganisms involved in

bioremediation processes increases exponentially and decrease naturally during remediation process thereby assisting in nutrient cycling compared to the use of chemical additives which may persist in the environment.

Therefore, this study encourages exploit of alternative sources of materials such as use of bio natural gas other than fossil fuels and means that introduces PAHs in the ecological system. Adequate environmental conditions that support degradation such as moisture, temperature, nutrient content and availability, pH, and use of proven strains of microorganisms with enzyme ability for degrading polycyclic aromatic hydrocarbons should be applied effectively to encourage complete biodegradation processes.

8. CONCLUSION

This review discussed the biodegradation and impact of polycyclic aromatic hydrocarbons compounds in the natural environment. PAHs have deleterious effect in surroundings because of its toxic nature and the time it takes prior to biodegradation. The pollutant affects the environment negatively by exposing humans to health risks, diminishes microbial populations, reduces soil quality and nutrients, and destroys aquatic lives. Microorganisms are excellent tools for the biodegradation of toxic compounds in nature. Microbial ability to feed on pollutants in an environment remains a key to the bioconversion of chemicals into harmless forms such as biomass, carbon dioxide and water. The use of eco-friendly technologies rather than means that introduces PAHs in an environment will be helpful in diminishing concentrations of pollutants in the environment. It is expected that the current research will open up more research in biodegradation of pollutants especially PAHs, as to find novel microorganisms and enzymes that can effectively degrade pollutants.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Lawal AT, Fantke P. Polycyclic aromatic hydrocarbons: A review. Cogent Environmental Science. 2017;3(1).
2. Kafilzadeh F. Distribution and sources of polycyclic aromatic hydrocarbons in water and sediments of the Soltan Abad River,

- Iran. The Egyptian Journal of Aquatic Research. 2015;41 (3):136-164.
3. Johnsen AR, Wickly, Harms H. Principle of microbial PAH-degradation in soil. Environmental pollution 2005;133(1):71-84.
 4. Choi B, Lee S, Jho EH. Removal of TPH, UCM, PAHs, and Alk-PAHs in oil contaminated soil by thermal desorption. Applied Biological Chemistry. 2020;63(83).
 5. Ghosal D, Ghosh S, Dutta TK, Ahn Y. Current State of knowledge in microbial degradation of Polycyclic Aromatic Hydrocarbons (PAHs): A review. Frontier Microbiology. 2016;7:13-69.
 6. Abdulrazaq A, Abdulsalam A, Rotimi AL, Abdulbasit AA, et al. Classification, potential routes and risk of emerging contaminants/pollutants. Intech Open; 2020.
DOI: 10.5772/intechopen.94447
 7. Domingo JL, Nadal M. Human dietary exposure to polycyclic aromatic hydrocarbons: A review of the scientific literature. Food Chemistry and Toxicology. 2015;86:144-153.
 8. Mumtaz M, George J. Toxicological profile for polycyclic aromatic hydrocarbons, U.S. Department of health and human services, Agency for Toxic Substances; 1995.
 9. Lim H, Sadiqsis L, Galvao M, Westerholm R, Dreij K. Polycyclic aromatic compounds in particulate matter and indoor dust at preschools in Stockholm, Sweden: Occurrence, sources and genotoxic potential in vitro. Science of the Total Environment. 2021;755(1):142709.
 10. Laitinen A, Michaux A, Aaltonen O. Soil cleaning by carbon dioxide extraction - A review. Environmental Technology. 1994; 15:715-727.
 11. Hong S, Yim UH, Ha SY, Shim WJ, Jeon S, Lee S, Khim JS. Bioaccessibility of AhR-active PAHs in sediments contaminated by the Hebei spirit oil spill: Application of Tenax extraction in effect-direct analysis. Chemosphere. 2016;144:706-712.
 12. Wilson SC, Hewitt U, Miller M, Campiglia AD. Water analysis of the sixteen environmental protection agency-polycyclic aromatic hydrocarbons via solid-phase nanoextraction – gas chromatography/mass spectrometry. Journal of Chromatography A. 2014;1-8.
 13. Brazkova M, Krastanov A. Polycyclic aromatic hydrocarbons: Sources, effect, and biodegradation in Biotechnologies and food technologies conference. 2013; (52):10.2.
 14. Mumtaz M, George J. Toxicological profile for polycyclic aromatic hydrocarbons, U.S. Department of health and human services. Agency for Toxic Substances; 1997.
 15. Ahmed F, Fakhruddin AN. A review on environmental contamination of petroleum hydrocarbons and its biodegradation. International Journal of Environmental Sciences & Natural Resources. 2018; 11(3):1-7.
 16. Gupte A, Tripathi A, Patel H, Rudakiya D, Gupte S. Bioremediation of polycyclic aromatic hydrocarbons (PAHs). A perspective. Open Biotechnology Journal; 2019.
 17. Eze CN, Eze EA, Okobo UJ. Effect of crude oil-pollution on soil and aquatic bacteria and fungi. Journal of Experimental Biology and Agricultural Sciences. 2020a; 8(2):176-184.
 18. Mackay D, Calcott D. "Partitioning and physical chemical properties of PAHs" in the Handbook of Environmental Chemistry, ed. Nelson AH. (Berlin Springer-verlag). 1998;325-346.
 19. Cerniglia CE. Biodegradation of polycyclic aromatic hydrocarbons. Biodegradation. 1992;3(2-3): 351-368.
 20. Abdel-Shafy HI, Mansour MS. A review on polycyclic aromatic hydrocarbons: source, environmental impact, effect on human health and remediation. Egyptian Journal of Petroleum. 2016;25(1):107-123.
 21. Edokpayi JN, Odiyo JO, Popoola OE, Msagati TA. Determination and distribution of polycyclic aromatic hydrocarbons in Rivers, sediments and waste water effluents in Vhembe district, South Africa. Int J Environ Res Public Health. 2016; 13(4):387.
 22. Kanaly RA, Harayama S. Biodegradation of high-molecular weight polycyclic aromatic hydrocarbons by bacteria. J Bacteriol. 2000;182(8):2059-67.
 23. Rahman MM, Rahim A, Mowla SG, Chowdhury FR. Severe haemolytic anaemia due to ingestion of Naphtalene containing oil. JCPSP. 2012;22(11):740-1.
 24. Mammi-Galani E, Chalvatzaki E, Lazaridis M. Personal exposure and dose of inhaled ambient particulate matter bound metals in five European cities. Aerosol Air Qual Res. 2016;16(6):1452-63.
 25. Goudarzi G, Alavi N, Babaei AA, Geravandi S, Idani E, Salmanzadeh S et

- al. Investigation of ambient polycyclic aromatic hydrocarbons in a populated Middle Eastern city. *Polycyclic Aromat Compd.* 2020;23:1-6.
26. Lu K, Mahbub R, Fox JG. Xenobiotics interaction with the intestinal microflora. *ILAR J.* 2015;56(2):218-27.
27. Eze CN, Orjiakor PI, Ebeifenadi UJ. Changes in soil microbial respiration and physicochemical properties following Bonny light crude oil contamination of sandy loam soil. *IFE J Sci.* 2020b; 22(1):153-63.
28. Alimba CG, Faggio C. Microplastics in the marine environmental pollution and mechanisms of toxicological profile. *Environ Toxicol Pharmacology.* 2019;68:61-74.
29. Schimel J, Balsler TC, Wallenstein M. Microbial stress-response physiology and its implications for ecosystem function. *Ecology.* 2007;88(6):1386-94.
30. Növak H. Influence of qPCR workflow on target gene enumeration from environmental samples in the case of bioremediation potential estimation [doctoral dissertation]; 2012.
31. Kaiser G. Horizontal gene transfer in Bacteria in [bio.libretexts.org-Libretexts_Biology](https://www.libretexts.org); 2022.
32. McGowan C, Fulthorpe R, Wright A, Tiedje JM. Evidence for interspecies gene transfer in the evolution of 2,4-dichlorophenoxyacetic acid degraders. *Appl Environ Microbiol.* 1998;64(10):4089-92.
33. Bansode R, Lal Chand Malav SK, Mahesh Kumar Malav. *Xenobiotic Compounds and Its Types*; 2015. Available:<https://www.researchgate.net/publication/275715148>.
34. UNEP. *Microbial degradation of xenobiotics.* United Nations Environment Programme; 1990.
35. Kumar P. Major Metabolites after degradation of xenobiotics and enzymes involved in these pathways; 2019. Available: www.science. Available from: direct.com.
36. Pannu JK, Singh A, Ward OP. Influence of peatnut oil on microbial degradation of polycyclic aromatic hydrocarbons. *Can J Microbiol.* 2003;49(8):508-13.
37. Fritsche W, Hofrichter M. *Aerobic degradation by microorganisms.* Biotechnology set. 2nd ed. 2008; 144-67.
38. Zhong H, Zeng G, GM, Yuan X, Lan C, Tam NF. Influence of growth medium on cometabolic degradation of polycyclic aromatic hydrocarbons by phingomonas strain PheB4. *Appl Microbiol Biotechnol.* 2007;7(1):175-86.
39. Sefidi-Heris Y, Hajizadeh N. Bacterial biodegradation of phenolics hydrocarbons in. *Microorganisms for Sustainability.* 2022; 139-62.
40. Elliot R, Singhal N, Swift S. Surfactants and bacterial bioremediation of polycyclic aromatic hydrocarbon contaminated soil—unlocking the targets. *Crit Rev Environ Sci Technol.* 2010;41(1):78-124.
41. Bezza FA. Biosurfactant assisted bioremediation of petroleum and polycyclic aromatic hydrocarbons in aquatic and soil media [doctoral dissertation]. 2016;10-28: 28-34.
42. Juhasz AL, Naidu R. Bioremediation of high molecular weight polycyclicaromatic hydrocarbons: a review of the microbial degradation of benzo[a]pyrene. *Int Biodeterior Biodegrad.* 2000;45(1-2):57-88.
43. Bisht S, Pandey P, Bhargava B, Sharma S, Kumar V, Sharma KD. Bioremediation of polyaromatic hydrocarbons (PAHs) using rhizosphere technology. *Braz J Microbiol.* 2015;46(1):7-21.
44. Chen SH, Aitken MD. Salicylate stimulates the degradation of high molecular weight polycyclic aromatic hydrocarbons by *Pseudomonas saccharophila* P15. *Environ Sci Technol.* 1999;33(3):435-9.
45. Gupta S, Pathak B. Mycoremediation of polycyclic aromatic hydrocarbons. In: *Abatement of environmental pollutants; trends and strategies.* 2012;127-49.
46. Xu M, Wu M, Zhang Y, Zhang H, Liu W, Chen G et al. Biodegradation of polycyclic aromatic hydrocarbons (PAHs) by bacterial mixture. *Int J Environ Sci. Technol.* 2022; 19(5):3833-44.
47. Pozdnyakova NN. Involvement of the liginolytic system of white rot and litter decomposing fungi in the degradation of polycyclic aromatic hydrocarbons. *Biotechnol Res Int.* 2012;2012:243217.
48. Cerniglia CE. Fungal metabolism of polycyclicaromatic hydrocarbons: past, present, and future applications in bioremediation. *J Ind Microbiol Biotechnol.* 1997;19(5-6):324-33.
49. Cerniglia CE, Sutherland JB. Degradation of polycyclic aromatic hydrocarbons by fungi. In: *Timmis KN, McGenity TJ, van der*

- meer JR, de Lorenzo V, editors (Berlin Springer), Handbook of hydrocarbon and lipid microbiology.2010;2080-110.
50. Wong DWS. Structure and action of ligninolytic enzymes. Appl Biochem Biotechnol. 2009;157(2):174-209.
 51. Moen MA, Hammel KE. Lipid peroxidation by the manganese peroxidase of *Phanerochaete chrysosporium* is the basis of for phenanthrene oxidation by the intact fungus. Appl Environ Microbiol. 1994;60(6):1956-61.
 52. Acevedo F, Pizzul L, Castillo MD, González ME, Cea M, Gianfreda L et al. Degradation of polycyclic aromatic hydrocarbons by free and nanoclay-immobilized manganese peroxidase from *Anthracozyllum discolor*. Chemosphere. 2010;80(3):271-8.
 53. Nikiforova SV, Pozdnyakova NN, Makarov OE, Chernyshova MP, Turkovskaya OV. Chrysene bioconversion by the white rot fungus *Pleurotus ostreatus*. Microbiology. 2010;79(4):456-60.
 54. Bezalel L, Harder Y, Cerniglia CE. Enzymatic mechanism involved in phenanthrene degradation by white rot fungus *Pleurotus ostreatus*. Appl Environ Microbiol. 1997;63:2495-501.
 55. Bamforth SM, Singleton I. Bioremediation of polycyclic aromatic hydrocarbons: current Knowledge and future directions. J Chem Technol Biotechnol. 2005;80(7):723-36.
 56. Sack U, Heinze TM, Deck J, Cerniglia CE, Martens R, Zadrazil F, et al. Comparison of phenanthrene and degradation by different wood decaying fungi. Appl Environ Microbiol. 1997;63(10):3919-25.
 57. Oswald WJ. The role of micro algae in liquid waste treatment and reclamation. In: Lembi CA, Waaland JR, editors Algae and human affairs. Cambridge: Cambridge University Press. 1998;225-81.
 58. Asghari S, Rajabi F, Tarrahi R, Salehi-Lisar SY, Asnaashari S, Omid Y et al. Potential of the green microalga *Chlorella vulgaris* to fight against fluorene contamination: Evaluation of antioxidant systems and identification of intermediate biodegradation compounds. J Appl Phycol. 2020;32(1):411-9.
 59. Cerniglia CE, Baalen CV, Gibson DT. Metabolism of naphthalene by cyanobacterium *Oscillatoria* sp. Strain. JCM. J Gen Microbiol. 1980;116:485-94.
 60. Warshawsky D, Cody T, Radike M, Reilman R, Schumann B, LaDow K et al. Biotransformation of benzo [a] pyrene and other polycyclic aromatic hydrocarbons and heterocyclic analogs by several green algae and other algal species under gold and white light. Chem Biol Interact. 1995;97(2):131-48.
 61. Haritash AK, Kaushik CP. Biodegradation aspects of polycyclic aromatic hydrocarbons (PAHs): a review. J Hazard Mater. 2009;169(1-3):1-15.
 62. Patel JG, Kumar JIN, Kumar RN, Khan SR. Biodegradation capability and enzymatic variation of potentially hazardous polycyclic aromatic hydrocarbons—anthracene and pyrene by *Anabaena fertilissima*. Polycyclic Aromat Compd. 2016;36(1):72-87.
 63. Lei AP, Hu ZL, Wong YS, Tam NF. Removal of fluoranthene and pyrene by different microalgal species. Bioresour Technol. 2007;98(2):273-80.
 64. Hong S, Yim UH, Ha SY, Shim WJ, Jeon S, Lee S et al. Bioaccessibility of AhR-active PAHs in sediments contaminated by the Hebei spirit oil spill: application of Tenax extraction in effect-direct analysis. Chemosphere. 2016;144:706-12.
 65. GarcideLasera MP, Olmos-Espejel Jde J, Diaz-Flores G, Montano-Montiel A. Biodegradation of benzo(a)pyrene by two freshwater microalgae *Selenastrum capricornutum* and *Scenedesmus acutus*: a comparative study useful for bioremediation. Environ Sci Pollut Resour International. 2016;23:3365-75.
 66. Luo L, Wang P, Lin L, Luan T, Ke L, Tam NFY. Removal and transformation of high molecular weight polycyclic aromatic hydrocarbons in water by live and dead microalgae. Process Biochem. 2014;49(10):1723-32.
 67. Ke L, Luo L, Wang P, Luan T, Tam NF. Effects of metals on biosorption and biodegradation of mixed polycyclic aromatic hydrocarbons by a freshwater green alga *Selenastrum capricornutum*. Bioresour Technol. 2010;101(18):6961-72.
 68. Borde X, Guiesse B, Delgado O, Munoz R, Kaul R, H et al. Synergistic relationships in algal-bacterial microcosms for treatment of aromatic plants. 2003;86:293-300.
 69. Muñoz R, Guiesse B, Mattiasson B. Phenanthrene biodegradation by an algal-bacterial consortium in two-phase

- partitioning bioreactors. Appl Microbiol Biotechnol. 2003;61(3):261-7.
70. Gupta G, Kumar V, Pal AK. Microbial degradation of high molecular weight polycyclic aromatic hydrocarbons with emphasis on pyrene. Polycyclic Aromat Compd. 2019;39(2):124-38.
71. Koyande AK, Chew KW, Rambabu K, Tao Y, Chu D-T, Show P-L. Microalgae: A potential alternative to health supplementation for humans. Food Sci. Hum Wellness. 2019;8(1):16-24.
72. Mackay D, Calcott D. Partitioning and physical chemical properties of PAHs. In: the Handbook of Environmental Chemistry Nelson AH, editor. Berlin: Springer-verlag. 1998;325-45.

© 2022 Onwujekwe et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/92878>