



Neonatal Jaundice: Knowledge, Attitude and Practice among Pregnant Women Attending the Antenatal Clinic of Rivers State University Teaching Hospital, Nigeria

Boma Awoala West ^{a,b*}, Josephine Enekele Aitafo ^{a,b}
and Basil Omiebi Altraide ^{c,d}

^a Department of Paediatrics, Rivers State University Teaching Hospital, Nigeria.

^b Department of Paediatrics and Child Health, Rivers State University, Nkpolu, Port Harcourt, Nigeria.

^c Department of Obstetrics and Gynaecology, Rivers State University Teaching Hospital, Nigeria.

^d Department of Obstetrics and Gynaecology, Rivers State University, Nkpolu, Port Harcourt, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJTDH/2021/v42i2330559

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/83226>

Original Research Article

Received 21 October 2021

Accepted 25 December 2021

Published 27 December 2021

ABSTRACT

Aim: This study was undertaken to assess the knowledge, attitude and practice towards neonatal jaundice among pregnant women.

Study Design: A cross-sectional questionnaire-based study.

Place and Duration of Study: The study was carried out in the antenatal clinic of the Rivers State University Teaching Hospital from 1st March to 30th June 2021.

Results: Majority of the pregnant women were within the age group 26-34years 92(61.7%), resided in urban areas 117(78.5%) and from South-south geo-political zone 91(61.1%). They were mostly multiparous 65(43.6%) with tertiary level of education 100(67.1%). More than half 82(55.0%) of the pregnant women had heard of neonatal jaundice. Majority 131(89.3%) did not know the causes of neonatal jaundice, did not know the area of the body jaundice appeared first 143(96.0%), did not know the complications of neonatal jaundice 106(71.1%), did not know danger signs of neonatal jaundice 141(94.6%) and none knew any preventive method. Majority of the pregnant women

132(88.6%) would visit the hospital if their babies developed jaundice and 131(87.9%) would consent to blood transfusion if the need arose. Only 25(16.8%) pregnant women had had child(ren) with neonatal jaundice, of which 20(80.0%) took child(ren) to the hospital and 12(60.0%) presented to the hospital within 24hours. Only 1(0.7%) woman had good knowledge level, 120(80.5%) had good attitude and 12(8.1%) had good practice levels towards neonatal jaundice. There was statistically significant association between knowledge level and attitude (P value=.003) and practice levels (P value=.001).

Conclusion: Pregnant women attending the antenatal clinic of Rivers State University Teaching Hospital had insufficient knowledge and poor practice but good attitude towards neonatal jaundice. Emphasis should therefore be placed on health education especially during antenatal period on neonatal jaundice and other prevailing health issues as this would improve their knowledge level and thus their attitude and practice.

Keywords: Neonatal jaundice; knowledge; attitude; practice; pregnant women.

1. INTRODUCTION

Neonatal jaundice (NNJ) or hyperbilirubinaemia is the yellowish colouration of the sclera, skin or mucous membranes of the neonate due to the accumulation of bilirubin in the blood [1,2]. It is a common phenomenon, with an incidence rate of 60% - 80% reported, the lower incidence occurring in term babies and the higher incidence in preterm babies [1,2,3]. About 75% of neonatal admissions have been attributed to NNJ [2,4] as well as readmissions after discharge from birth hospitalizations [5,6]. The Nigerian Society of Neonatal Medicine (NISONM) also estimated that hyperbilirubinaemia accounted for 1 out of 5 neonatal admissions and at least 5% of all neonatal mortalities in Nigeria [5]. A previous study carried out in our centre also showed a high prevalence of NNJ of 34% [7].

Bilirubin is the product of the breakdown of red blood cells. These are frequently broken down as a physiologic response. Jaundice becomes visible in the neonate when the total serum bilirubin level reaches more than/or equal to 85 $\mu\text{mol/L}$ [8]. Hyperbilirubinaemia thus occurs due to the massive breakdown of red blood cells, immaturity of the liver, reduced bilirubin excretion from the liver and increased absorption/circulation of bilirubin from the intestine. It may be conjugated or unconjugated, the latter being the most encountered [9] and may be physiologic or pathologic depending on the aetiology. Some of the risk factors of NNJ are sepsis, prematurity, maternal diabetes mellitus, race, drugs, cephalhaematoma, breastfeeding, blood group incompatibilities between the baby and mother, enzyme deficiencies such as G6PD deficiency, male sex, induction of labour with oxytocin etc [2,4,8].

Jaundice, a physical sign observed during the neonatal period is an important cause of neonatal morbidity and mortality [10,11]. The unconjugated (indirect) bilirubin is neurotoxic and the reason for the morbidity associated with hyperbilirubinaemia.

Two main modalities of treatment of NNJ includes exposure of the baby to ultraviolet light (phototherapy) in mild to moderate cases and exchange blood transfusion in severe cases. Delay in the treatment of NNJ could lead to acute bilirubin encephalopathy(kernicterus) resulting from bilirubin staining of the basal ganglia with diffuse neuronal damage which could lead to long term neurologic sequelae such as cerebral palsy, seizure disorders, hearing impairments etc and even death [8,12,13]. Prevention of these deleterious effects of hyperbilirubinaemia calls for early recognition and prompt treatment and thus early recourse to the health care facility [4,8].

It is pertinent to note that an adequate knowledge, right attitude, and practice on the part of mothers to neonatal jaundice will help in mitigating the scourge of neonatal jaundice. Early postnatal discharges from the hospital of apparently healthy neonates could also contribute to the menace of NNJ thus mothers should be equipped with the skill of recognition of jaundice in their babies so as to seek prompt medical attention.

Igboanus et al. [14] in their community study in Northern Nigeria among women of reproductive age reported poor knowledge in 91% and poor attitude to NNJ in 62% of the women studied. This poor knowledge and attitude towards NNJ were similarly observed by other researchers [2,15,16]. It is therefore imperative that mothers

have good knowledge and attitude towards NNJ so as to reduce its morbidity and mortality.

No study of this nature has been carried out in the Rivers State University Teaching Hospital therefore the above study was done to assess the knowledge, attitude and practice towards NNJ of pregnant mothers attending the antenatal clinic. This would generate evidence-based information that will help formulate policies that would improve maternal knowledge, their attitude and practices thereby improve neonatal outcomes.

2. MATERIALS AND METHODS

This was a cross-sectional questionnaire-based study carried out over a 4-months period from 1st March to 30th June 2021 in the antenatal clinic (ANC) of the Rivers State University Teaching Hospital (RSUTH). The ANC which is a unit in the Department of Obstetrics and Gynaecology sees an average of 60 patients per clinic day and it is run by consultants, resident doctors and house-officers every Mondays to Fridays while nurses give health talks each clinic day before commencement of the clinic consultations.

The Rivers State University Teaching Hospital is a tertiary hospital owned by the Rivers State Government and receives referral from Primary Healthcare Centres and General Hospitals in all the 23 Local Government areas as well as from private health facilities and neighbouring states. The RSUTH is comprised of the Department of Obstetrics and Gynaecology in addition to other Departments in the hospital such as Paediatrics, Surgery, Internal medicine, Family medicine, Radiology, Pathology, Pharmacy, Nursing etc.

Before commencement of the study, a research team was formed comprising of 3 research assistants who were house officers. A structured training of the research assistants was carried out by the lead researcher on the proper administration of the pre-tested and validated questionnaire including inclusion and exclusion criteria. Inclusion criteria were all pregnant women who gave consent to partake in the study while the exclusion criteria were pregnant women who did not give consent.

The study was explicitly explained to the pregnant women attending the ANC and an informed consent obtained. An average of 3-4 pregnant women who gave consent were recruited each day in 3 out of the 5 clinic days.

A convenient sampling size of 149 pregnant women were recruited.

The proforma was administered to every pregnant woman recruited. The information obtained included the socio-demographic characteristics of the participants including the parity, questions which assessed the knowledge of NNJ, attitude and practice towards NNJ. Eight items were used to assess the knowledge of NNJ. Correct answers were scored 1 each and wrong answers were scored 0 and then summed and converted to percentages with a total of 100%. Assessments were then graded good knowledge (70-100%), fair knowledge (50-69.9%) and poor knowledge (0-49.9%). A total of 2 items were used to assess the attitude towards NNJ. A correct answer was scored 1 while wrong answer 0. All items were summed and converted to a total of 100%. Assessments were graded as good attitude ($\geq 50\%$) and poor attitude (0-49.9%). Two items were used to assess the practice of participants towards NNJ. Correct answer was scored 1 and wrong answer 0. All items were summed and converted to 100% and graded as good practice ($\geq 50\%$) and poor attitude (0-49.9%). The reliability test score (Cronbach alpha) for the scale used for assessing the knowledge, attitude, and practice of mothers towards NNJ were 0.500, 0.832 and 0.846 respectively. The reliability coefficient (r) for knowledge was below the required coefficient value of 0.7 but good for attitude and practice.

All the information obtained was entered into an excel spreadsheet and data analysed using SPSS version 23. The data was presented as frequency, percentages and bar chart for categorical variables and as means and standard deviations for continuous variables. Fischer's Exact test was used to test for association while Logistics regression was carried out to test for the strength of the relationship between the variables. Statistical significance was at P value $< .05$ at 95% confidence interval.

3. RESULTS

3.1 Socio-Demographic Characteristics of Respondents

Of 149 respondents, majority were within age group 26-34 years 92(61.7%), reside in urban areas 117(78.5%) and from South-south geopolitical zone 91(61.1%). They were mainly multiparous 65(43.6%), engaged in business/trading 48(32.2%) with tertiary level of education 100(67.1%), Table 1.

Table 1. Socio-demographic characteristics of respondents

Variables	Frequency, n=149 (%)
Age group (years)	
17-25	12 (8.1)
26-34	92 (61.7)
≥ 35	45 (30.2)
Place of residence	
Urban	117 (78.5)
Rural	32 (21.5)
Geo-political zone	
South-south	91 (61.1)
South-east	53 (35.6)
South-west	2 (1.3)
North	3 (2.0)
Parity	
Nulliparous	39 (26.2)
Primiparous	45 (30.2)
Multiparous	65 (43.6)
Mother's occupation	
Civil servant	14 (9.4)
Public servant	34 (22.8)
Professional	6 (4.0)
Artisan	19 (12.8)
Business/Trading	48 (32.2)
Unemployed/Housewives	28 (18.8)
Mother's level of education	
No formal education	2 (1.4)
Primary	6 (4.0)
Secondary	41 (27.5)
Tertiary	100 (67.1)

3.2 Knowledge of Neonatal Jaundice among Pregnant Mothers

Majority had heard of NNJ 82(55.0%), correctly defined NNJ 77(51.7%) but only 16(10.7%) could mention at least one correct cause of NNJ. Six (4.0%) pregnant women had knowledge of the area of the body jaundice appeared first in babies while 43(28.9%) and 8(5.4%) knew the complication(s) and danger(s) signs of NNJ respectively. Majority knew NNJ is treatable 121(87.7%) and knew appropriate treatment for jaundice 95(63.8%). Ninety-five (63.8%) pregnant women knew NNJ was preventable, but none knew any preventive method. Only 1(0.7%) respondent was assessed to have good knowledge of NNJ while 34(22.8%) and 114(76.5%) had fair and poor knowledge of NNJ respectively, Table 2.

3.3 Sources of Information on Neonatal Jaundice

Of the 82 pregnant women who had heard of NNJ, the commonest source of information on NNJ was from the hospital 70(52.2%) followed by from neighbours 33(24.6%) and relatives 30(22.4%) while the least were from school, market and self-knowledge grouped as others 3(2.2%) with equal distribution, Fig. 1.

3.4 Attitudes of Pregnant Mothers towards Neonatal Jaundice

Majority of respondents would visit the hospital once jaundice was observed in their babies 132(88.6%) whereas most respondents who would not visit the hospital preferred exposure to early morning sun 8 (44.4%). One hundred and

thirty-one respondents (87.9%) would consent to blood transfusion as a form of treatment if the need arose. For those who would not consent to blood transfusion, no reason was given by

majority of the respondents 9(50.0%). One hundred and twenty (80.5%) respondents had good attitude towards NNJ and 29(19.5%) had poor attitude, Table 3.

Table 2. Knowledge of neonatal jaundice among pregnant mothers

Variables	Frequency, n=149 (%)
Ever heard of NNJ	
Yes	82 (55.0)
No	67 (45.0)
Definition of NNJ	
Correct answer	77 (51.7)
Wrong answer/No idea	72 (48.3)
Cause(s) of NNJ	
Mentioned at least one correct answer	16 (10.7)
Wrong answer/No idea	133 (89.3)
Area of body jaundice appears 1st	
Correct answer	6 (4.0)
Wrong answer/No idea	143 (96.0)
Complication(s) of NNJ	
Correct answer	43 (28.9)
Wrong answer/No idea	106 (71.1)
Danger sign(s) of NNJ	
Correct answer	8 (5.4)
Wrong answer/No idea	141 (94.6)
NNJ being treatable	
Yes	121 (81.2)
No	28 (18.8)
Knows appropriate treatment for jaundice	
Yes	95 (63.8)
No	54 (36.2)
Knows NNJ is preventable	
Yes	95 (63.8)
No	54 (36.2)
Knows at least one prevention method	
Yes	0
No	149 (100.0)

Table 3. Attitudes of pregnant mothers towards neonatal jaundice

Variables	Frequency, n=149 (%)
Would visit the hospital once NNJ is noticed	
Yes	132 (88.6)
No	17 (11.4)
If No, would prefer (multiple answers), n=18	
Early morning sun	8 (44.4)
Glucose water	3 (16.7)
Abidec drop	2 (11.1)
No specific answer	5 (27.8)
Would consent to blood transfusion if need arises	
Yes	131 (87.9)
No	18 (12.1)
If No, why	
Religion	5 (27.8)
Not comfortable with blood transfusion	4 (22.2)
No reason given	9 (50.0)




Fig. 1. Sources of information on neonatal jaundice

3.5 Practice of Pregnant Mothers towards Neonatal Jaundice

Most of the respondents had no child(ren) with NNJ 124(83.2%). Of the 25(16.8%) respondents whose child(ren) had NNJ, majority took their child(ren) with NNJ to the hospital 20(80.0%). For those who did not visit the hospital with their

jaundiced baby, majority preferred mainly the exposure of their child(ren) to early morning sun 4(40.0%). Most of the respondents presented their babies to the hospital within 24 hours after jaundice was noticed 12(60.0%) with most of them being alive 24(96.0%). Twelve (8.1%) respondents had good practice while 137(91.9%) had poor practice, Table 4.

Table 4. Practice of pregnant mothers towards neonatal jaundice

Variables	Frequency, n=149 (%)
Child(ren) ever had NNJ	
Yes	25 (16.8)
No	124 (83.2)
Took child to the hospital for treatment	
Yes	20 (80.0)
No	5 (20.0)
If No, mother's preference (multiple answers), n=10	
Early morning sun	4 (40.0)
Glucose water	2 (20.0)
Ampiclox drop	2 (20.0)
Abidec drop	2 (20.0)
Time of presentation after jaundice was noticed (hours)	
< 24	12 (60.0)
24-48	3 (15.0)
> 48	5 (25.0)
Outcome of child(ren) with NNJ	
Alive	24 (96.0)
Died	1 (4.0)

3.6 Association between Sociodemographic Characteristics and Knowledge of NNJ

There was no statistical significance in the association between respondent's age, place of residence, geo-political zone, parity, occupation and level of education and the level of knowledge of the respondents (P value $> .05$), Table 5.

3.7 Relationship between Knowledge, Attitude and Practice Level of NNJ among Respondents

There was statistically significant association between the knowledge level and attitude level (P value = .003) as well as the knowledge level and practice level (P value = .001), Table 6.

Table 5. Association between sociodemographic characteristics and knowledge of NNJ

Variables	Knowledge level		P value
	Good/Fair, n=35 (%)	Poor, n=114 (%)	
Age group (years)			
17-25	2 (5.7)	10 (8.8)	.868
29-33	23 (65.7)	44 (60.7)	
≥ 35	10 (28.6)	35 (30.7)	
Place of residence			
Urban	29 (82.9)	88 (77.2)	.639
Rural	6 (17.1)	26 (22.8)	
Geo-political zone			
South-south	23 (65.7)	68 (59.6)	.806
South-east	11 (31.4)	42 (36.8)	
South-west	0	2 (1.8)	
North	1 (2.9)	2 (1.8)	
Parity			
Nulliparous	5 (14.3)	34 (29.8)	.186
Primiparous	12 (34.3)	33 (28.9)	
Multiparous	18 (51.4)	47 (41.2)	
Mother's occupation			
Civil servants	2 (5.7)	12 (10.5)	.326
Public servants	11 (31.4)	23 (20.2)	
Professionals	3 (8.6)	3 (2.6)	
Artisan	4 (11.4)	15 (13.2)	
Business/Trading	8 (22.9)	40 (35.1)	
Unemployed/Housewives	7 (20.0)	21 (18.4)	
Mother's level of education			
No formal education	0	2 (1.8)	.837
Primary	1 (2.9)	5 (4.4)	
Secondary	8 (22.9)	33 (28.9)	
Tertiary	26 (74.3)	74 (64.9)	

Table 6. Relationship between knowledge, attitude and practice level of NNJ among respondents

Variables	Knowledge level		P value
	Good/Fair, n=35 (%)	Poor, n=114 (%)	
Attitude level			
Good	34 (97.1)	86 (75.4)	.003 OR(95% CI):8.2(1.173-57.567)
Poor	1 (2.9)	28 (24.6)	
Practice level			
Good	8 (22.9)	4 (3.5)	.001 OR(95% CI):3.4(2.004-5.711)
Poor	27 (77.1)	110 (96.5)	

4. DISCUSSION

A little over half of the pregnant women in our study had heard of NNJ and could correctly define it. This is less than the finding by Egube et al. [17] in a study carried out in Benin-city and the study by Eneh et al. [18] in a previous study in Port Harcourt but similar though slightly higher than the findings of Salia et al in Ghana [19]. Though the age group of most of the caregivers in all four studies was similar (25-35 years), the study by Egube et al. [17] in Benin had a much higher sample size (389 respondents) when compared to that of the present study and this may account for the difference reported. The study by Eneh et al. [18] also had a higher sample size of 255 respondents. The study by Salia et al. [19] had caregivers with similar demographics (age-group, marital status and level of education) but was carried out in Ghana.

Although majority of the pregnant women in our study knew what NNJ was and could correctly define it only 28.9% recognised the complications of NNJ and only 5.4% of the caregivers knew any danger sign. This was similar to the finding in an earlier study in Port Harcourt by Eneh et al. [18] who documented only 22% being able to recognise brain damage as a possible complication of NNJ. This is similar to that reported by the study in Benin [17] where majority of the mothers did not know any danger sign and could not identify any complications of NNJ. This poses a very serious challenge as the caregivers even when they recognise the presence of NNJ, may not present early enough to the hospital for appropriate care as they may not recognise the danger signs even when present.

Majority of the pregnant women in our study (87.7%) knew NNJ was treatable and up to 63.8% of them knew appropriate treatment measures. This was unlike the finding reported earlier by Eneh et al. [18] where up to 50.7% of mothers did not know appropriate treatment measures (33% of the mothers recommended exposure to sunlight for treatment and 26.7% recommended glucose water for treatment). This difference in knowledge base could be attributed to the fact that only 54.2% of the mothers in the study by Eneh et al. [18] had tertiary education unlike the 67.1% of the pregnant women in our study who had tertiary education. A study by Aggarwal et al. [20] in India similarly reported only 15% of mothers knew appropriate treatment measures for NNJ with up to 20% of the mothers

recommending the use of incantations, wrapping of black thread around the cord and exposure to sunlight for treatment. This is not surprising as about 15% of the mothers in the latter study had no formal education and only 39% had tertiary education. Cultural beliefs, traditional infant care practices and reduced knowledge could be factors possibly responsible for this observation.

Though most of the pregnant women in our study (63.8%) knew NNJ was preventable none of them knew any preventive measures. This is like the findings of other authors [17,18,19,20,21]. This buttresses the fact that maternal education about NNJ, its causative factors, treatment modalities and preventive measures are highly required and advocated for to reduce neonatal morbidity and mortality.

The commonest source of information on NNJ as reported by our study was from the hospital (52.2%) then neighbours and then relatives. Similarly, the study in Abia state by Onyearugha et al. [21] reported 50% of caregivers had health workers as their source of information. Eneh et al. [18] reported a slightly lower figure of 44.7% of mother's having their source of information as health care workers. With the current trend of the increasing use of social media as a source of information dissemination, the health sector would have to engage more in this regard ie using social media as an effective means of dissemination of information concerning NNJ, its' risk factors, preventive and treatment measures.

Most of the respondents in our study would take their babies to the hospital as soon as jaundice was noticed and would consent to blood transfusion if indicated. This is unlike that reported in India [20] where only 20% of mothers would take their affected infant to the hospital. This could also be attributable to the fact that pregnant women in the present study were more educated than in the latter study (most of the mothers in our study had tertiary education whereas only 39% of the mothers in the study in India [20] had tertiary education and 15% of mothers were completely illiterate). It is worthy of note that 25% of pregnant women in our study who would not consent to blood transfusion attributed this to their religious beliefs. Severe NNJ is known to cause neurologic complications including seizure disorders, hearing deficit and cerebral palsy [22]. It has been reported as the causative factor in as many as 1/3rd of all cerebral palsy patients in a paediatric neurology clinic [23]. A previous study carried out in our

centre also showed NNJ accounting for the 3rd commonest predisposing factor of neurologic disorders after perinatal asphyxia and meningitis [24]. Some centres have reported as high as 17-22% of babies with severe NNJ receiving exchange blood transfusions as urgent treatment to rapidly lower their bilirubin levels [25,26]. Thus, the need to engage religious and cultural leaders to serve as advocates in the fight against NNJ to reduce infant morbidity and mortality cannot be overemphasized.

Majority of mothers whose babies had NNJ took their babies within 24 hours to the hospital for treatment. Those who did not, mostly used early morning sun, Ampiclox and Abidec drops as well as glucose water for supposed treatment. Aggarwal et al. [20] in contrast reported only 20% of affected babies were taken to the hospital by their mothers within 24 hours. Others used incantations, herbs, application of special black thread as well as early morning sun as supposed therapy. Again, the effects of poor education and cultural beliefs are implicated as being responsible for this difference.

Our study showed majority (76.5%) of mothers had poor knowledge, about 22.8% had fair knowledge and only 1(0.7%) mother had good knowledge. This observation was somewhat surprising as majority of the mothers in our study had tertiary education. This therefore calls for more education on neonatal jaundice and other health issues not only during the antenatal clinic health talk sessions but also via the media like television and radio as well as via social media. Our study showed majority of mothers had poor practice but good attitude with regards to NNJ. This observation was not surprising as poor knowledge influences one's practice. The fact that majority had good attitude would suggest that with increase in dissemination of appropriate evidence-based information about NNJ, it's preventive measures and management, pregnant women at our centre could have improved/better practice subsequently. Our finding was similar to that reported by Aggarwal et al. [20] in India where majority of the mothers had poor knowledge, poor attitude and in addition, poor practice. Again, poor education and sociocultural factors may be responsible for this difference. In Nigeria, Ogunfowora et al. [2] reported poor knowledge and practice to NNJ while Igboanusi et al. [14] documented poor knowledge in 91% and poor attitude in 62% of respondents. In contrast, Salia et al. [19] documented slightly less than half of caregivers demonstrated good knowledge (45.5%) and attitude (47.5%) while

58.9% had good practices regarding NNJ. This difference could be attributed to difference in geographic location and possible difference in the perception of NNJ. The level of education of the respondents was not documented in the latter study although in the present study, more than 2/3rd had tertiary education.

Our study found no significant association between Knowledge and sociodemographic factors. However, there was significant association between Knowledge and mother's attitude (P value=.003) and practice (P value=.001). Similarly, Ogunlesi et al [27] found no significant association of knowledge with religion and marital status but found significant association between knowledge and practice as observed in the present study. They reported that mothers with good knowledge had better care-seeking behaviour [27]. Salia et al. [19] in Ghana reported caregivers who had prior awareness and received education on NNJ were 3 times more likely to have good knowledge than those without previous education. Thus, efforts to promote well informed and improved caregivers' attitude will improve maternal health-seeking behaviour and reduce disabilities and death through early detection and intervention of infants with NNJ [19,20,21,25,27].

5. CONCLUSION

Pregnant women in Rivers State University Teaching Hospital had insufficient knowledge and poor practice towards NNJ but had good attitude towards NNJ. The need to intensify health education in pregnant women on neonatal jaundice and other health related issues cannot be over emphasized as this will improve their attitude towards neonatal jaundice as well as their practice.

CONSENT

As per international standard or university standard, patients' written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

ACKNOWLEDGEMENT

We acknowledge the nurses in the antenatal clinic of the Rivers State University Teaching

Hospital, the research assistants and the non-clinical staff. We also appreciate the pregnant women who gave consent to participate in the study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Ogunfowora OB, Daniel OJ. Neonatal jaundice and its management: Knowledge, attitude and practice of community health workers in Nigeria. *BMC Public Health*. 2006;6:19.
- Ogunfowora OB, Adefuye PO, Fetuga MB. What do expectant mothers know about neonatal jaundice? *Int Electronic J Health Education*. 2006;9:134-140.
- Waqar T, Arham QA, Razzaq A. Neonatal jaundice: A review and guidelines for its management in resource limited countries. Available: <https://www.researchgate.net/publication/259309673>
- Melton K, Akinbi HY. Neonatal jaundice. Strategies to reduce bilirubin induced complications. *Postgrad Med*. 1999; 106:167-178.
- Olusanya BO, Osibanjo FB, Mabogunje CA, Slusher TM, Olowe SA. The burden and management of neonatal jaundice in Nigeria: A scoping review of the literature. *Niger J Clin Pract*. 2016;19:1-17.
- Young Infants Clinical Signs Study Group. Clinical signs that predict severe illness in children under age 2 months: A multicentre study. *Lancet*. 2008;371:135-142.
- West BA, Josiah AE. Neonatal jaundice: Prevalence, risk factors and outcome among inborn babies in a neonatal unit in Port Harcourt, Nigeria. *Indian J Basic Applied Med Res*. 2020;9:92-103. DOI: 10.36848/IJMAMR/2020/18215.55572
- American Academy of Pediatrics. Provisional committee for quality improvement and subcommittee on hyperbilirubinaemia Practice parameter: Management of hyperbilirubinaemia in the healthy term newborn. *Pediatrics*. 1994;94:558-562.
- Porter MI, Dennis BL. Hyperbilirubinaemia in the term newborn. *Am Family Phys*. 2002;65:599-607.
- Zupan J. Prenatal mortality in developing countries. *N Engl J Med*. 2005;352:2047-2048.
- Ezechukwu IC, Ugochukwu EF, Egbuonu I, Chukwuka JO. Risk factors from neonatal mortality in regional tertiary hospital in Nigeria. *Niger J Clin Pract*. 2004;7:50-52.
- Watchko JK. Kernicterus and the molecular mechanisms of bilirubin-induced central nervous systems injury in newborns. *Neuromuscular Med*. 2006;8:513-529.
- Scrafford CG, Mullany LC, Khatry SK, Leclerg SC, Darmstadt GL, Tielsch JM. Incidence and risk factors for neonatal jaundice among newborns in Southern Nepal. *Trop Med & Inter Health*. 2013;18:1317-1328.
- Igboanusi CJ, Nmadu AG, Joshua IA, Onoja-Alexander MO, Olatubosun B. Knowledge and attitude on neonatal jaundice among women of reproductive age group in rural community in Northern Nigeria. *Sahel Med J*. 2020;23:184-190
- Khalesi N, Rakhshani F. Knowledge, attitude and behaviour of mothers on behaviour of mothers on neonatal jaundice. *J Pak Med Med Assoc*. 2008;58:671-674.
- Amirshaghagi A, Ghabili K, Shoja MM, Kooshaver H. Neonatal jaundice: Knowledge and practice of Iranian mothers with icteric newborns. *Pak J Biol Sci*. 2008;11:942-945.
- Egube BA, Ofili A, Isara A, Onakewhor JU. Neonatal jaundice and its' management: Knowledge, attitude and practice among expectant mothers attending antenatal care at University of Benin Teaching Hospital, Benin City, Nigeria. *Nig J of Clin Pract*. 2013;16(2):199-94. DOI: 10.4103/1119-3077.110147
- Eneh AU, Ugwu RO. Perception of neonatal jaundice among women attending children out patient and immunization clinics of the University of Port Harcourt Teaching Hospital, Post Harcourt. *Niger J Clin Pract*. 2009;12(2):187-191.
- Salia SM, Afaya A, Wuni A, Ayanore MA, Salia E, Kporvi DD et al. Knowledge, Attitudes and practices regarding neonatal jaundice among caregivers in a tertiary health facility in Ghana. *PLoS One*. 2021;16(6):e0251846. DOI: 10.1371/journal.pone.0251846.r008
- Aggarwal B, Agrawal A, Chaudhary P, Gupta G, Rana S, Gupta S. Neonatal

- jaundice: Knowledge, attitude, beliefs and practices of postnatal mothers in a tertiary care hospital in Uttarakhand, India. Indian J Child Health. 2017;4(4):603-608.
21. Onyearugha CN, Chapp-Jumbo A, George IO. Neonatal Jaundice: Evaluating the knowledge and practice of expectant mothers in Aba, Nigeria. Journal of health Science Research. 2016;1(2):42-7. ISSN 2456-2688. DOI: 10.18311/jhsr/2016/v1/i2/4918
 22. Ibe BC. Neonatal Jaundice. In: Azubuike JC, Nkangineme KE, editors. Paediatrics and child health in a tropical region. Owerri: African Educational Services. 1999;204-211.
 23. Asindi AA. The pattern of neurological disabilities in children seen at the University of Calabar Teaching Hospital. Nig J Paediatr. 1986;13:127-131.
 24. West BA, Onubogu UC. Pattern and predisposing factors of neurological disorders in a neurology outpatient clinic in Port Harcourt, Nigeria. Med Res Chronicles. 2019;6:218-228.
 25. Ibekwe RC, Ibekwe MU, Muoneke VU. Outcome of Exchange Blood transfusions done for Neonatal Jaundice in Abakaliki, South Eastern Nigeria. J Clin Neonatol. 2012;1(1):34-37. DOI: 10.4103/2249-4847.92239
 26. Sgro M, Campbell D, Shah V. Incidence and causes of severe neonatal hyperbilirubinaemia in Canada. CMAJ. 2006;175:587-590.
 27. Ogunlesi TA, Abdul AR. Maternal knowledge and care-seeking behaviours for newborn jaundice in Sagamu, Southwest Nigeria. Niger J Clin Pract. 2015;18(1):33-40. DOI: 10.4103/1119-3077.146976

© 2021 West et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/83226>