



Onchocercal Skin Disease Profiles Following Ivermectin Treatment in the Middle Imo River Basin, Nigeria

A. A. Amaechi ^{a*}, J. I. Iwunze ^a, C. M. Nwokeji ^b and J. Egejuru ^c

^a Entomology and Parasitology Unit, Department of Zoology, Imo State University Owerri, Nigeria.

^b Department of Medical Parasitology, Imo state University Teaching Hospital Orlu, Nigeria.

^c Out Patient Department (OPD), Federal Medical Center Owerri, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. Authors AAA and NCM conceived and designed the study. Authors AAA, IJI and EJ collected the data in the field. Authors EJ and IJI analyzed the data. Author AAA wrote the first manuscript. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJTDH/2021/v42i2330561

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/64339>

Original Research Article

Received 07 November 2020

Accepted 18 January 2021

Published 27 December 2021

ABSTRACT

In Nigeria infection with onchocerciasis has been under control with the introduction of ivermectin in most affected communities for a decade. To assist 'stop mass drug administration (MDA)' decision, a cross-sectional survey evaluated the impact of ivermectin on onchocercal Skin Diseases (manifestations) from 540 participants in 5 endemic communities. The rapid epidemiological assessment (REA) method with palpable nodules and skin depigmentation (leopard skin, LS) as diagnostic indices were used. The prevalence profiles of palpable nodules and leopard skin was 5.9% and 12.2% respectively. Prevalence of clinical features varied among the farming community ($P < 0.05$). Similarly, age group 21-30 years had significant prevalence of clinical features than other age groups ($P < 0.05$). Males had significantly higher prevalence of these indices (palpable nodules 6.8%; LS 13.3%) than females (palpable nodules 4.6%; LS 11.1%). Although, Onchocercal Skin Diseases (clinical manifestations) were observed in all age groups, it occurred most in older groups. These groups with nodules could portend as reservoir for transmission. The implications of the results were discussed in the context of the on-going prediction of possible elimination of onchocerciasis in Africa and 'stop MDA' decision.

*Corresponding author: Email: amaechiaustine@gmail.com;

Keywords: Ivermectin treatment; diagnostic indices; onchocercal skin disease.

1. INTRODUCTION

Onchocerciasis is a leading cause of visual impairment and blindness in many developing countries. The main complications are severe eye disease characterized by popular or hypopigmented lesions and intense itching. The disease is caused by the filarial nematode *Onchocerca volvulus* that is transmitted by *Simulium damnosum* species of black flies [1]. According to WHO [2], the disease affected about 17 to 18 million people in 37 countries of the world with approximately 123 million being at risk of infection. The disease is widespread in Nigeria and causes blindness in most rural communities. Of all the countries of the world, Nigeria has the largest number of persons with onchocerciasis accounting for about a third of the global prevalence with about 40 million at risk of infection [3].

Onchocerciasis in Nigeria is solely transmitted by members of *S. damnosum* complex which is widespread in the Savannah forest, Savannah mosaic and rainforest areas. They are known to be common in the rocky sections of the River Niger and its tributaries [4]. In Imo State rapid epidemiological mapping of onchocerciasis (REMO) study in sentinel villages (Ezinachi, Umulolo, Umuele, Ndioji and Isiokwe) recorded a palpable nodule range of 40-56% and 24-48% of leopard skin (LS). This lead to Imo State being classified as an endemic State and necessitating community- wide ivermectin treatment [5-6], study showed that blindness associated with onchocerciasis is fewer in the rainforest zone compared to the Savannah zone. The pattern is less clear in central Africa with severe blinding as well as less blinding onchocerciasis presents in both Savannah and rainforest belts. Onchocercal skin diseases (OSD) is found in East Africa where blindness is rare [1]. Skin disease and severe prurities (troublesome itching) has recently been recognized to be affecting more than 50% of the populations in some rural communities in the rainforest belt with relative low blindness prevalence [7].

The clinical presentation of the disease is varied from changes in the skin (dermatitis), lymphatic system and ocular (eye) changes leading to blindness. The symptoms include skeletal pain, prurities, skin lesions, impaired vision (patial or total blindness) and premature death. The infective microfilariae is introduced by the bite of

black flies which latter form swellings called nodules. These nodules vary in sizes as they contain adult worms. In patients in Africa, nodules tend to be located on the lower tonso, pelvis and lower extremities as against patients in Central and South America (Aapredbook Publications 2009). The relationship between the sites of biting by the vector and the localization of the nodules showed that many nodules were found on the lower parts of the body whereas in Central America as many as 70% were found on the head [8]. The localization of the nodules has been related to the area of bitten by different vector species [9]. In an effort to know whether onchocerciasis transmission had been halted / interrupted since on-going treatment characterized by varied compliance, this study evaluated nodules and dermatitis as signs of onchocerciasis in the study area.

2. MATERIALS AND METHODS

2.1 Study Area and Population

The study area is Okigwe Local Government Area of Imo State Nigeria. Okigwe is located in the Northern part and is one of the 27 LGAs of the State. There are 115 villages in Okigwe, all has been receiving mectizan/ ivermectin since 1995 [10]. The communities where the study was carried out were all in the Imo River Basin of Nigeria located between Lat. 5°40'1-5°57'1N and Long. 7°10'1-7°26'1E. They are drained by several fast flowing streams and rivulets which empty into the main river in the area, the Imo River [11]. The seasons are well defined: a wet season from October to March and a dry season from April to September. The range of annual rainfall is between 1700 and 2000 mm with average relative humidity of about 80% and mean annual temperature of 29°C [12].The vegetation is forest. Subsistence farming, fishing, hunting, wine tapping and petty trading are the main occupation of the people who are Ibo ethnic group.

2.2 Subject Examination

Five farming communities; Amano, Amuro, Ihube, Umulolo and Aku along the fast flowing rivers were selected for the study. They population census with the help of field assistants was conducted in each communities. Out of a combined population ,540 subjects male and females aged 5 years and above were

chosen based on household records. 150 from Amuro and Umulolo and 80 respectively from Aku, Amano and Ihube endemic communities. Subjects were asked to gather at the health center in the communities. A health talk was made to them for enlightenment on the existing health problem. Their verbal consent were obtained thereafter before they were physically examined; leopard skin, palpation for presence of onchocercal dermatitis and nodules, scrotal and clitoral enlargements, hanging groin, itching and virtual acuity by the optometrist in the team. Subjects' records on history and use of ivermectin as well as demographic data (age, sex, occupation and marital status) were taken.

2.3 Data Analysis

Simple percentages were used to estimate the prevalence rate of various parameters by the subjects. Chi-Square was used to ascertain significance in sex and age on the prevalence of clinical manifestations among subjects at 0.05 level of probability.

3. RESULTS

A total of 5 farming communities were surveyed between May and November, 2019. The highest

infection rate was found in Umulolo (15.4%) followed by Amuro (13.1%), Aku, Ihube and Amano had 9.3%, 8.5% and 8.3% respectively. The overall prevalence was 54.6% (Table 1). Prevalence of clinical features varied among the farming community ($P < 0.05$). Results of onchocerciasis manifestations revealed that leopard skin was the commonest (12.2%) feature. Lizard skin (11.3%), onchocercal nodules (5.9%), itching (8.0%), musculoskeletal pains (10.0%) and poor vision (3.5%) were also reported. Leopard skin was highest in Amuro (21.3%) and Ihube (18.8%) and lowest in Amano (7.3%). The prevalence of onchocercal nodules was highest in Amuro (8.7%) and lowest in Ihube (2.5%). Onchodermatitis was highest in Ihube (5.0%) and lowest in Aku, Amano and Amuro (0.0%). Poor vision was highest in Aku (7.3%) and lowest in Amuro (0.0%). No scrotal and clitoral enlargements were recorded. Prevalence of infections increases with age group (Table 2). Males had significantly higher prevalence than females ($p < 0.05$; Fig. 1). Similarly, age group 21-30 years had significant prevalence of Clinical Features than other age group ($P < 0.05$). Leopard skin, lizard skin, onchodermatitis, poor vision, itching and musculoskeletal pains were noted from 21-30 years old. Hanging groin was found in 31 years and above (Table 3).




Plate 1. Showing subjects with Onchocercal nodules

Table 1. Results of Onchocercal clinical features of the subjects

Clinical features	Amano	Amuro	Umulolo	Aku	Ihube	Total
Musculo-skeletal pain	09(11.3)	11(7.3)	13(8.7)	13(8.7)	08(10.0)	54(10.0)
Poor vision	01(1.3)	00(0.0)	07(2.0)	06(7.5)	05(6.3)	19(3.5)
Onchocercal nodules	04(5.0)	13(8.7)	08(5.3)	05(6.3)	02(2.5)	32(5.9)
Leopard skin	17(21.3)	11(7.3)	14(9.3)	09(11.3)	15(18.8)	66(12.2)
Lizard skin	07(8.8)	17(11.3)	23(15.3)	11(13.8)	03(3.8)	61(11.3)
Itching	07(8.8)	16(10.7)	13(8.7)	03(3.8)	04(5.0)	43(8.0)
Hanging groin	00(0.0)	03(2.0)	00(0.0)	03(3.8)	05(6.3)	11(2.0)
Scrotal/Clitoral Elanrgement	00(0.0)	00(0.0)	00(0.0)	00(0.0)	00(0.0)	00(0.0)
Rashes and Onchodermatitis	00(0.0)	00(0.0)	05(3.3)	00(0.0)	04(5.0)	09(1.7)
Total	45(8.3)	71(13.1)	83(15.4)	50(9.3)	46(8.5)	295(54.6)

Table 2. Age-related Distribution of Onchocerciasis among the subjects

Communities	5-10		11-20		21-30		31-40		41-50		51+	
	No Exam	No(%) +ve	No Exam	No(%) +ve	No Exam	No(%) +ve	No Exam	No(%) +ve	No Exam	No(%) +ve	No Exam	No(%) +ve
Amano	7	1(5.9)	7	0(0.0)	17	4(23.5)	15	3(20.0)	18	3(17.7)	13	3(23.1)
Amuro	33	1(3.0)	13	2(15.4)	25	3(12.0)	23	3(13.0)	31	2(6.5)	25	3(12.0)
Umulolo	26	1(3.8)	17	2(11.8)	31	3(9.7)	24	2(8.3)	23	3(13.0)	29	5(17.2)
Aku	13	0(0.0)	4	1(25.0)	13	3(23.1)	19	2(10.5)	16	2(12.5)	15	2(13.3)
Ihube	15	0(0.0)	3	1(33.0)	13	4(30.8)	19	3(15.8)	23	5(21.7)	10	2(20.0)
Total	94	3(3.2)	44	6(13.6)	99	17(17.2)	100	13(13.0)	111	15(13.5)	92	15(16.3)

Table 3. Results of Onchocercal clinical features by Age and Gender

Age groups (Years)	Sex	No Exam	Musclo-skeletal pain	Poor vision	Nodules	Leopard skin	Lizard skin	Hanging groin	Rashes and Onchodermatitis	Itching	Scrotal/Clitoral enlargement	Total
05-10	M	61	0	0	0	0	0	0	0	0	0	0
	F	43	0	0	0	0	0	0	0	0	0	0
11-20	M	21	0	0	0	0	0	0	0	0	0	0
	F	13	0	0	0	0	0	0	0	0	0	0
21-30	M	51	3	2	0	4	3	0	2	0	0	14
	F	48	2	0	0	5	0	0	1	4	0	12
31-40	M	47	7	4	3	7	5	1	1	7	0	35
	F	53	3	2	3	3	7	0	0	5	0	23
41-50	M	52	11	3	5	13	7	4	1	3	0	47
	F	59	5	1	3	7	12	1	1	7	0	37
51+	M	47	14	3	11	13	8	3	2	10	0	64
	F	45	9	4	7	14	19	2	1	7	0	63
Total	M	279	35(12.5)	12(4.3)	19(6.8)	37(13.3)	29(10.4)	8(2.9)	6(2.2)	20(7.0)	0(0.0)	160(29.6)
	F	261	19(7.3)	7(2.7)	13(4.6)	29(11.1)	32(12.3)	3(1.1)	3(1.1)	23(8.8)	0(0.0)	135(25.0)
Grand total		540	54(10.0)	19(3.5)	32(5.9)	66(12.2)	61(11.3)	11(2.0)	9(1.7)	43(8.0)	0(0.0)	295(54.6)




Fig. 1. Prevalence by gender across all study subjects

In occupation prevalence, farmers (19.5%) had the highest prevalence followed by others (teachers and applicants) 12.2%, Traders and

school children had 10.8% and 4.1% respectively (Fig. 2). Plates 1 and 2 showed subjects with onchocercal nodules and leopard skin.




Plate 2. Showing subjects with Leopard skin




Fig. 2. Prevalence of Onchocerciasis with relation to occupation in the study area

4. DISCUSSION AND CONCLUSION

The advent of ivermectin for the treatment of onchocerciasis and its sustainability for large scale application has been a major breakthrough against onchocercal skin diseases. The severity of these morbidity indicators especially palpable nodules and dermatitis are used to assess clinical improvement of ivermectin intervention. The results of this study have shown that the epidemiological profile has changed to hypo endemicity status. The five communities surveyed in the middle Imo River Basin had prevalence with one or two clinical features. Umulolo and Amuro communities located close to River Lolo and Ibii had highest manifestations (15.4% and 13.1%) while other farming communities (Ihube, Aku and Amano) which resides >5KM away had varied prevalences. These Rivers provide adequate breeding sites for black flies that have the capacity to bite 60-100 KM from these sites or windblown 300-500 KM away [13]. Lolo and Ibii Rivers are tributaries of Imo River which serve these communities and may account for the prevalence [14]. The varied percentage reduction in these communities may be due to the difference in treatment coverage pattern in the River Basin. The relationship between ivermectin community coverage and community morbidity reduction was evidence in rapid epidemiological assessment (REA) results. Remarkable reduction on the common onchocerciasis symptoms were identified; nodules (5.9%), poor/impaired vision (3.5%), itching (8.0%), leopard skin (12.2%) among others. Similar to previous studies [15], these indicators (nodule and leopard skin) showed that prevalence increased with age and males had higher prevalence than females. Nodular rates in the five communities ranged from 2.5% in Ihube to 8.7% in Amuro with overall rate of 5.9%. Leopard skin ranged from 7.3% in Amuro to 21.3% in Amano with overall rate of 12.2%. The nodule rate shows a decrease from the >20% nodules rate specified for definite CDTI area [16], which indicated that annual ivermectin treatment is effective. The reported claim by a participant that his nodule size has reduced after several annual treatments provides support to the questionnaire and clinical studies elsewhere [17,13]. The observation on leopard skin among older age (>20 years) and the absence among younger age group (< 20 years) support the popular consensus that the symptom is a long standing manifestation induced by degenerating microfilariae, a very important diagnostic tool in the communities [18].

The continued bulk of infection on the middle and elderly people remains significant and singled out these working groups (most active and productivity farming age) as the most at risk. Occupational (farming, fishing, animal rearing /hunting etc) or residential exposure could be attributed to this observation. The infection reported among age group 5-10 (children) may be due to early contact with infective black flies bites by their visits to breeding sites or learning under trees inside school premises which is a common habit among the schools especially during hot weather. Three factors; length of time spent near river or work (women hardly exceed 3:00 pm while men stays till 6:00pm), culture which permit men to fend for their families while women give helping hands (men are full time farmers, fishermen, hunters etc while women combine farming with trading or other indoor activities) and dressing pattern (women are traditionally attired in dressing that cover them than men). These could explain the observed difference between males and females prevalence rates.

[19] had posited that the efficacy of ivermectin in controlling onchocerciasis burden of any community depends on the consistency of the treatment regimen. Somewhere in Ghana [20], demonstrated that skin microfilaria repopulation commenced in 12-90% of patients studied in various communities. The absence of complete clearance of onchocercal skin disease may be linked to intervals between treatment beyond 90 days which may allow the effective transmission of the parasite by the black fly vectors. These study areas also shared common boundaries with endemic areas of Abia State, Hence an obstacle to 'stop MDA' decision. We argue that associated operations research should be undertaken to allow data from the same samples and age groups to help stop MDA by technical committees.

CONSENT

It is not applicable.

ETHICAL APPROVAL

The State Ministry of Health, Local Government Area Health Unit and the village heads were notified before the study commenced through a letter of intent from the Research Team under the approval of the Head of Department of Zoology, Imo State University, Owerri Nigeria. The purpose of the study was explained in each

village to the village heads and the elders, and their consent was obtained for the mobilization by the respective communities.

ACKNOWLEDGEMENTS

This study received the financial support of the TETFUND Institutional Based Research project intervention for Research in Tropical Disease. We thank the entire staff of Zoology Department Imo State University Owerri, Nigeria for logistic support and assistance in the field. The authors highly appreciate our Post graduate students for useful in puts.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. WHO. Onchocerciasis and its control. Report of a WHO Expert Committee on onchocerciasis control. World Health Organisation Technical Report. 1995;852;1-104.
2. WHO. The world Health Report. Conquering suffering, Enriching humanity. World Health Organisation. Geneva; 1997.
3. Nwoke BEB. The Socioeconomic aspects of human onchocerciasis in Africa: present appraisal. Journal of Hygiene Epidemiology and Microbiol Immunology. 1990;59(1):37-44
4. Eneanya CI and Nworgu OC. Social and psychological aspects of onchocercal skin disease in Nkweue-Ezunala, Anambra State, Nigeria. Nigerian Journal of Parasitology. 2001;22(1,2):11-16.
5. Nwoke BEB, Edungbola LD, Mencia BS, Njoku AJ, Abanobi OC, Nkwogu EU, et al. Human onchocerciasis in the rainforest zone of south eastern Nigeria.1: Rapid assessment for community diagnosis in the Imo River Basin, Nigeria. Nigerian Journal of Parasitology. 1994;15:46-58.
6. Dozie INS, Onwuliri COE, Iwuagwu FO. Onchocerciasis in Imo State, Nigeria. Clinical and epidemiological studies of anterior segment lesions. Nigerian Journal of Parasitology. 2003;24:89-94.
7. Kale OO. Onchocerciasis the burden of the disease. Annals of Tropical Medicine and Parasitology. 1998;92:5109-5115.
8. Adetokunbo OL, Herbert MG. Short Textbook of Public health Medicine for the Tropical. Power Book LTD, London. 2003;229.
9. Ubachukwu PO. Human onchocerciasis: epidemiological status of Uzo-Uwani local Government Area of Enugu State Nigeria. Nigerian Journal of Parasitology. 2004;25:95-102.
10. Global River Blindness Onchocerciasis. Annual Report/ sentinel village evaluation Reports. GRBP and Carter Center Inc. USA; 2000.
11. Atlas of Imo State of Nigeria. Department of Land Survey and Urban population. Ministry of works and Transport. 1st ed. Government Press Owerri; 1984.
12. Dozie INS, Onwuliri COE, Nwoke BEB, Anosike JC, Okoli NJC. Onchocerciasis in Imo State Nigeria. The prevalence and distribution in the middle Imo River Basin, Nigeria. Nigerian Journal of Parasitology. 2006;27:16-22.
13. Emukah EC, Osuala E, Miri ES, Onyenama J, Amadiogwu S, Korve K, et al. A longitudinal study of impact of repeated mass ivermectin treatment of clinical manifestation of onchocerciasis in the Imo State Nigeria. Journal of Tropical Medicine and Hygiene. 2004;70(5):556-610 Clinical manifestation.
14. Akogun OB, Onwuliri COE. Hyper endemic onchocerciasis in persons under twenty years of age in an endemic area of Northern Nigeria. Annals of Tropical Medicine and Parasitology. 1991;53:10-29.
15. Ezigbo OR, Nwoke BEB, Ukaga CN, Emukah EC. Impact assessment of repeated mass ivermectin treatment on onchocerciasis in Abia State, Nigeria. Journal of Biology, Agriculture and HealthCare. 2013;3(2).
16. Noma M, Nwoke BEB, Nutall I, Tambala PA, Enyong P, Romme J, et al. Rapid epidemiological mapping of onchocerciasis (REMO): Its application by the African Programme for Onchocerciasis Control (APOC). Annals of Tropical Medicine and Parasitology. 2002;96:29-39.
17. Ukaga CN, Dozie INS, Nwoke BEB. Validation of reports of nodule dissolution after repeated ivermectin treatment of onchocerciasis in south eastern Nigeria. East African Medical Journal. 2001;78:4-6.

18. Pajiah M and Eneanya CE Nodules and dermatitis as signs of onchocerciasis in some communities of Aniocha North Local Government Area, Delta State Nigeria. Nigerian Journal of Parasitology. 2019;40(1):97-102.
19. Winnen M, Piaisier AP, Alley ES, Nageslkeke NJD, van Qortmarsen G, Boatin, et al. Can ivermectin mass treatment eliminate onchocerciasis in Africa? Bulletin of World Health Organisation. 2002;80:384-390.
20. Osei-Atweneboana MY, Eng JKL, Boakye DA, Gyapang JO Pichard PK Prevalence and intensity of *Onchocerca volvulus* infection and efficacy of ivermectin in endemic communities in Ghana a two-phase epidemiological study. Lancet. 2007;369:2021-2029.

© 2021 Amaechi et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/64339>